





APPROVED: 11 November 2022 doi: 10.2903/j.efsa.2022.7666

# The European Union One Health 2021 Zoonoses Report

European Food Safety Authority European Centre for Disease Prevention and Control

## **Abstract**

This report of the European Food Safety Authority and the European Centre for Disease Prevention and Control presents the results of zoonoses monitoring and surveillance activities carried out in 2021 in 27 MSs, the United Kingdom (Northern Ireland) and nine non-MSs. Key statistics on zoonoses and zoonotic agents in humans, food, animals and feed are provided and interpreted historically. In 2021, the first and second most reported zoonoses in humans were campylobacteriosis and salmonellosis, respectively. Cases of campylobacteriosis and salmonellosis increased in comparison with 2020, but decreased compared with previous years, In 2021, data collection and analysis at the EU level were still impacted by the COVID-19 pandemic and the control measures adopted in the MSs, including partial or total lockdowns. Sixteen MSs and the United Kingdom (Northern Ireland) achieved all the established targets in poultry populations for reduction in Salmonella prevalence for the relevant serovars. Salmonella samples from carcases of various animal species and samples for Campylobacter quantification from broiler carcases were more frequently positive when performed by the competent authorities than when own-checks were conducted. Yersiniosis was the third most reported zoonosis in humans, followed by Shiga toxin-producing Escherichia coli (STEC) and Listeria monocytogenes infections. L. monocytogenes and West Nile virus infections were the most severe zoonotic diseases, with the most hospitalisations and highest case fatality rates. Overall, MSs reported more foodborne outbreaks and cases in 2021 than in 2020. S. Enteritidis remained the most frequently reported causative agent for foodborne outbreaks. Salmonella in 'eggs and egg products' and in 'mixed foods' were the agent/food pairs of most concern. Outbreaks linked to 'vegetables and juices and products thereof' rose considerably compared with previous years. This report also provides updates on brucellosis, Coxiella burnetii (Q fever), echinococcosis, rabies, toxoplasmosis, trichinellosis, tuberculosis due to Mycobacterium bovis or M. caprae, and tularaemia.

© 2022 European Food Safety Authority and European Centre for Disease Prevention and Control.

**Keywords:** Campylobacter, foodborne outbreaks, *Listeria*, monitoring, parasites, *Salmonella*, zoonoses

Requestor: European Commission

**Question number:** EFSA-Q-2021-00762 **Correspondence:** zoonoses@efsa.europa.eu



18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (thps://onlinelibrary.wiley.com/terms

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

**Declarations of interest:** The declarations of interest of all scientific experts active in EFSA's work are available at https://ess.efsa.europa.eu/doi/doiweb/doisearch.

**Acknowledgements:** EFSA and the ECDC wish to thank the members of the EFSA Scientific Network for Zoonoses Monitoring Data and of the ECDC Food and Waterborne Diseases and Zoonoses Network, the ECDC Emerging and Vector-borne Diseases Network and the ECDC Tuberculosis Network, who provided the data and reviewed the report; the members of the Scientific Network for Zoonoses Monitoring Data for their endorsement of this scientific report; the EFSA staff members (Frank Boelaert, Giusi Amore, Valentina Rizzi, Anca Stoicescu, Maria Teresa Da Silva Felício, Sandra Luisa De Almeida Florentino Correia, Michaela Hempen, Winy Messens, Sotiria-Eleni Antoniou, Francesca Baldinelli, Alessandro Broglia, Sofie Dhollander, Lisa Kohnle and Lina Mur), the ECDC staff members (Taina Niskanen, Joana Haussig and Veronica Cristea) and the EFSA and ECDC contractor: Istituto Superiore di Sanità (Gaia Scavia, Ilaria Altieri, Fabrizio Anniballi, Antonino Bella, Simone Cacciò, Adriano Casulli, Alessandra Ciervo, Elisabetta Delibato, Michele Luca D'Errico, Aurora Garcia Fernandez, Maria Angeles Gomez Morales, Francesca Iacoponi, Arnold Knijn, Marco Lalle, Claudia Lucarelli, Stefano Morabito, Marco Ortoffi, Paolo Pasquali, Pierfrancesco Barbariol, Flavia Riccardo, Elisabetta Suffredini, Rosangela Tozzoli, Eleonora Ventola and Laura Villa); Istituto Zooprofilattico delle Venezie (Giorgia Angeloni, Lisa Barco, Grazia Manca, Katia Capello, Giulia Cento, Laura Ciot, Paola De Benedictis, Laura Gagliazzo, Annie Ishebabi, Elena Lazzaro, Monica Lorenzetto, Marzia Mancin, Claudio Mantovani, Martina Simonato and Marica Toson); Istituto Zooprofilattico Sperimentale dell'Abruzzo e del Molise 'G. Caporale' (Paolo Calistri, Enrico De Albentiis, Giuliano Garofolo, Diana Palma, Fabrizio De Massis, Francesco Pomilio, Carla Ippoliti, Americo Bonanni, Susanna Tora, Alessio Di Lorenzo, Mauro Ferella, Andrea De Ruvo and Annamaria Conte); Istituto Zooprofilattico Sperimentale della Lombardia ed Emilia Romagna B. Ubertini – IZSLER (Maria Beatrice Boniotti, Nadia Vicari, Maria Lodovica Pacciarini, Silvia Bellini and Gabriele Casadei) for the support provided to this scientific report; the French Agency for Food, Environmental and Occupational Health & Safety (Adrien Asseré, Christophe Cordevant, Aurélie Couesnon, Radu Blaga, Florence Cliquet, Corinne Danan, Nolwenn Dheilly, Claire Ponsart, Dana Pottratz, Emmanuelle Robardet and Elodie Rousset).

**Amendment:** Few minor editorial corrections (two names added in the acknowledgments and two numbers changed in the first paragraph on page 184) were carried out that do not materially affect the contents or outcome of this scientific output. To avoid confusion, the original version of the output has been removed from the EFSA Journal, but is available on request.

**Suggested citation:** EFSA and ECDC (European Food Safety Authority and European Centre for Disease Prevention and Control), 2022. The European Union One Health 2021 Zoonoses Report. EFSA Journal 2022;20(12):7666, 273 pp. https://doi.org/10.2903/j.efsa.2022.7666

**ISSN:** 1831-4732

© 2022 European Food Safety Authority and European Centre for Disease Prevention and Control.

This is an open access article under the terms of the Creative Commons Attribution-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited and no modifications or adaptations are made.



The EFSA Journal is a publication of the European Food Safety Authority, a European agency funded by the European Union.





# **Table of contents**

Abstract		1
	tion	12
Legal ha	sis of European Union-coordinated zoonoses monitoring	12
Reportin	g requirements	12
	f Reference	12
	rces and report production	13
Data SUL	irces and report production	
Data ana	alyses and presentation	15
	y of human zoonoses data for 2021	17
	son of human zoonoses data for 2021, 2020 and previous years	20
	s included in compulsory annual monitoring (in accordance with Directive 2003/99 List A)	22
1.	Campylobacter	22
1.1.	Key facts	23
1.2.	Surveillance and monitoring of Campylobacter in the EU	23
1.2.1.	Humans	23
1.2.2.	Food and animals	24
1.3.	Data analyses	25
1.4.	Results	25
1.4.1.	Overview of key statistics, EU, 2017–2021	25
	Human campylobacteriosis	26
		29
1.4.3.	Campylobacter in food	
1.4.4.	Campylobacter in animals	34
1.5.	Discussion	35
2.	Salmonella	37
2.1.	Key facts	38
2.2.	Surveillance and monitoring of Salmonella in the EU	39
2.2.1.	Humans	39
2.2.2.	Food, animals and feed	39
2.3.	Data analyses	40
2.3.1.	Comparison between competent authority and food business operator sampling results	40
2.3.2.	Statistical trend analyses for poultry monitoring data	41
2.3.3.	Descriptive analyses of Salmonella serovars.	41
2.3.3.	Results	42
2. <del>4</del> . 2.4.1.	Overview of key statistics, EU, 2017–2021	
		42
2.4.2.	Human salmonellosis	43
2.4.3.	Salmonella in food	46
2.4.4.	Salmonella in animals	56
2.4.5.	Salmonella in feed	68
2.4.6.	Salmonella serovars in humans, food and animals	69
2.5.	Discussion	76
3.	Listeria monocytogenes	78
3.1.	Key facts	78
3.2.	Surveillance and monitoring of <i>Listeria monocytogenes</i> in the EU	79
3.2.1.	Humans	80
3.2.2.	Food, animals and feed	80
3.3.	Data analyses	81
3.3.1.	Data on RTE food in the context of Regulation (EC) No 2073/2005 on microbiological criteria	81
3.3.2.	Other monitoring data for <i>Listeria monocytogenes</i> in RTE food	82
	Monitoring data for <i>Listeria monocytogenes</i> in animals and feed	82
3.4.	Results	83
3.4.1.	Overview of key statistics, EU, 2017–2021	83
3.4.2.	Human listeriosis	84
	Listeria monocytogenes in RTE food	86
3.4.4.	Listeria spp. in animals	89
3.5.	Discussion	90
4.	Shiga toxin-producing Escherichia coli (STEC)	92
4.1.	Key facts	93
4.2.	Surveillance and monitoring of STEC infections in the EU	93
4.2.1.	Humans	93
4.2.2.	Food and animals	94
4.3.	Data analyses	94
4.3.1.	Occurrence in food and animals	94
т.Э.1.	Occurrence in 1000 driu driinidis	34

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903j.efsa.2022.7666 by Istituto Zooprofilativo Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [1812.2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rate of use; OA articles are governed by the applicable Creative Commons License



4.3.2.	Serogroups and virulence features in food and animals	94
4.4.	Results	
4.4.1.	Overview of key statistics, EU, 2017–2021	
4.4.2.	STEC infections in humans	
	STEC in food	98
4.4.4.	STEC in animals	
4.4.5.	Focus on STEC strain features: virulence genes and serogroups	
4.5.	Discussion	
5.	Tuberculosis due to <i>Mycobacterium bovis</i> or <i>Mycobacterium caprae</i>	
5.1.	Key facts	
5.2.	Surveillance and monitoring of tuberculosis due to Mycobacterium bovis or Mycobacterium caprae in	
	the EU	107
5.2.1.	Humans	
	Animals	
5.3.	Results	
	Overview of key statistics, EU, 2017–2021	
5.3.2.	Tuberculosis due to <i>Mycobacterium bovis</i> and <i>Mycobacterium caprae</i> in humans	
	Mycobacterium in food	112
5.3.4.	Tuberculosis in bovine animals	
5.4.	Discussion	
6.	Brucella	
6.1.	Key facts	120
6.2.	Surveillance and monitoring of <i>Brucella</i> in the EU	
6.2.1.	Humans	
6.2.2.	Food and animals	
6.3.	Results	
	Overview of key statistics, EU, 2017–2021	
	Human brucellosis	
	Brucella in food	
	Brucella in animals	
6.4.	Discussion	
7.	Trichinella	
7.1. 7.2.	Key facts	
7.2. 7.2.1.	Humans	
	Animals	
7.2.2. 7.3.	Results	
	Overview of key statistics, EU, 2017–2021.	
7.3.1. 7.3.2.	Human trichinellosis.	
7.3.2. 7.3.3.	Trichinella infection in food and animals	
7.3.3. 7.4.	Discussion	
	Echinococcus	
8.1.	Key facts	
8.2.	Surveillance and monitoring of cystic and alveolar echinococcosis in humans and animals in the EU	149
	Humans	149
8.2.2.	Animals	149
8.3.	Results	151
	Overview of key statistics, EU, 2017–2021	151
	Human echinococcosis	
	Echinococcus spp. in animals and food	
8.4.	Discussion	
-	ne outbreaks (in accordance with Directive 2003/99/EC)	
1.	Key facts	
2.		
3.	Data analyses	
4.	Results and discussion	166
	Overview of countries reporting foodborne outbreak data in 2021	166
4.2.	Overview of causative agents in foodborne outbreaks in 2021	173
4.2.1.	Bacteria	173
4.2.2.	Bacterial toxins	179
4.2.3.	Viruses	179
4.2.4.	Parasites	
4.2.5.	Other causative agents	180

18314732, 2022, 12, Downloaded from https://efs.aonlinelibrary.wiey.com/doi/10.2903j.drs.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18122023]. See the Terms and Conditions (https://onlinelibrary.wiey.com/terms-and-conditions) on Wiley Online Library for rules of use; OA article are governed by the applicable Creative Commons.



4.2.6.	Outbreaks caused by unknown/unspecified agents	
4.3.	Overview of food vehicles implicated in foodborne outbreaks	182
4.3.1.	Food vehicles implicated in strong-evidence outbreaks	182
4.3.2.	Food vehicles implicated in weak-evidence outbreaks	
4.3.3.	Top-10 agent/food pairs in strong-evidence outbreaks associated with the highest impact on health	
1.5.5.		188
1 1	,	194
4.4.		
4.5.	Contributing factors in strong-evidence foodborne outbreaks	
4.6.	Temporal trends by causative agent	
4.6.1.	Temporal trend at EU level	
4.6.2.	Country-specific temporal trends by causative agent over the period 2012–2021	
4.6.3.	Temporal trends by implicated food vehicle	197
4.7.	Waterborne outbreaks	198
4.8.	ECDC and EFSA rapid outbreak assessment (ROA) of multi-country foodborne outbreaks	
5.	Conclusions	
5.1.	Health impact, causative agents and trends	
5.2.	Food vehicles and places of exposure	
	es monitored in accordance with the epidemiological situation (Directive 2003/99 List B)	200
	is monitored in accordance with the epidemiological situation (Directive 2003/99 List B)	202
1.	Yersinia	
1.1.	Key facts	203
1.2.	Surveillance and monitoring of <i>Yersinia</i> in the EU	
1.2.1.	Humans	203
1.2.2.	Food and animals	203
1.3.	Results	204
1.3.1.	Overview of key statistics, EU, 2017–2021	
1.3.2.	Human yersiniosis	
1.3.3.	Yersinia in food	
1.3.4.	Yersinia in animals	
1.4.	Discussion	
2.	Toxoplasma gondii	
2.1.	Key facts	
2.2.	Surveillance and monitoring of <i>Toxoplasma</i> in the EU	210
2.2.1.	Humans	210
2.2.2.	Animals and food	211
2.3.	Results	
2.3.1.	Human congenital toxoplasmosis	
2.3.2.	Toxoplasma gondii in food and animals	
2.4.	Discussion	21/
3.	Rabies	
3.1.	Key facts	
3.2.	Surveillance and monitoring of rabies in the EU	
3.2.1.	Humans	
3.2.2.	Animals	
3.3.	Results	
3.3.1.	Overview of key statistics, EU, 2017–2021	216
3.3.2.	Humans	217
3.3.3.		218
3.4.		221
4.	Q fever	
4.1.		222
4.2.	·	223
4.2.1.	Humans	
4.2.2.	Animals	
4.3.	Results	
4.3.1.	Overview of key statistics, EU, 2017–2021	224
		226
4.3.3.	Coxiella burnetii in animals	228
4.4.	Discussion	229
5.		230
5.1.		230
5.2.	WNV ecology and epidemiology	
5.3.	Surveillance and monitoring of West Nile virus in the EU	
5.3.1.	Humans	
J.J.I.	Tuttatis	۷۵.

18314732, 2022, 12, Downloaded from https://efs.aonlinelibrary.wiey.com/doi/10.2903j.drs.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18122023]. See the Terms and Conditions (https://onlinelibrary.wiey.com/terms-and-conditions) on Wiley Online Library for rules of use; OA article are governed by the applicable Creative Commons.



5.3.2.	Animals	231
5.4.	Results	231
5.4.1.	Overview of key statistics, EU, 2017–2021	231
5.4.2.	West Nile virus infections in humans	233
5.4.3.	West Nile virus infections in animals	
5.4.4.	Joint analysis of trends and seasonality	
5.5.	Discussion	
6.	Tularaemia	
6.1.	Key facts	
6.2.	Surveillance and monitoring of tularaemia in the EU.	
6.2.1.	Humans	
6.2.2.	Animals	
6.3.	Results	
6.3.1.	Overview of key statistics, EU, 2017–2021	
6.3.2.	Tularaemia in humans	
6.3.3.	Tularaemia in animals	
6.4.	Discussion	
7.	Other zoonoses and zoonotic agents	
7.1.	Bacillus spp	
7.2.	Caliciviruses (including norovirus)	
7.3.	Chlamydia spp.	
7.4.	Clostridium spp.	
7.5.	Hepatitis virus	
7.6.	Proteus spp	2 <del>4</del> 7
7.7.	Staphylococcus spp. and staphylococcal enterotoxins	
7.8.	Cysticercus spp	
7.9.	Leishmania	248
7.10.	Sarcocystis spp.	248
	Other	
Microb	piological contaminants subject to food safety criteria (Regulation (EC) No 2073/2005)	249
1.	Histamine	
1.1.	Histamine data in the context of Regulation (EC) No 2073/2005	249
1.2.	Other monitoring or surveillance data for histamine	
2.	Staphylococcal enterotoxins	
3.	Cronobacter sakazakii	
Note o	on outbreak data of SARS-CoV-2 in minks, mustelids and raccoon dogs, in EU	250
Refere	ences	250
Abbre	viations	260
	ry codes	
	Idix A – Number of tested samples for <i>Listeria monocytogenes</i> for the main ready-to-eat (RTE) food	
	pries, by reporting MS and non-MS countries, 2021	
Appen	dix B – Occurrence of <i>Listeria monocytogenes</i> at all sampling stages combined in ready-to eat (RTE)	
food c	rategories using a detection method, EU, 2019–2021	264
	dix C – Atlas of STEC serogroups in food, animals and humans in reporting MSs, EU, 2017–2021	
	dix D – STEC serogroups in food and animals, by reporting MS, EU, 2021	
	dix E – Atlas of STEC serogroups in food and animals in reporting MSs, EU, 2021	
	and a contract of the contract	-, -

1831/4732, 2022, 12. Downloaded from https://drs.o.onlinelibrary.wiley.com/doi/10.2903j.drs.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2021]. See the Terms and Conditions (thps://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License



# **List of Figures**

Figure 1	Reported numbers of cases and notification rates for confirmed human zoonoses in the EU, 2021 Data on congenital toxoplasmosis are not shown since 2021 data are not available yet	18
Figure 2	Trends in reported confirmed human cases of campylobacteriosis in the EU, by month, 2017–2021	29
Figure 3	Trend in reported confirmed human cases of non-typhoidal salmonellosis in the EU by month, 2017–2021	45
Figure 4	Prevalence of poultry flocks (breeding flocks of <i>Gallus gallus</i> , laying hens, broilers, breeding turkeys and fattening turkeys) positive for target <i>Salmonella</i> serovars, EU MSs and non-MS countries, 2021	<del>5</del> 7
Figure 5	Trend in the estimated prevalence of poultry flocks positive for <i>Salmonella</i> spp. and target <i>Salmonella</i> serovars, at EU level for different poultry populations, 2007–2021	67
Figure 6	Trend in reported confirmed human cases of <i>S</i> . Enteritidis infections acquired in the EU, by month, 2017–2021	71
Figure 7	Sankey diagram of the distribution of the top five EU-level <i>Salmonella</i> serovars involved in human salmonellosis cases acquired in the EU, reported from specified food–animal	73
Figure 8	categories, by food–animal source, EU, 2021  Overview of <i>Listeria monocytogenes</i> testing along the food chain according to the sampling stage, the sampler and the objective of the sampling, EU	81
Figure 9 Figure 10	Trends in reported confirmed human cases of listeriosis in the EU by month, 2017–2021  Trends in reported confirmed human cases of STEC infection in the EU by month,	85
_	2017–2021	98
Figure 11	Map of the number of confirmed tuberculosis cases due to <i>Mycobacterium bovis</i> and <i>Mycobacterium caprae</i> in individuals of EU origin, and national herd prevalence of tuberculosis in the bovine population in EU MS and non-MS countries, 2021	111
Figure 12	Status of countries for infection with the <i>Mycobacterium tuberculosis</i> complex ( <i>M. bovis</i> , <i>M. caprae</i> and <i>M. tuberculosis</i> ) in the bovine animal population, EU MSs and non-MSs, 2021	112
Figure 13	Prevalence of cattle herds infected with <i>Mycobacterium tuberculosis</i> complex in disease-free (DFS) zones, EU, 2012–2021	114
Figure 14	Prevalence of cattle herds infected with <i>Mycobacterium tuberculosis</i> complex in non-disease-free (non-DFS) zones, EU, 2012–2021	115
Figure 15	Prevalence of bovine tuberculosis-infected herds in non-disease-free zones of five MSs and the United Kingdom (Northern Ireland), 2005–2021	116
Figure 16	Trends in reported confirmed human cases of brucellosis in the EU, by month, 2017–2021	124
Figure 17	Number of confirmed, domestically acquired brucellosis cases in humans and national prevalence of <i>Brucella</i> -positive cattle herds, and sheep and goat herds, in EU MSs and non-MS countries, 2021	125
Figure 18	Status of countries as regards brucellosis in bovine animal population ( <i>Bison</i> ssp., <i>Bos</i> ssp., <i>Bubalus</i> ssp.), MSs and non-MSs, 2021	126
Figure 19	Prevalence of <i>Brucella</i> -positive cattle herds, in non-DF zones, EU, 2012–2021	128
Figure 20	Prevalence of <i>Brucella</i> -positive cattle herds, in two MSs with non-DFS zones (Italy and Portugal) and in one non-DFS MS (Greece), 2005–2021	129
Figure 21	Status of countries as regards ovine and caprine brucellosis, MSs and non-MS countries, 2021	130
Figure 22	Prevalence of <i>Brucella</i> -positive sheep and goat herds, in non-DFS zones, EU, 2012–2021	132
Figure 23	Prevalence of <i>Brucella</i> -positive sheep and goat herds, in two MSs with non-DFS zones (Italy and Portugal) and in one non-DFS MS (Greece), 2005–2021	133
Figure 24	Trend in reported confirmed human cases of trichinellosis in the EU by month, 2017–2021	140
Figure 25	Trichinella spp. in domestic pigs and farmed wild boar, in EU MSs and non-MS countries, 2012–2021	143
Figure 26	Pooled prevalence of <i>Echinococcus multilocularis</i> in red and Arctic foxes within the EU and adjacent countries, depicting the current epidemiological situation in Europe (Oksanen et al.,	
Fig. 27	2016)	150
Figure 27	Approximate geographical distribution of the <i>Echinococcus granulosus sensu lato</i> species causing human cystic echinococcosis in Europe (2000–2021) (Casulli et al., 2022)	151
Figure 28	Cumulative proportion (%) of test-positive animals for <i>Echinococcus granulosus sensu lato</i> , by intermediate host species, in EU MSs and non-MSs, 2017–2021	158
Figure 29	Map of the cumulative proportion (%) of test-positive animals for <i>Echinococcus granulosus</i> sensu lato in MSs and non-MSs, by country, 2017–2021	159
Figure 30	Relative variations (%) in foodborne outbreaks reported in 2021 compared with 2020, and mean annual totals for the pre-COVID-19 pandemic period 2017–2019, by reporting country	170

1831/372, 2022, 12, Downloaded from https://efsa.onlinelibmry.wiley.com/oi/10.2903j.efsa.2022.7666 by. Istituto Zooprofilatioo Speriment dell'Umbria e del Marche (IZSUM). Wiley Online Library on [1812/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/erms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons. License



Figure 31	Trends in the number of strong-evidence and weak-evidence outbreaks (left axis) and the outbreak reporting rate (per 100,000) (right axis) in the EU and reporting EU MSs and non-	173
Figure 32	MS countries, 2012–2021  Overview of EU MSs and non-MS countries reporting data on foodborne outbreaks, by causative agent, 2021	181
Figure 33	Number of foodborne outbreaks by causative agent, reported to the EU by MSs, 2012–2021	196
Figure 34	Trends in reported confirmed human cases of yersiniosis in the EU, by month, 2012–2021	206
Figure 35	Choropleth map of the number of tested and positive foxes, and the geographical distribution of the rabies cases reported in foxes in EU MSs and non-EU countries, 2021	218
Figure 36	Choropleth map of the number of tested and positive bats, and the geographical distribution of the rabies cases reported in bats in EU MSs and non-EU countries, 2021	219
Figure 37	Choropleth map of the number of tested and positive pets, and the geographical distribution of the rabies cases reported in pets in EU MSs and non-EU countries, 2021	220
Figure 38	Choropleth map of the number of tested and positive farmed animals, and the geographical distribution of the rabies cases reported in farmed animals in EU MSs and non-EU countries,	
Figure 39 Figure 40	Trend in reported confirmed human cases of Q fever in the EU by month, 2017–2021  Geographical distribution of locally acquired West Nile virus infections among humans (NUTS 3 level) and outbreaks notified to ADIS among equids and birds (XY coordinates) across the	221 228
Figure 41	EU, 2021 transmission season	233 238
Figure 42	Reported human cases of West Nile virus infection in the EU MSs, by month, 2017–2021	239
Figure 43 Figure 44	Outbreaks of West Nile infection in animals in the EU MSs, by month, 2017–2021 Trends in reported confirmed human cases of tularaemia in the EU, by month and year,	240
	2017–2021	244
List of Tabl	es	
Table 1	Categorisation of the data used in the EU One Health Zoonoses 2021 Summary Report (adapted from Boelaert et al. (2016))	15
Table 2	Reported hospitalisations and deaths due to zoonoses in confirmed human cases and among foodborne outbreak cases in the EU, 2021	19
Table 3	Number of confirmed human cases and notification rates (per 100,000 population) in 2021 by zoonosis, including the absolute and relative (%) difference with regard to 2020, and the mean annual cases and mean notification rates (per 100,000 population) for the 2017–2019	
Table 4	period, EU	20
Table 5	main animal species, EU, 2017–2021	26
Table 6	population in EU MSs and non-MS countries, by country and year, 2017–2021	27 31
Table 7	sampler and reporting MS, EU, 2021  Occurrence of <i>Campylobacter</i> in the main food categories, EU, 2021 and 2017–2020	33
Table 8	Summary of <i>Campylobacter</i> statistics related to major animal species, reporting EU MSs and non-MS countries, 2021	34
Table 9	Summary of Salmonella statistics related to humans, major food categories and the main animal species, EU, 2017–2021	42
Table 10	Reported confirmed human cases of salmonellosis and notification rates per 100,000	
Table 11	population in EU MSs and non-MS countries, by country and year, 2017–2021 Proportion (%) of <i>Salmonella</i> -positive samples from official sampling as part of the verification of <i>Salmonella</i> FSC according to Regulation (EC) No 2073/2005, by stage in the	44
Table 12	food chain, EU, 2021	46
Table 13	after dressing, but before chilling, by sampler and reporting MS, EU, 2021	48
Table 14	(neck skin samples) after chilling, by sampler and reporting MS, EU, 2021	49
	(neck skin samples) after chilling, by sampler and reporting MS, EU, 2021	50

18314732, 2022, 12, Downloaded from https://drs.aonlinelibrary.wiey.com/doi/10.29033j.drs.2022.7666 by Istituto Zooprofilattico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiey.com/terms-and-conditions) on Wiley Online Library for rules of use; OA attacks are governed by the applicable Creative Commons



1831/372, 2022, 12, Downloaded from https://efsa.onlinelibmry.wiley.com/oi/10.2903j.efsa.2022.7666 by. Istituto Zooprofilatioo Speriment dell'Umbria e del Marche (IZSUM). Wiley Online Library on [1812/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/erms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons. License

Table 15	Comparisons of proportions (%) of <i>Salmonella</i> -positive single samples from bovine carcases after dressing but before chilling, by sampler and reporting MS, EU, 2021	51
Table 16	Comparisons of proportions (%) of Salmonella-positive single samples from sheep carcases	
Table 17	after dressing but before chilling, by sampler and reporting MS, EU, 2021	52
Table 18	after dressing but before chilling, by sampler and reporting MS, EU, 2021	53
Table 19	after dressing but before chilling, by sampler and reporting MS, EU, 2021  Occurrence of Salmonella in the main food categories, EU, 2021 and 2017–2020	53 55
Table 20	Salmonella in breeding flocks of Gallus gallus during the production period (all types of	55
Tubic 20	breeding flocks, flock-based data) in countries running control programmes in accordance with Regulation (EC) No 2160/2003, 2021	58
Table 21	Salmonella in laying hen flocks of Gallus gallus during the production period (flock-based	-
	data) in countries running control programmes in accordance with Regulation (EC) No 2160/2003, 2021	59
Table 22	Salmonella in broiler flocks of Gallus gallus before slaughter (flock-based data) in countries	
	running control programmes in accordance with Regulation (EC) No 2160/2003, 2021	60
Table 23	Comparisons of the prevalence of target Salmonella serovar-positive broiler flocks, by	
	sampler and by reporting MS, EU, 2021	61
Table 24	Salmonella in breeding flocks of turkeys during the production period (flock-based data) in countries running control programmes in accordance with Regulation (EC) No 2160/2003,	62
Table 25	2021	62
Table 25	Comparisons of the prevalence of target <i>Salmonella</i> serovar-positive flocks of breeding	<b>C</b> 2
Table 26	turkeys, by sampler and by reporting MS, EU, 2021	63
Tubic 20	running control programmes, 2021	64
Table 27	Comparisons of the prevalence of target <i>Salmonella</i> serovar-positive flocks of fattening	٠.
	turkeys, by sampler and by reporting MS, EU, 2021	65
Table 28	Summary of <i>Salmonella</i> statistics related to major animal species, reporting MSs and non-MS countries, 2021	68
Table 29	Distribution of reported confirmed cases of human salmonellosis in the EU, 2019–2021, for	00
	the 20 most frequent <i>Salmonella</i> serovars in 2021	69
Table 30	Distribution of reported confirmed cases of human salmonellosis acquired in the EU, 2019–	
	2021, for the six most frequent Salmonella serovars in 2021	70
Table 31	Distribution of <i>Salmonella</i> isolates (number and percentage of positive sampling units) with and without serotype identification among the different selected sources (food and	71
Table 32	animals), EU, 2021	71
Table 32	turkeys, pigs and bovines), EU, 2021	74
Table 33	Listeria monocytogenes FSC as described in Regulation (EC) No 2073/2005 for the different RTE categories across the food chain, 2021	81
Table 34	Summary of <i>Listeria monocytogenes</i> statistics related to invasive human infections and on	01
Tubic 5 i	sampled major RTE food categories in the EU, 2017–2021	83
Table 35	Reported confirmed human cases of invasive listeriosis and notification rates per 100,000	00
	population in EU MSs and non-MS countries, by country and year, 2017–2021	84
Table 36	Proportions (%) of positive single samples at the manufacturing and distribution stage from	
	official sampling by CAs in the context of verification of implementation by FBOps of the	
	Listeria monocytogenes FSC according to Regulation (EC) No 2073/2005, EU, 2021	86
Table 37	Summary of <i>Listeria</i> species statistics related to major animal species, MSs, 2021	89
Table 38	Summary of STEC statistics related to humans, major food categories and the main animal	
Table 20	species, EU, 2017–2021	95
Table 39	Reported human cases of STEC and notification rates per 100,000 population in EU MSs	96
Table 40	and non-MS countries, by country and year, 2017–2021  Occurrence of Shiga toxin-producing <i>E. coli</i> (STEC) in the main food categories, EU, 2021	90
Iabic TU	and 2017–2020	99
Table 41		101
Table 42	Virulotypes ( <i>stx</i> type and presence of <i>eae</i> ) in food, animal and human STEC isolates	
	causing severe infection (haemolytic-uraemic syndrome (HUS), hospitalisation and bloody	
	diarrhoea) in 2021 and comparison with those associated with severe disease in humans	
	during 2012–2017, EU	102



Table 43	Virulotypes ( <i>stx</i> subtype) of food and human STEC isolates causing severe infection (haemolytic-uraemic syndrome, hospitalisation and bloody diarrhoea) in 2021 and	
	comparison with those associated with severe disease in humans during 2012–2017, EU	102
Table 44	Summary statistics related to tuberculosis due to <i>Mycobacterium bovis</i> and <i>Mycobacterium caprae</i> in humans and bovine animals (stratified by disease-free <sup>(a)</sup> and non-disease-free	
	status zones), EU, 2017–2021	108
Table 45	Reported cases of human tuberculosis due to <i>Mycobacterium bovis</i> and <i>Mycobacterium</i>	
	caprae and notification rates per 100,000 population in EU MS and non-MS countries by	
	country and year, 2017–2021	110
Table 46 Table 47	Status of countries as regards bovine tuberculosis and related prevalence, EU, 2021	113
	species (stratified by disease-free status <sup>(a)</sup> and non-disease-free status zones), EU, 2017–	122
T-bl- 40	2021	122
Table 48	Reported confirmed human cases of brucellosis and notification rates per 100,000 population in EU MSs and non-MSs countries, by country and year, 2017–2021	123
Table 49	Status of countries as regards brucellosis in bovine animals ( <i>Bos</i> ssp., <i>Bubalus</i> ssp. and	123
Table 49	Bison ssp.) and related prevalence, EU, 2021	127
Table 50	Status of countries for brucellosis in sheep ( <i>Ovis</i> ssp.) and goat ( <i>Capra</i> ssp.) populations	127
	and related prevalence, EU, 2021	131
Table 51	Summary of <i>Trichinella</i> statistics related to humans and main animal species, EU, 2017–	
	2021	137
Table 52	Reported confirmed human cases of trichinellosis and notification rates per 100,000	
	population in EU MSs and non-MS countries, by country and year, 2017–2021	138
Table 53	Trichinella monitoring results for domestic pigs and farmed wild boar in reporting EU MSs	
	and non-MS countries, by housing conditions <sup>(a)</sup> , 2021	141
Table 54	Trichinella monitoring results for hunted wild boar and wild boar with unspecified habitat,	
	bears, foxes and domestic solipeds, in reporting EU MSs and non-MS countries, 2021	143
Table 55	Trichinella monitoring results in other wild animals in reporting EU MSs and non-MS	
	countries, 2021	145
Table 56	Summary of echinococcosis in humans, caused by <i>Echinococcus multilocularis</i> or	
	Echinococcus granulosus sensu lato (s.l.) in the most relevant definitive and intermediate	454
T	animal hosts in the EU, 2017–2021	151
Table 57	Reported human cases of cystic and alveolar echinococcosis and notification rates per	450
Table 58	100,000 population in EU MSs and non-MS countries, by country and year, 2017–2021	153
Table 36	Reported human cases of cystic and alveolar echinococcosis in EU MSs and non-MS countries, by country, year and <i>Echinococcus</i> species, 2017–2021	154
Table 59	Echinococcus multilocularis monitoring results in wild and domestic animals in reporting EU	134
Table 33	MSs and non-MSs, 2021	156
Table 60	Echinococcus granulosus sensu lato monitoring results in wild and domestic animals in	150
	reporting EU MSs and non-MSs, 2021	160
Table 61	Number of foodborne outbreaks, human cases, hospitalisations and deaths, in reporting EU	
	MSs and non-MS countries, 2021	168
Table 62	Number of foodborne outbreaks, human cases, hospitalisations and deaths, by causative	
	agents, in reporting EU MSs, 2021	175
Table 63	Frequency distribution of strong-evidence foodborne outbreaks, by food vehicle, in reporting	
	EU MSs, 2021	185
Table 64	Top 10 pathogen/food vehicle pairs causing the highest number of strong-evidence	
	outbreaks in reporting EU MSs, 2021	189
Table 65	Top-10 pathogen/food vehicle pairs causing the highest number of cases in strong-evidence	
Table 66	outbreaks in reporting EU MSs, 2021	190
Table 66	Top-10 pathogen/food vehicle pairs causing the highest number of hospitalisations in	100
Table 67	strong-evidence outbreaks in reporting EU MSs, 2021	192
Table 07	evidence outbreaks in reporting EU MSs, 2021	193
Table 68	Frequency distribution of strong-evidence foodborne outbreaks by place of exposure, in	193
Table 00	reporting EU, MSs, 2021	195
Table 69	Summary of <i>Yersinia</i> statistics related to humans, major food categories and main animal	1,7,3
	species, EU, 2017–2021	204
Table 70	Reported confirmed human cases of yersiniosis and notification rates per 100,000	
	population in EU MSs and non-MS countries, by country and year, 2017–2021	205
Table 71	Occurrence of <i>Yersinia</i> in the main food categories, EU, 2021 and 2017–2020	207

10

18314732, 2022, 12, Downloaded from https://drs.aonlinelibrary.wiey.com/doi/10.29033j.drs.2022.7666 by Istituto Zooprofilattico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiey.com/terms-and-conditions) on Wiley Online Library for rules of use; OA attacks are governed by the applicable Creative Commons



Table 72	Summary of <i>Yersinia enterocolitica</i> statistics related to major animal species, reported by EU	200
Table 73	MSs and non-MS countries, 2021	208
	100,000 live births in EU MSs and non-MS countries, by country and year, 2016–2020	212
Table 74	Summary of <i>Toxoplasma gondii</i> statistics in the main animal species, EU, 2017–2021	213
Table 75	Summary of rabies lyssavirus statistics related to humans and the main animal reservoirs,	216
Table 76	EU, 2017–2021	216
Table 70	2017–2021	224
Table 77	Reported confirmed human cases of Q fever and notification rates per 100,000 population	
	in EU MSs and non-MS countries, by country and year, 2017–2021	227
Table 78	Summary of WNV infection statistics related to humans, birds and equids EU, 2017–2021	232
Table 79	Locally acquired human WNV cases (confirmed and probable) and notification rates per 100,000 population in EU MSs and non-MS countries, by country and year, 2017–2021	234
Table 80	Locally acquired human WNND cases (confirmed and probable) and notification rates per	234
	100,000 population in the EU MSs and non-MS countries, by country and year, 2017–2021	235
Table 81	Summary of WNV surveillance/monitoring results in animals reported to EFSA, and WNV	
	outbreaks notified to ADIS, by EU MSs and non-MS countries, 2021	236
Table 82	Summary of <i>Francisella tularensis</i> statistics related to humans and the main animal species,	242
Table 83	EU, 2017–2021	242
Table 05	population in EU MSs and non-MS countries, by country and by year, 2017–2021	243
Table 84	Francisella tularensis monitoring results for animals, by species in reporting EU MSs and	0
	non-MS countries, 2021	245

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903j.efsa.2022.7666 by Istituto Zooprofilativo Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [1812.2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rate of use; OA articles are governed by the applicable Creative Commons License



## Introduction

# Legal basis of European Union-coordinated zoonoses monitoring

The European Union (EU) system for the monitoring and collection of information on zoonoses is based on Zoonoses Directive 2003/99/EC¹, which obliges EU Member States (MSs) to collect relevant and, when applicable, comparable data on zoonoses, zoonotic agents, antimicrobial resistance and foodborne outbreaks. In addition, MSs shall assess the trends and sources of these agents, as well as outbreaks in their territory, submitting an annual report each year by the end of May to the European Commission covering the data collected. The European Commission should subsequently forward these reports to the European Food Safety Authority (EFSA). EFSA is assigned the tasks of examining these data and publishing the EU Annual Summary Reports. In 2004, the European Commission entrusted EFSA with the task of setting up an electronic reporting system and database for monitoring zoonoses (EFSA Mandate No 2004-0178, continued by M-2015-0231²).

Data collection on human diseases from MSs is conducted in accordance with Decision 1082/2013/EU<sup>3</sup> on serious cross-border threats to health. In October 2013, this decision replaced Decision 2119/98/EC on setting up a network for the epidemiological surveillance and control of communicable diseases in the EU. The case definitions to be followed when reporting data on infectious diseases to the European Centre for Disease Prevention and Control (ECDC) are described in Decision 2018/945/EU<sup>4</sup>. ECDC has provided data on zoonotic infections in humans, as well as their analyses, for the EU Summary Reports since 2005. Since 2008, data on human cases have been received via The European Surveillance System (TESSy), maintained by ECDC.

# Reporting requirements

In accordance with List A of Annex I of Zoonoses Directive 2003/99/EC, data on animals, food and feed must be reported on a mandatory basis for the following eight zoonotic agents: *Salmonella*, *Campylobacter*, *Listeria monocytogenes*, Shiga toxin-producing *Escherichia coli* (STEC), *Mycobacterium bovis*, *Brucella*, *Trichinella* and *Echinococcus*. In addition, and based on the epidemiological situations in the MSs, data must be reported on the following agents and zoonoses (List B of Annex I of the Zoonoses Directive): (i) viral zoonoses: calicivirus, hepatitis A virus, influenza virus, rabies, viruses transmitted by arthropods; (ii) bacterial zoonoses: borreliosis and agents thereof, botulism and agents thereof, leptospirosis and agents thereof, psittacosis and agents thereof, tuberculosis due to agents other than *M. bovis*, vibriosis and agents thereof, cysticercosis and agents thereof, toxoplasmosis and agents thereof, cryptosporidiosis and agents thereof, cysticercosis and agents thereof, toxoplasmosis and agents thereof; and (iv) other zoonoses and zoonotic agents such as *Francisella* and *Sarcocystis*. Furthermore, MSs provided data on certain other microbiological contaminants in foods: histamine, staphylococcal enterotoxins and *Cronobacter sakazakii*, for which food safety criteria are set down in the EU legislation.

In accordance with Article 9 of the Directive, MSs shall assess the trends and sources of zoonoses, zoonotic agents and antimicrobial resistance in their territory and each MS shall send to the European Commission every year by the end of May a report on the trends and sources of zoonoses, zoonotic agents and antimicrobial resistance. Reports and any summaries of these shall be made publicly available.

# **Terms of Reference**

In accordance with Article 9 of Directive 2003/99/EC, EFSA shall examine the national reports and data submitted by the EU MSs regarding their 2021 zoonoses monitoring activities as described above and publish an EU Summary Report on the trends and sources of zoonoses, zoonotic agents and antimicrobial resistance in the EU.

The 2021 data of MSs on antimicrobial resistance in zoonotic agents are published in a separate EU Summary Report.

<sup>&</sup>lt;sup>1</sup> Directive 2003/99/EC of the European Parliament and of the Council of 17 November 2003 on the monitoring of zoonoses and zoonotic agents, amending Council Decision 90/424/EEC and repealing Council Directive 92/117/EEC. OJ L 325, 12.12.2003 pp. 31–40.

<sup>&</sup>lt;sup>2</sup> See mandate M-2015-0231 in OpenEFSA Question: https://open.efsa.europa.eu/questions/EFSA-Q-2020-00787

<sup>&</sup>lt;sup>3</sup> Decision No. 1082/2013/EU of the European Parliament and of the Council of 22 October 2013 on serious cross-border threats to health and repealing Decision No 2119/98/EC. OJ L 293, 5.11.2013, pp. 1–15.

<sup>&</sup>lt;sup>4</sup> Commission Implementing Decision 2018/945/EU on the communicable diseases and related special health issues to be covered by epidemiological surveillance as well as relevant case definitions. OJ L 170, 6.7.2018, pp. 1–74.



# **Data sources and report production**

Since 2019, the annual EU Summary Reports on zoonoses, zoonotic agents and foodborne outbreaks have been renamed the 'EU One Health Zoonoses Summary Report' (EUOHZ), which is co-authored by EFSA and ECDC.

Since 2020, the production of the annual EUOHZ report has been supported by the ZOE (**Z**oonoses under a **O**ne health perspective in the **E**U) Consortium's Work-package 1 composed of the *Istituto Superiore di Sanità* (Rome, Italy), the *Istituto Zooprofilattico Sperimentale delle Venezie* (Padova, Italy), the French Agency for Food, Environmental and Occupational Health & Safety (Maisons-Alfort, France), the *Istituto Zooprofilattico Sperimentale dell'Abruzzo e del Molise* (Teramo, Italy) and the *Istituto Zooprofilattico Sperimentale dell'Abruzzo e del Molise* (Teramo, Italy), under the coordination of the *Istituto Zooprofilattico Sperimentale dell'Abruzzo e del Molise* (Teramo, Italy).

The efforts made by the MSs, the reporting non-MSs and the European Commission in the reporting of zoonoses data and in the preparation of this report are gratefully acknowledged.

The MSs, other reporting countries, the European Commission, members of EFSA's Scientific Panels on Biological Hazards (BIOHAZ) and Animal Health and Welfare (AHAW) and the relevant European Union Reference Laboratories (EURLs) were consulted while preparing the EUOHZ 2021 report.

The EUOHZ 2021 report focuses on the most relevant information on zoonoses and foodborne outbreaks within the EU in 2021. Where substantial differences when compared to the previous years were observed, they have been reported.

In 2020, data collection was affected by the reduction in the number of EU MSs from 28 to 27, due to the withdrawal of the United Kingdom from the EU.<sup>5</sup> On 1 February 2020, the United Kingdom became a third country. In descriptive tables, data from the United Kingdom were included in the EU statistics for 2019 and previous years, whereas the 2020 statistical data from the United Kingdom, when available (for EFSA data), were assigned to the non-MSs group. Human data from the United Kingdom were not collected by ECDC as of 2020. With regard to trend analyses of human data, only countries having contributed data for all the years of the considered period were taken into account. For trend analyses of the estimated prevalence at EU level of *Salmonella* in poultry populations covered by National Control Programs, any data provided by the reporting MSs were taken into account in the model. The United Kingdom data were only included when available for 2019 and previous years.

In 2021, the only United Kingdom data that were reported to EFSA were from Northern Ireland. In accordance with the Agreement on the withdrawal of the United Kingdom from the European Union, and in particular with the Protocol on Ireland/Northern Ireland, the European Union requirements on data sampling are also applicable to and in the United Kingdom with respect to Northern Ireland. Therefore, pursuant to Article 5(4) and Section 24 of Annex 2 of the Protocol on Ireland/Northern Ireland, which is an integral part of the Agreement on the withdrawal of the United Kingdom of Great Britain and Northern Ireland from the European Union and the European Atomic Energy Community, for the purpose of this report, references to MSs should be read as including Northern Ireland, despite it being part of the United Kingdom. Hence, the European Union requirements on data sampling were also applicable to Northern Ireland and data transmitted by the United Kingdom (Northern Ireland) have been assigned to the MSs group. For the collection of data, EFSA aligned with the guidelines of the Commission concerning customs registration, which lay down the following abbreviations and terminology:

- GB, which stands for 'the United Kingdom' and refers to: Great Britain, Northern Ireland, the Channel Islands and the Isle of Man.
- XI, which stands for 'the United Kingdom (Northern Ireland)', and is used when the United Kingdom is identified with respect to Northern Ireland.
- XU, which stands for 'the United Kingdom (excluding Northern Ireland)', and is used when the United Kingdom excluding Northern Ireland is identified.

18314732, 2022, 12, Downloaded from https://fs.aonlinelibrary.wiley.com/inchibrary.wiley.com/inchibrary.wiley.com/em-and-conditions) on Wiley Online Library on [18/12/2023], See the Terms and Conditions (http://oinineibrary.wiley.com/em-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons

<sup>&</sup>lt;sup>5</sup> Agreement on the withdrawal of the United Kingdom of Great Britain and Northern Ireland from the European Union and the European Atomic Energy Community. OJ L 29, 31.1.2020, p. 7 ('Withdrawal Agreement').

<sup>6</sup> https://taxation-customs.ec.europa.eu/system/files/2020-11/use\_of\_gb\_and\_xi\_codes\_guidance.pdf



#### Human data collection for 2021

In the 2021 EUOHZ report, the analyses of data from infections in humans were prepared by the Food- and Waterborne Diseases and Zoonoses (FWD) domain (brucellosis, campylobacteriosis, congenital toxoplasmosis, echinococcosis, listeriosis, salmonellosis, Shiga toxin-producing *E. coli* infection, trichinellosis and yersiniosis), the Emerging and Vectorborne Diseases (EVD) domain (Q fever, rabies, tularaemia and West Nile virus (WNV) infection) and the tuberculosis (TB) domain (TB due to *Mycobacterium bovis* and *M. caprae*) at ECDC. Please note, as explained above, that the numbers presented in the report may differ from those in national reports due to differences in the case definitions used at the EU and at national levels, or due to differing dates of data submission and extraction. The latter may also result in some divergence in the case numbers presented in the different ECDC reports.

TESSy is a software platform in which data on 56 diseases and special health issues are collected. Both aggregated and case-based data are reported to TESSy by MSs and other European countries. Although aggregated data did not include individual case-based information, both reporting formats were included, when possible, to calculate the number of cases and country-specific case notification rates. Human data used in the report were extracted from TESSy as of 20 July 2022 for EVD, as of 25 July 2022 for FWD and as of 30 September 2022 for TB due to *M. bovis* and *M. caprae*. The denominators used for calculating notification rates were based on the human population data from Eurostat on 01 January 2022.

Data on human cases were received from 27 MSs and from three non-MSs (Iceland, Liechtenstein and Norway). Switzerland reported its data on human cases directly to EFSA. For 2021, Liechtenstein reported distinct data on human cases for the first time since 2008, whereas in the past years, these data were reported together with Switzerland.

Since the United Kingdom became a third country on 1 February 2020, human data from the United Kingdom were not collected by ECDC for 2020 and afterwards.

When interpreting data, data quality issues should be considered, as well as the differences between MS surveillance systems; comparisons between countries should therefore be undertaken with caution.

# Data collection on food, animals, feed and foodborne outbreaks

In 2021, 27 MSs and the United Kingdom (Northern Ireland) submitted data and national zoonoses reports on monitoring results in food, animals, feed and foodborne outbreaks. In addition, data and reports were submitted by four non-MSs and the European Free Trade Association (EFTA) countries: Iceland, Norway, Switzerland and Liechtenstein.<sup>7</sup> For some food, animal and feed matrices, and for foodborne outbreaks, EFSA received data and reports from the following pre-accession countries: Albania, Bosnia and Herzegovina, the Republic of North Macedonia, Montenegro and Serbia.

Data were submitted electronically to the EFSA zoonoses database, through EFSA's Data Collection Framework (DCF). MSs could also update their data from previous years.

The deadline for data submission was 31 May 2022. Two data validation procedures were carried out, from 2 June to 14 June 2022 and from 7 July to 15 July 2022, respectively. Validated data on food, animals, feed and foodborne outbreaks used in the report were extracted from the EFSA zoonoses database on 26 July 2022.

The draft EUOHZ report was sent to the MSs for consultation on 12 October 2022 and comments were collected by 26 October 2022. The utmost effort was made to incorporate comments within the available time frame. Data amended after the data validation period that ended on 26 July 2022 have been not considered in summary calculations or other analyses, and footnotes to tables and figures have been added to inform on these late data corrections. The report was finalised by 11 November 2022 and published online by EFSA and ECDC on 8 December 2022.

A detailed description of the terms used in the report is available in EFSA's manuals for reporting on zoonoses (EFSA, 2022, 2022b,c).

<sup>&</sup>lt;sup>7</sup> Based on the customs union treaty of the Principality of Liechtenstein with Switzerland, Liechtenstein is part of the Swiss customs territory. Due to the strong connection between the veterinary authorities of Liechtenstein and Switzerland, and Liechtenstein's integration into the Swiss system in the veterinary field, in principal, all legislation, rules and data on contagious diseases are identical for both Switzerland and Liechtenstein. If not mentioned otherwise, the Swiss data also include the data from Liechtenstein.

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903j.efsa.2022.7666 by Istituo Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

The national zoonoses reports submitted in accordance with Directive 2003/99/EC are published on the EFSA website together with the EU One Health Zoonoses Report. They are available online at <a href="http://www.efsa.europa.eu/en/biological-hazards-data/reports">http://www.efsa.europa.eu/en/biological-hazards-data/reports</a>.

# **Data analyses and presentation**

# Comparability and quality of data

#### **Humans**

For data on human infections, please note that the numbers presented in this report may differ from national zoonoses reports due to differences in case definitions used at the EU and national levels or because of differing dates of data submission and extraction. Results are not directly comparable among the MSs.

## Food-animals-feed and foodborne outbreaks

For data on food, animals and feed, please note that the numbers presented in this report may differ from national zoonoses reports due to differing dates of data submission and extraction.

The data obtained by the EFSA DCF can vary according to the level of data quality and harmonisation. Therefore, the type of data analyses suggested by EFSA for each zoonosis and matrix (food, animals, feed or foodborne outbreaks) strongly depended on this level of harmonisation and can either be a descriptive summary of submitted data, the follow-up of trends (trend watching) or the (quantitative) analyses of trends. Data analyses were carried out according to Table 1, as adapted from Boelaert et al. (2016). Food, animals, feed and foodborne outbreak data can be classified into three categories according to the zoonotic agent monitored and the design of the monitoring or surveillance carried out. It follows that the type of data analyses that can be implemented is conditioned by these three distinct categories.

**Table 1:** Categorisation of the data used in the EU One Health Zoonoses 2021 Summary Report (adapted from Boelaert et al., 2016)

Category	Type of analysis		Type/comparability between MSs	Examples
Ī	Descriptive summaries at the national level and EU level		Programmed harmonised monitoring or surveillance	Salmonella national control programmes in poultry, bovine tuberculosis, bovine and small
	EU trend watching (trend monitoring)	$\vdots$	Comparable between MSs	ruminant brucellosis, <i>Trichinella</i> in pigs at slaughterhouse
	Spatial and temporal trend analyses at the EU level	0	Results at the EU level are interpretable	
II	Descriptive summaries at national level and EU level		Monitoring or surveillance not fully harmonised	Foodborne outbreak data; official samplings related to process hygiene criteria for carcases at the
	EU trend watching (trend monitoring)	···	Not fully comparable between MSs	slaughterhouse for <i>Salmonella</i> and <i>Campylobacter</i> and to food safety criteria for <i>Campylobacter</i> , <i>L. monocytogenes</i> , <i>Salmonella</i> and
	No EU trend analysis		Caution needed when interpreting results at the EU level	STEC in the context of Regulation (EC) No. 2073/2005; Rabies monitoring, West Nile virus



com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilattico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

Category	Type of analysis	Type/comparability between MSs	Examples
III	Descriptive summaries at national level and EU level	Non-harmonised monitoring or surveillance data with no (harmonised) reporting requirements	Campylobacter, Yersinia, Q fever, Francisella tularensis, Taenia spp., Toxoplasma and other zoonoses
	No EU trend watching (trend monitoring) No EU trend analysis	Not comparable between MSs; extreme caution needed when interpreting results at the EU level	

#### Rationale of the table of contents

In keeping with the rationale of the zoonoses listing in Annex I of Directive 2003/99/EC, for the mandatory reporting of foodborne outbreaks and of the above-mentioned categorisation of food, animal and feed data (Table 1), the following table of contents has been adopted for the 2021 EUOHZ report.

Zoonoses and zoonotic agents included in compulsory annual monitoring (Directive 2003/99/EC List A):

- 1) Campylobacter
- 2) Salmonella
- 3) Listeria
- 4) Shiga toxin-producing Escherichia coli
- 5) Tuberculosis due to Mycobacterium bovis and Mycobacterium caprae
- 6) Brucella
- 7) Trichinella
- 8) Echinococcus

Foodborne and waterborne outbreaks (according to Directive 2003/99/EC).

Zoonoses and zoonotic agents monitored according to the epidemiological situation (Directive 2003/99/EC List B):

- 1) Yersinia
- 2) Toxoplasma gondii
- 3) Rabies
- 4) Q fever
- 5) West Nile virus
- 6) Tularaemia
- 7) Other zoonoses and zoonotic agents

Microbiological contaminants subject to food safety criteria (Regulation (EC) No 2073/2005).

After the section on 'Microbiological contaminants subject to food safety criteria (Regulation (EC) No 2073/2005)', a note was added in the 2021 EUOHZ report on outbreak data of SARS-CoV-2 in minks, mustelids and raccoon dogs, in EU.

### **Chapter sections**

The 2021 EUOHZ Report presents a harmonised structure for each chapter:

- 'Key facts',
- 'Monitoring and surveillance' in the EU for the specific disease,
- 'Results', summarising the major findings of 2021 as regards trends and sources, starting with a table displaying summary statistics for the last 5 years (2017–2021) for human cases, food matrices and major animal species,
- A 'Discussion' section. For foodborne and waterborne outbreaks, the main findings are presented and discussed in a joint 'Results and discussion' section and key messages are summarised in the 'Conclusions' section.

For each chapter, overview tables present the data reported by each country. However, for the tables summarising MS-specific results and providing EU-level results for food, animals and feed, unless stated otherwise, data from industry own-check programmes, hazard analysis and critical

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (thps://onlinelibrary.wiley.com/terms

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

control point (HACCP) sampling, as well as data from suspect sampling, selective sampling and outbreak or clinical investigations are excluded. Moreover, regional data reported by countries for food, animals and feed, without statistics at the national level were also excluded from these summary tables.

## **Data analyses**

Statistical trend analyses in humans were carried out to evaluate the significance of temporal variations in the EU (further details in each chapter) in years 2017–2021. The number of confirmed cases for the EU by month is presented as a trend figure for years 2012–2021. All countries that consistently reported cases – or reported zero cases over the whole reporting period – were included. The trend figure also shows a centred 12-month moving average over the last 5 years, illustrating the overall trend by smoothing seasonal and random variations. Moreover, the same trend analysis was carried out separately for each country (MS and non-MS countries). Analyses were carried out considering confirmed cases only, except for WNV infection, for which total locally acquired cases (i.e. probable and confirmed cases) were considered.

The notification rates were calculated taking into account the coverage of the human population under surveillance (percentage of national coverage). For countries where surveillance did not cover the whole population, the estimated coverage – if provided – was used to calculate the country-specific rate. Cases and populations of those countries not providing information on national coverage or reporting incomplete data were excluded from the EU notification rate.

ESRI ArcMap 10.5.1 was used to map the data. Choropleth maps with graduated colours over five class scales of values, according to the natural breaks function proposed by the ArcGIS software, were used to map the proportion of positive sample units across the EU and other reporting countries. In the maps included in the present report, EU MSs and the United Kingdom (Northern Ireland) were represented with a blue label, whereas all the non-EU MSs (including the EFTA countries: Iceland, Norway, Switzerland and Liechtenstein; and the pre-accession countries: Albania, Bosnia and Herzegovina, the Republic of North Macedonia, Montenegro and Serbia) were represented with an orange label.

Statistical trend analysis of foodborne outbreaks was performed to evaluate the significance of temporal variations at the single MS level over the 2012–2021 period.

Summary data and figures on food, animals, feed and foodborne outbreaks used to produce this report, as well as additional information on related projects and internet sources are published for this report on the EFSA Knowledge Junction on the Zenodo general-purpose open-access repository here. All country-specific data on food, animals, feed and foodborne outbreaks updated until 30 November 2022 are also available at this URL.

With the present report, EFSA has also published interactive communication tools:

- the EFSA story maps on *Campylobacter* (here), *Salmonella* (here), *Listeria monocytogenes* (here) and foodborne outbreaks (here).
- the EFSA dashboards on *Campylobacter* (here), *Salmonella* (here), *Listeria monocytogenes* (here) and foodborne outbreaks (here).


Data used in these communication tools were extracted from the EFSA zoonoses database on 26 July 2022.

# Summary of human zoonoses data for 2021

The numbers of confirmed human cases of the zoonoses presented in this report are summarised in Figure 1. In 2021, campylobacteriosis was confirmed as the most commonly reported zoonosis (as it has been since 2005). It accounted for more than 62% of all the reported confirmed human cases in 2021. After campylobacteriosis, salmonellosis, yersiniosis, STEC infections and listeriosis were the most frequently reported. The severity of the diseases was descriptively analysed based on hospitalisations and the outcomes of reported cases (Table 2). Based on severity data, listeriosis and West Nile virus infection were the two most severe diseases with the highest case fatality and hospitalisation rates. Almost all confirmed cases with available hospitalisation data for these two diseases were hospitalised (96.5% of cases for listeriosis and 84.3% of cases for West Nile virus infection, respectively). The highest number of deaths was associated with listeriosis (N = 196; 13.7%), followed by salmonellosis (N = 71; 0.18%) and STEC infections (N = 18; 0.41%). For West Nile infections, the number of deaths was lower (N = 11), but the fatality rate was higher (7.2%) than salmonellosis and STEC infections.



With regard to foodborne outbreaks (FBOs), *Salmonella*, Norovirus and *Campylobacter* accounted for the highest number of outbreaks and cases.



Data on congenital toxoplasmosis are not shown since 2021 data are not available yet.

Note: The total number of confirmed cases is indicated in parentheses at the end of each bar.

(a) Regarding West Nile virus infection, the total number of locally acquired cases was used (includes probable and confirmed cases).

**Figure 1:** Reported numbers of cases and notification rates for confirmed human zoonoses in the EU, 2021

Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.v



Table 2: Reported hospitalisations and deaths due to zoonoses in confirmed human cases and among foodborne outbreak cases in the EU, 2021

		9	Survei	illance data	on hun	nan cas	es (sour	ce: EC	DC)			Foodborne	Foodborne outbreaks (FBO) (source: EFSA)						
		Hospitalisation					Deaths												
Disease	Confirmed human cases	Status available		Reporting MSs <sup>(b)</sup>	Cases and proportion of hospitalised cases		Outcome available		Reporting MSs <sup>(b)</sup>	Deaths and case fatality		Outbreaks	Cases	Hospitalisations and proportion of hospitalised cases					
	N	N	%	N	N	%	N	%	N	N	%	N	N	N	%	N	%		
Campylobacteriosis	127,840	45,121	35.3	15	10,469	23.2	91,177	71.3	16	26	0.03	249	1,051	134	12.7	6	0.6		
Salmonellosis	60,050	30,951	51.5	16	11,785	38.1	38,658	64.4	16	71	0.18	773	6,755	1,123	16.6	1	0.1		
Yersiniosis	6,789	1,564	23.0	13	508	32.5	3,596	53.0	21	0	0	21	125	14	11.2	0	0		
STEC infections	6,084	2,133	35.1	17	901	42.2	4,366	71.8	20	18	0.41	31	275	47	13.5	0	0		
Listeriosis	2,183	956	43.8	16	923	96.5	1,427	65.4	14	196	13.7	23	104	48	46.2	12	11.5		
Tularaemia	876	221	25.2	10	112	50.7	341	38.9	11	2	0.59	0	0	0	_	0	_		
Echinococcosis	529	121	22.9	13	51	42.1	270	51.0	16	0	0	0	0	0	_	0	_		
Q fever	460	NA	NA	NA	NA	NA	270	58.7	11	4	1.5	0	0	0	_	0	_		
Brucellosis	162	60	37.0	10	36	60.0	59	36.4	11	0	0	1	2	2	100	0	_		
West Nile virus infection <sup>(a)</sup>	152	83	54.6	6	70	84.3	152	100	8	11	7.2	NA	NA	NA	NA	NA	NA		
Trichinellosis	77	26	33.8	6	10	38.5	28	36.4	6	0	0	1	2	0	100	0	_		
Rabies	0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA		

Data on congenital toxoplasmosis are not shown since 2021 data are not available yet.

MSs: Member States.

NA: Not applicable as the information is not collected for this dsease.

(a): Total number of locally acquired infections (probable and confirmed cases).

(b): Not all countries observed cases for all diseases.



# Comparison of human zoonoses data for 2021, 2020 and previous years

In order to estimate the trends of human infections, especially in relation to the COVID-19 pandemic, the 2021 data (number of cases and notification rates) were compared with those from 2020 as well as with the mean annual number of cases and mean notification rates from the prepandemic years (2017–2019). Moreover, some statistics were also collected to obtain indications concerning the effects of the withdrawal of the United Kingdom from the EU. The absolute and relative difference between the number of cases and the notification rates reported in the EU for 2021 compared with 2020, and for 2021 compared with the annual average for each disease in the 2017–2019 period, were estimated (Table 3).

For all zoonoses except echinococcosis, Q fever, WNV infection and trichinellosis, there was an increase in the number of confirmed cases for 2021 when compared to 2020, whereas the comparison in terms of the number of confirmed cases for 2021 vs. the annual mean for the 2017–2019 period showed a decrease in 2021 for all zoonoses except tularaemia.

For STEC infections ( $\pm$ 36.9%), tularaemia ( $\pm$ 33.3%), salmonellosis ( $\pm$ 14.3%), listeriosis ( $\pm$ 14.0%), brucellosis ( $\pm$ 12.3%), yersiniosis ( $\pm$ 11.8%) and campylobacteriosis ( $\pm$ 2.1%), there was an increase in the relative difference in notification rates ( $\pm$ 100,000 population) in 2021 as compared with 2020, whereas for all the other zoonoses, the opposite trend was observed. The same comparison (relative difference) between the 2021 notification rates and the annual averages of the 2017–2019 notification rates, conducted to evaluate the effect of the pandemic, revealed a sharp decrease in WNV infections ( $\pm$ 76.4%), brucellosis ( $\pm$ 54.8%), Q fever ( $\pm$ 8.8%), campylobacteriosis ( $\pm$ 8.6%), trichinellosis ( $\pm$ 8.6%), echinococcosis ( $\pm$ 23.0%) and salmonellosis ( $\pm$ 9.9%) and listeriosis ( $\pm$ 4.3%). For each disease, the relative difference in EU notification rates was also calculated based on EU-27 data only (i.e. excluding data reported by the United Kingdom for 2017, 2018 and 2019) (Table 3) in order to provide evidence of the effect of the withdrawal of the United Kingdom from the EU. The comparison of the relative difference in notification rates based on EU and EU-27 data showed the same trend for almost all of the zoonoses with very few exceptions (e.g. yersiniosis).

**Table 3:** Number of confirmed human cases and notification rates (per 100,000 population) in 2021 by zoonosis, including the absolute and relative (%) difference with regard to 2020, and the mean annual cases and mean notification rates (per 100,000 population) for the 2017–2019 period, EU

Listeriosis			Cases (N	)	Notific	cation rates	(N confirm	ned cases <sup>;</sup>	*100,000)					
Zoonosis	EU		2020	2017-		20	20	2017–2019 (mean annual)						
200110313	level <sup>(a)</sup>	2021	2020 difference	2019 (mean annual)	2021	Absolute difference	Relative difference (%)	Absolute difference	Relative difference (%)					
Campylobac-	EU	127,840	+7,296	-110,076	41.1	+0.86	+2.1	-23.2	-36.1					
teriosis	EU-27			<b>-47,665</b>				-16.1	-28.1					
Salmonellosis	EU	60,050	+7,360	-30,401	15.7	+1.96	+14.3	-3.8	-19.6					
	EU-27	,		-20,638				-4.7	-23.1					
Yersiniosis	EU	6,789	6,789	6,789	6,789	6,789	6,789	+1,128	-153	1.9	+0.20	+11.8	+0.19	+11.3
	EU-27			+15				-0.19	-8.9					
STEC infections	EU	6,084	6,084	6,084	+1,595	-1,262	2.1	+0.55	+36.9	+0.19	+9.9			
	EU-27			+211				+0.26	+14.2					
Listeriosis	EU	2,183	+296	-363	0.49	+0.06	+14.0	+0.02	+4.3					
	EU-27			-203				-0.01	-2.0					
Tularaemia	EU	876	+235	+252	0.20	+0.05	+33.3	+0.08	+66.7					
	EU-27			+252				+0.06	+42.9					
Echinococcosis	EU	529	-15	-283	0.15	-0.01	<b>−7.5</b>	-0.04	-23.0					
	EU-27			-280				-0.06	-30.4					



18314732, 2022, 12, Downloaded from https://drs.onlinelibrary.wiley.com/doi/10.2903j.drs.2022.7666 by Istituto Zooprofilation Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (thtps://onlinelibrary.wiley.com/terms

			Cases (N	)	Notific	cation rates	(N confirm	ned cases	*100,000)													
Zoonosis	EU			2017-		20	20	2017-2019 (mean annual)														
Zoonosis	level <sup>(a)</sup>	2021	2020 difference	2019 (mean annual)	2021	Absolute difference	Relative difference (%)	Absolute difference	Relative difference (%)													
Q-fever	EU	460	-63	<b>-415</b>	0.11	-0.01	-12.0	-0.07	-38.8													
	EU-27			-399				-0.09	-45.8													
West Nile	EU	152	-178	-573	0.03	-0.04	-53.9	-0.11	<b>−76.4</b>													
virus <sup>(b)</sup>	EU-27			-573				-0.13	− <b>79.5</b>													
Brucellosis	EU	162	+30	-178	0.03	< 0.01	+12.3	-0.04	-54.8													
	EU-27																	-170				-0.04
Tuberculosis	EU	111	111	+12	-71	0.03	< 0.01	+12.2	-0.01	-28.3												
caused by <i>M. bovis,</i> <i>M caprae</i>	EU-27			-38				-0.01	-25.3													
Trichinellosis	EU	77	<del>-4</del> 0	-33	0.02	-0.01	-32.5	<-0.01	-17.6													
	EU-27			-33				-0.01	-28.6													
Rabies	EU	0	0	-2	0	0	_	0	_													
	EU-27			-2				0	_													

<sup>(</sup>a): Data from the United Kingdom are taken into account for 2017–2019 because the United Kingdom was an EU MS, but it became a third country on 1 February 2020. To calculate the difference between 2021 and 2017–2020 (mean annual statistics), data from the United Kingdom for 2017–2019 were included in the 'EU Total' difference, whereas human data from the United Kingdom were not collected by ECDC for 2020 and 2021 ('EU-27').

<sup>(</sup>b): For West Nile virus infection, the total number of locally acquired cases was used (including probable and confirmed cases).



//efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico

Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms


and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

# Zoonoses included in compulsory annual monitoring (in accordance with Directive 2003/99 List A)

# 1. Campylobacter

Summary data substantiating this chapter, as well as additional information on related projects and internet sources are published for this report on the EFSA Knowledge Junction on Zenodo here. Summary statistics on *Campylobacter* food monitoring data and on campylobacteriosis foodborne outbreaks reported in the framework of Directive 2003/99/EC, with downloadable files, are retrievable using the EFSA *Campylobacter* dashboard and the EFSA foodborne outbreaks dashboard, respectively available here and here. Summary statistics on human surveillance data with downloadable files are retrievable using the ECDC Surveillance Atlas of Infectious Diseases here.


# Campylobacter in the EU, 2021



# Human cases in foodborne outbreaks



# Foodborne outbreaks





# 1.1. Key facts

- Campylobacteriosis is the most commonly reported foodborne gastrointestinal infection in humans in the EU and has been so since 2007.
- In 2021, the number of confirmed cases of human campylobacteriosis was 127,840, corresponding to an EU notification rate of 41.1 per 100,000 population. This was an increase of 2.1% compared with 2020 (40.2 per 100,000 population).
- In 2020, ECDC recorded the lowest number of human cases of campylobacteriosis in any year since surveillance began in 2007, owing to the impact of the COVID-19 pandemic and the withdrawal of the United Kingdom from the EU.
- Compared with the rate before the COVID-19 pandemic (2017–2019 annual mean), the 2021 EU notification rate decreased by 36.1% and 28.1%, with and without the data from the United Kingdom, respectively. The overall trend for the campylobacteriosis in the period 2017–2021 in the EU showed no statistically significant increase or decrease.
- In most of the cases (99.1%), where the origin was known, the infection was acquired in the EU.
- Twenty-four MSs reported data in 2021 in the context of the *Campylobacter* process hygiene criterion, set out in Regulation (EC) No 2073/2005. Fifteen MSs reported 8,063 test results from official controls, with 42.1% *Campylobacter*-positive samples and 18.4% exceeding the limit of 1,000 CFU/g. Twenty MSs reported 53,351 test results from the monitoring of food business operators, with 27.2% positive samples and 16.4% exceeding the limit of 1,000 CFU/g. Eleven MSs reported results from both samplers, showing that the number of samples exceeding the limit was significantly higher in official samples (19.4%) than in own-checks (7.3%).
- In 2021, 3,220 'ready-to-eat' and 14,158 'non-ready-to-eat' results from food sampling units were reported by 12 and 16 MSs, respectively. In the former category, 0.31% of the units were positive to *Campylobacter* as compared to 10.9% in the latter. The highest level of contamination in the 'non-ready-to-eat' food category was in 'meat and meat products' with 11.9% of positive units. Overall, *Campylobacter* was isolated from all fresh meat categories, with meat from turkeys and broilers showing the highest percentage of *Campylobacter*-positive sampling units at 12.9% and 11.5%, respectively.
- Campylobacter spp. was detected by 16 MSs and three non-MSs in more than 30 different animal categories in 2021. The vast majority of units tested in the EU (N = 10,162) were from broilers with a proportion of positives of 10.5%. The proportion of positive sampling units for pigs, cattle and 'cats and dogs' was 41.3%, 13.5% and 12.3%, respectively.
- Alongside the present report, EFSA has also published two new interactive communication tools on *Campylobacter*: the EFSA story map (available here) and the dashboard (available here).

# 1.2. Surveillance and monitoring of *Campylobacter* in the EU

## **1.2.1.** Humans

In 2021, 27 EU MSs reported information on campylobacteriosis in humans. Surveillance of campylobacteriosis is mandatory in 22 EU MSs. In five MSs (Belgium, France, Greece, Italy and the Netherlands), notification is based on a voluntary system. In 2020–2021, Spain did not receive data from all the regions that normally report, so the case number might not be complete. The EU case definition was used by 21 MSs. Four MSs used a different case definition for reporting (Denmark, France, Germany and Italy), Greece did not use any case definition and the Netherlands did not specify which case definition it used. All MSs, except four (Belgium, Greece, Italy and the Netherlands) had a comprehensive surveillance system.

The campylobacteriosis surveillance systems cover the whole population in all MSs except four (France, Italy, the Netherlands and Spain). The estimated coverage of the surveillance system is 20% in France and 64% in the Netherlands. These estimated proportions of population coverage were used in the calculation of notification rates for these two MSs. No estimates of population coverage in Italy and Spain were provided, so no notification rates were calculated for these two MSs. All countries reported case-based data except Belgium, Bulgaria and Greece, which reported aggregated data.



## 1.2.2. Food and animals

*Campylobacter* is monitored along the food chain during the primary production stage (farm animals), during harvest/slaughter, manufacturing and at the distribution stage.

# Campylobacter data in the context of Regulation (EC) No 2073/2005

A regulatory limit (microbiological process hygiene criterion (PHC)) of 1,000 CFU/g of *Campylobacter* on the neck skins of chilled broiler carcases was set by Regulation (EC) N. 2073/2005<sup>8</sup> (point 2.1.9 of Chapter 2 of Annex I). This limit applies to a set of 50 pooled samples from 10 consecutive sampling sessions. As of 2021, a maximum number of 15 samples with values exceeding this limit are considered as acceptable. Food business operators (FBOp) failing to comply with this limit are required to carry out corrective actions involving validation and verification of their food safety management procedures based on HACCP principles and good manufacturing practices (GMP). The PHC has been in force since 1 January 2018. On 14 December 2019, Commission Implementing Regulation (EU) 2019/627<sup>9</sup> came into force, harmonising sampling procedures for official controls. This legislation requires the Competent Authority (CA) to verify whether the FBOp is correctly implementing and checking the PHC on broiler carcases by choosing one of two approaches: implementing ad hoc official sampling or collecting all the information from the samples taken by the FBOp. The results obtained in official controls enable improved trend watching and trend analyses (Table 1).

# Other monitoring data for food and animals

Campylobacter monitoring data from food and animals submitted to EFSA in compliance with Chapter II 'Monitoring of zoonoses and zoonotic agents' of the Zoonoses Directive 2003/99/EC<sup>11</sup> are collected without a harmonised procedure. These data allow descriptive summaries at EU level, but they do not support EU-level trend analyses and trend watching (Table 1).

In 2021, general data on food and animals reported to EFSA by MSs and non-MSs were obtained mainly from official sampling, industry sampling HACCP and own-checks, as part of national monitoring and surveillance, and/or organised surveys. In addition, for animal data, other reported samples were obtained from clinical investigations by private veterinarians and industry (e.g. artificial insemination centres).

The occurrence of *Campylobacter* reported in the main food categories for the year 2021 and for the 4-year period of 2017–2020 was descriptively summarised, making a distinction between RTE and non-RTE food. Data sets were extracted using the strategy of 'objective sampling', meaning that the reporting MS collected the samples as part of a planned strategy based on the selection of random samples that are statistically representative of the population to be analysed.

Other *Campylobacter* monitoring data, intended solely for monitoring antimicrobial resistance, are obtained from selected animal species and their carcases/meat, using the harmonised sampling scheme set out in the Commission Implementing Decision (EU) 2020/1729. These antimicrobial resistance results are published in a separate EU Summary Report.

The detection and confirmation of *Campylobacter* in food and animals is generally based on culture, with the use of international standards or equivalent validated methods. Species identification is carried out using biochemical and molecular methods (PCR based), as well as matrix-assisted laser desorption/ionisation time-of flight mass spectrometry (MALDI-TOF MS).

EFSA Journal 2022;20(12):7666

<sup>&</sup>lt;sup>8</sup> Commission Regulation (EC) No 2073/2005 of 15 November 2005 on microbiological criteria for foodstuffs OJ L 338, 22.12.2005, pp. 1–26.

<sup>&</sup>lt;sup>9</sup> Commission Implementing Regulation (EU) 2019/627 of 15 March 2019, laying down uniform practical arrangements for the performance of official controls on products of animal origin intended for human consumption in accordance with Regulation (EU) 2017/625 of the European Parliament and of the Council and amending Commission Regulation (EC) No 2074/2005 regarding official controls. OJ L 131, 17.5.2019, pp. 51–100.

This means official sampling using the same method and sampling area as food business operators. At least 49 random samples shall be taken in each slaughterhouse each year. The number of samples may be reduced in small slaughterhouses and based on a risk evaluation.

Directive 2003/99/EC of the European Parliament and of the Council of 17 November 2003 on the monitoring of zoonoses and zoonotic agents, amending Council Decision 90/424/EEC and repealing Council Directive 92/117/EEC. OJ L 325, 12.12.2003, pp. 31–40.

18314732, 2022, 12, Downloaded from https://fs.aonlinelibrary.wiley.com/doi/10.2033/jcfsa.202272666 by Istituto Zooperfilation Sperimental dell'Umbria e delle Marche (ZSUM), Wiley Online Library on [18122023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Cereive Common

## EFSA story map on Campylobacter

The EFSA story map on *Campylobacter* is a new interactive communication tool developed by EFSA in 2022, available online (here) and geared to the general public. This story map provides general information on the pathogen and its epidemiology, including information on where the pathogen can be found, how people and animals get infected, the occurrence of this pathogen in different sources, the disease it causes and how to prevent infection. In addition, this story map also illustrates the monitoring activities implemented in the EU and the role of EFSA with respect to these activities. Users can easily display and explore the content of the different sections in the story map, browsing the dynamic maps, images, text and multimedia features.

## 1.3. Data analyses

A comparison was made of *Campylobacter* results exceeding 1,000 CFU/g from the neck skins of broiler carcases after chilling, as obtained by the CA and FBOp as part of the *Campylobacter* PHC in compliance with Regulation (EC) No 2073/2005. The significance of any differences was verified by the one-tailed Fisher's exact probability test in cases where the expected values of any of the cells in a contingency table were below 5; otherwise, the one-tailed z test was used. The official control sampling results by the CA and the own-check results by the FBOp were expressed as prevalence ratios with an exact binomial confidence interval of 95%. A p-value < 0.10 (Clayton and Hills, 2013) was considered as significant, in order to highlight every possible indication of differences between the data collected by the FBOp and the CA. R software (www.r-project.org, version 4.0.5) was used to conduct the above analyses.

# EFSA dashboard on Campylobacter

The EFSA dashboard on *Campylobacter* (available online here) is a graphical user interface for searching and querying the large amount of data collected each year by EFSA from EU MSs and other reporting countries based on Zoonoses Directive 2003/99/EC. The *Campylobacter* dashboard shows summary statistics for the monitoring results of the pathogen with regard to major food categories, *Campylobacter*-positive official samples exceeding the Process Hygiene Criterion limit of 1,000 CFU/g for chilled broiler carcases and the occurrence of *Campylobacter* in major food categories. The *Campylobacter* data and related statistics can be displayed interactively using charts, graphs and maps in the online EFSA dashboard. In this tool, the main statistics can also be viewed and downloaded in tabular format. Detailed information on the use and features of the *Campylobacter* dashboard can be found in the user guide available on Zenodo here and can also be downloaded from the online tool. Links to the dashboard are available in the relevant sections of this chapter.

## 1.4. Results

# 1.4.1. Overview of key statistics, EU, 2017–2021

Table 4 summarises EU statistics on human campylobacteriosis and on the occurrence and prevalence of *Campylobacter* in food and animals, respectively, during 2017–2021. In 2021, a slight increase was observed in notified human cases and the EU notification rate, despite the ongoing COVID-19 pandemic. However, the notification rate remained low compared with the pre-pandemic period of 2017–2019. The food data of interest in this report were classified into two major categories: 'meat and meat products' and 'milk and milk products', aggregated by year to obtain an annual overview of the volume of data submitted. Since 2019, the number of sampling units reported for 'meat and meat products' has increased sharply, probably owing to the Commission Implementing Regulation (EU) 2019/627 establishing compulsory reporting of *Campylobacter* PHC monitoring data (see above). Animal data were collected principally from broilers, cattle and pigs, showing comparable and consistent test numbers over the considered period. More detailed descriptions of these statistics are provided in the below subsections and in the chapter on foodborne outbreaks.

//efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico

dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms

**Table 4:** Summary of *Campylobacter* statistics related to humans, major food categories and the main animal species, EU, 2017–2021

	2021 <sup>(a)</sup>	2020	2019 <sup>(b)</sup>	2018 <sup>(b)</sup>	2017 <sup>(b)</sup>	Data source
Humans						
Total number of confirmed cases	127,840	120,544	220,639	246,570	246,538	ECDC
Total number of confirmed cases/100,000 population (notification rates)	41.1	40.2	60.6	66.0	66.1	ECDC
Number of reporting MSs	27	27	28	28	28	ECDC
Infection acquired in the EU	81,311	70,772	109,937	116,246	122,280	ECDC
Infection acquired outside the EU	704	1,586	6,514	7,685	6,583	ECDC
Unknown travel status or unknown country of infection	45,825	48,186	104,188	122,639	117,675	ECDC
Number of foodborne outbreak-related cases	1,051	1,319	1,770	2,365	3,608	EFSA
Total number of foodborne outbreaks	249	317	542	537	395	EFSA
Food <sup>(c)</sup>						
Meat and meat products <sup>(d)</sup>						
Number of sampling units	87,808	66,099	57,027	26,514	21,521	EFSA
Number of reporting MSs	25	25	25	26	22	EFSA
Milk and milk products(e)						
Number of sampling units	2,125	2,145	2,749	3,227	2,317	EFSA
Number of reporting MSs	11	11	11	13	13	EFSA
Animals						
Cattle (bovine animals)						
Number of sampling units <sup>(f)</sup>	7,529	4,387	6,850	4,220	7,312	EFSA
Number of reporting MS	11	7	10	8	10	EFSA
Gallus gallus (chicken)						
Number of sampling units <sup>(f)</sup>	10,162	13,628	10,472	14,093	10,133	EFSA
Number of reporting MSs	6	15	8	16	7	EFSA
Pigs						
Number of sampling units <sup>(f)</sup>	4,502	2,110	4,308	2,481	3,824	EFSA
Number of reporting MSs	14	4	11	5	10	EFSA

ECDC: European Centre for Disease Prevention and Control; EFSA: European Food Safety Authority; MSs: Member States.

- (a): Data on food and animal samples from the United Kingdom (Northern Ireland) are taken into account for 2021. In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/Northern Ireland, the EU requirements on data sampling are also applicable to Northern Ireland.
- (b): Data from the United Kingdom are taken into account for 2017–2019, because the United Kingdom was an EU MS, but it became a third country on 1 February 2020.
- (c): Summary statistics referring to MSs were obtained by totalling all sampling units (single samples, samples from batches and samples from slaughter batches), sampling stages (farm, packing centre, processing plant, cutting plant, slaughterhouse, catering, hospital or medical care facility, restaurant or cafe or pub or bar or hotel or catering service, retail, wholesale, border control posts, school or kindergarten, unspecified), sampling strategies (census, convenience sampling, selective sampling, objective sampling and unspecified) and samplers (official sampling, official and industry sampling, private sampling, unspecified).
- (d): 'Meat and meat products' refer to carcases and fresh meat/ready-to-eat (RTE), cooked and fermented products.
- (e): 'Milk and milk products' refer to raw and pasteurised milk and all dairy products including cheeses.
- (f): Summary statistics referring to MSs were obtained by totalling all sampling units (single animals, slaughter animal batches and herds or flocks).

For a further interactive look at *Campylobacter* monitoring results, dashboards have been implemented (*different filters can be applied to query the data*) (*link*).

## 1.4.2. Human campylobacteriosis

In 2021, 127,840 confirmed cases of human campylobacteriosis were reported by the 27 EU MSs, corresponding to an EU notification rate of 41.1 cases per 100,000 population (Table 4). This is an increase of 2.1% compared with 2020 (40.2 per 100,000 population) and a decrease of 36.1% and

28.1% compared with the average annual notification rate from 2017 to 2019 (pre-pandemic period), with and without United Kingdom data, respectively (Table 3). France, Ireland, Latvia, Malta and Portugal had an increased rate in 2021 compared with the pre-pandemic period. The highest country-specific notification rates in 2021 were observed in Czechia (152.4 cases per 100,000), Slovakia (111.7), Luxembourg (92.8) and Malta (73.2). The lowest rates in 2021 were observed in Poland, Romania, Bulgaria, Greece and Cyprus ( $\leq$  2.7 per 100,000) (Table 5).

For most (99.1%) of the reported campylobacteriosis cases of known origin, the infection was contracted in the EU (Table 4) as compared with 98.5% in 2020. Twenty countries reported data on the importation of cases. The proportion of domestic cases with known data was over 95% in all reporting countries except five MSs, which reported the highest proportion of travel-associated cases: Finland (34.2%), Sweden (19.1%), Denmark (12.7%), Portugal (6%) and Germany (5.1%). The proportion of travel-associated cases observed in 2021 (3.8%) compared with 2020 (3.7%) were very much at the same level, differing from the higher proportion of travel-associated cases in 2019 (10.8%), before the COVID-19 pandemic. However, the number of cases acquired outside the EU decreased considerably in 2021 compared with 2020 (Table 4). Of the 3,138 travel-associated cases reported by MSs with a known country of origin, 2,063 cases (65.7%) were linked to travel within the EU, with most of the infections being acquired in Spain, Croatia, Greece and Italy (28.1%, 12.6%, 10.5% and 9.5%, respectively). Turkey, Serbia, Bosnia and Herzegovina, Pakistan and Ukraine were the most frequently reported probable countries of infection outside the EU (29.8%, 7.7%, 7%, 3.1% and 3.1%, respectively). Campylobacteriosis cases were reported in all age groups, with the highest proportion of reported cases belonging to the youngest age group from 0 to 4 years (20,884 cases: 16.8%).

**Table 5:** Reported confirmed human cases of campylobacteriosis and notification rates per 100,000 population in EU MSs and non-MS countries, by country and year, 2017–2021

		2021	_		202	:0	201	.9	201	В	201	.7
Country	National coverage <sup>(a)</sup>	Data	Confirme and ra		Confirme and ra		Confirme		Confirmed and ra		Confirme	
	coverage	iormat	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate
Austria	Υ	С	6,019	67.4	5,406	60.7	6,572	74.2	7,999	90.7	7,204	82.1
Belgium	Υ	Α	3,273	28.3	5,693	49.4	7,337	64.0	8,086	70.9	8,649	76.2
Bulgaria	Υ	Α	130	1.9	127	1.8	229	3.3	191	2.7	195	2.7
Croatia	Υ	С	1,173	29.1	1,054	26.0	1,722	42.2	1,965	47.9	1,686	40.6
Cyprus	Υ	С	24	2.7	18	2.0	21	2.4	26	3.0	20	2.3
Czechia	Υ	С	16,305	152.4	17,517	163.8	22,894	215.0	22,895	215.8	24,326	230.0
Denmark	Υ	С	3,740	64.0	3,742	64.3	5,402	93.0	4,559	78.9	4,255	74.0
Estonia	Υ	С	185	13.9	265	19.9	347	26.2	411	31.2	285	21.7
Finland	Υ	С	1,798	32.5	2,074	37.5	4,382	79.4	5,099	92.5	4,289	77.9
France <sup>(b)</sup>	N	С	8,875	65.6	7,920	58.8	7,712	57.4	7,491	55.9	6,579	49.2
Germany	Υ	С	47,912	57.6	46,378	55.8	61,277	73.8	67,585	81.6	69,251	83.9
Greece	Υ	Α	260	2.4	218	2.0	366	3.4	357	3.3	344	3.2
Hungary	Υ	С	5,088	52.3	4,461	45.7	6,400	65.5	7,117	72.8	7,807	79.7
Ireland	Υ	С	3,147	62.9	2,419	48.7	2,776	56.6	3,044	63.0	2,779	58.1
Italy <sup>(c)</sup>	N	С	1,542	_	1,418	_	1,633	_	1,356	_	1,060	_
Latvia	Υ	С	158	8.3	104	5.5	133	6.9	87	4.5	59	3.0
Lithuania	Υ	С	357	12.8	684	24.5	1,221	43.7	919	32.7	990	34.8
Luxembourg	Υ	С	589	92.8	729	116.4	271	44.1	625	103.8	613	103.8
Malta	Υ	С	378	73.2	206	40.0	278	56.3	333	70.0	231	50.2
Netherlands <sup>(d)</sup>	N	С	2,692	24.1	2,549	25.2	3,415	34.1	3,091	34.6	2,890	32.5
Poland	Υ	С	616	1.6	414	1.1	715	1.9	719	1.9	874	2.3
Portugal	Y	С	973	9.4	790	7.7	887	8.6	610	5.9	596	5.8
Romania	Y	С	348	1.8	300	1.6	805	4.1	573	2.9	467	2.4
Slovakia	Υ	С	6,099	111.7	4,921	90.2	7,690	141.1	8,339	153.2	6,946	127.8




18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10/2903j\_cfsa.0222.7666 by Isitiuo Zooprofilatios Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [8/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

Slovenia Spain <sup>(c),(e)</sup> Sweden <b>EU Total 27</b> United Kingdom <b>EU Total</b> <sup>(f)</sup>		2021	L		202	0	201	9	2018	3	201	7
Country	National	Data	Confirmed and ra		Confirme			Confirmed cases and rates and rates		Confirmed cases and rates		
	coverage <sup>(a)</sup>	format <sup>(a)</sup>	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate
Slovenia	Y	С	856	40.6	811	38.7	1,085	52.1	1,305	63.1	1,408	68.2
Spain <sup>(c),(e)</sup>	N	С	11,244	_	6,891	_	9,658	_	18,410	_	18,860	_
Sweden	Υ	С	4,059	39.1	3,435	33.3	6,693	65.4	8,132	80.4	10,608	106.1
EU Total 27	-	_	127,840	41.1	120,544	40.2	161,921	54.0	181,324	58.3	183,271	59.0
United Kingdom	_	_	_	_	_	_	58,718	88.1	65,246	98.4	63,267	96.1
EU Total <sup>(f)</sup>	-	-	127,840	41.1	120,544	40.2	220,639	60.6	246,570	66.0	246,538	66.1
Iceland	Υ	С	58	15.7	95	26.1	136	38.1	145	41.6	119	35.2
Norway	Y	С	2,049	38.0	2,422	45.1	4,154	78.0	3,668	69.3	3,883	73.8
Liechtenstein	Υ	С	38	97.3	6,196	71.7	7,200	83.9	7,696	90.3	7,217	85.3
Switzerland <sup>(g)</sup>	Υ	С	6,755	77.9								

- -: Data not reported.
- (a): Y: yes; N: no; A: aggregated data; C: case-based data.
- (b): Sentinel surveillance; notification rates calculated with an estimated coverage of 20%.
- (c): The notification rate was not calculated since information on estimated coverage was not available.
- (d): Sentinel surveillance; notification rates calculated with an estimated coverage of 64% in 2021, 58% in 2019–2020, 52% in 2017–2018.
- (e): Data not complete in 2020–2021, rate not estimated.
- (f): Cases reported by the United Kingdom for the period 2017–2019 were also considered for this estimation (EU-28). When the United Kingdom data were collected for the period 2017–2019, the United Kingdom was an EU MS, but it became a third country on 1 February 2020.
- (g): Switzerland provided data directly to EFSA. The human data for Switzerland include data from Liechtenstein for the years 2017–2020.

Between 2012 and 2021, the number of confirmed campylobacteriosis cases reported in the EU showed a clear seasonal trend, peaking in the summer months. Annual winter peaks were also observed in January from 2012 to 2021, although peak numbers were lower than those observed during the summer. As in 2020, a decrease in the number of cases was observed in 2021, probably due to the COVID-19 pandemic. However, the overall campylobacteriosis trend in 2017–2021 showed no statistically significant increase or decrease (Figure 2). Belgium, Finland, Hungary, Poland, Slovenia and Sweden reported significantly decreasing trends (p < 0.05) during the period 2017–2021. Latvia and Portugal reported a significantly increasing trend over the same period.



Source: Austria, Cyprus, Czechia, Denmark, Estonia, Finland, France, Germany, Hungary, Ireland, Italy, Latvia, Luxembourg, Malta, the Netherlands, Poland, Romania, Slovenia and Sweden.

**Figure 2:** Trends in reported confirmed human cases of campylobacteriosis in the EU, by month, 2017–2021

Information on hospitalisation status was provided for 35.3% of all campylobacteriosis cases by 15 MSs. Of the cases with known hospitalisation status, 10,469 (23.2%) were hospitalised. The highest hospitalisation rates were reported in Latvia (95.6%), Cyprus (91.7%) and Poland (80%), where most of the reported cases were hospitalised. Outcomes were reported for 71.3% of all cases by 16 MSs. Twenty-six deaths from campylobacteriosis were reported in 2021, resulting in an EU case fatality rate of 0.03%. The average percentage of fatal outcomes observed has remained unchanged over the past 5 years (range 0.03–0.05%). Information on gender was provided for 123,990 confirmed cases in the EU, 54.7% were male and 45.3% female.

Campylobacter species information was provided by 22 MSs for 65.1% of confirmed cases reported in the EU, representing a slight increase compared with 2020 (64.7%). Of these cases, 88.4% were Campylobacter jejuni, 10.1% C. coli, 0.18% C. fetus, 0.12% C. upsaliensis and 0.09% Campylobacter lari. Other Campylobacter species accounted for 1.1% of cases, but most of those cases were reported at the national level as 'C. jejuni/C. coli/C. lari not differentiated'. Belgium, Bulgaria, Denmark, Greece and Sweden provided no information on species.

# 1.4.3. Campylobacter in food

# Campylobacter data in the context of Regulation (EC) No 2073/2005

The *Campylobacter* PHC monitoring data reported in Table 6 show test results obtained using a culture-based enumeration method (ISO 10272-2:2017), from the neck skins of chilled broiler carcases sampled at slaughterhouses within the EU. Fifteen MSs reported ad hoc official sampling results, 20 MSs reported monitoring results from FBOp and 11 MSs reported data from both samplers. In total, MSs reported 8,936 (31%) *Campylobacter*-positive units out of 28,823 neck skin units tested.

From ad hoc official sampling of 8,063 units, 1,486 (18.4%) exceeded the limit of 1,000 CFU/g. Moderate variability was observed in percentage test results exceeding the limit. In particular, Latvia reported none, six MSs (Bulgaria, Estonia, Hungary, Ireland, Netherlands, Romania) reported fewer

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/continelibrary.wiley.com/confinelibrary.wiley.c



18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903j.efsa.2022.7666 by Istituo Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

than 8.3% of units exceeding the limit, while eight MSs (Belgium, Croatia, Cyprus, Germany, Greece, Italy, Poland, Spain) reported a higher number of units above the limit, ranging from 14.1% to 51.4%. The number of *Campylobacter*-positive units, reported by 13 MSs totalled 3,119 (42.1%).

FBOp reported test results for 53,351 neck skin samples from own-check sampling activities, of which 8,759 (16.4%) exceeded the limit of 1,000 CFU/g. For four MSs (Estonia, Finland, Romania, Slovakia), the levels of positives exceeding the limit ranged from 0 to 0.41%. In Switzerland and Montenegro, 77 of 805 (9.5%) and 2 of 50 (4%) units exceeded the limit, respectively. The number of *Campylobacter*-positive units reported by 15 MSs totalled 5,817 (27.2%).

Eleven MSs reported results from both samplers and the overall total number of units exceeding the limit was significantly higher in official samples (19.4%, N = 1,090) than in own-checks (7.3%, N = 1,773). A far higher percentage of units exceeding the limit was also observed in official samples compared with FBOp in eight MSs (Belgium, Estonia, Germany, Greece, Italy, Poland, Romania, Spain). A substantial difference in percentage *Campylobacter*-positive units was also observed, with results for official controls (N = 2,263;45.6%) being higher than own-checks (N = 1,443;15.8%).

Overall, for the *Campylobacter* PHC monitoring data provided by 24 MSs, the percentage of units exceeding the limit was significantly higher in official samples (18.4%) than in FBOp samples (16.4%).

For a further interactive look at *Campylobacter* monitoring results, dashboards have been implemented (different filters can be applied to query the data) (link).



Comparison of proportions (%) of *Campylobacter*-positive single samples and samples exceeding the *Campylobacter* PHC limit collected from the neck skins of chilled broiler carcases sampled at the slaughterhouses in accordance with Regulation (EC) No. 2073/2005, by sampler and reporting MS, EU, 2021

		Competen	t Authority (C	A)	F	ood busines	s operator (FI	3Op)		
Country	N samples Tested	N (%) samples positive	N (%) samples above 1,000 CFU/g	CI <sub>95</sub> samples above 1,000 CFU/g	N samples Tested	N (%) samples positive	N (%) samples above 1,000 CFU/g	CI <sub>95</sub> samples above 1,000 CFU/g	p-value <sup>(b),(c)</sup>	Interpretation <sup>(c)</sup>
Austria	_	_	_	_	957	NA	61 (6.4)	[4.9; 8.1]	_	_
Belgium	633	NA	89 (14.1)	[11.4; 17.0]	2,421	NA	172 (7.1)	[6.1; 8.2]	< 0.001	CA > FBOp
Bulgaria	1,048	218 (20.8)	16 (1.5)	[0.88; 2.5]	_	-	_	_	_	_
Croatia	832	417 (50.1)	253 (30.4)	[27.3; 33.7]	_	_	_	_	_	_
Cyprus	220	180 (81.8)	113 (51.4)	[44.6; 58.1]	_	-	_	_	_	_
Czechia	_	_	_	_	4,110	2,573 (62.6)	1,620 (39.4)	[37.9; 40.9]	_	_
Denmark	_	_	_	_	1,150	164 (14.3)	86 (7.5)	[6,0; 9.2]	_	_
Estonia	12	1 (8.3)	1 (8.3)	[0.21; 38.5]	260	0	0	[0; 1.4] <sup>(a)</sup>	0.04	CA > FBOp
Finland	_	_	_	_	585	1 (0.17) <sup>(f)</sup>	1 (0.17)	[0; 0.95]	_	_
France	_	_	_	_	16,357	NA	4,389 (26.8)	[26.2; 27.5]	_	_
Germany	28	NA	9 (32.1)	[15.9; 52.4]	6,604	NA	510 (7.7)	[7.1; 8.4]	< 0.001	CA > FBOp
Greece	75	52 (69.3)	33 (44.0)	[32.5; 55.9]	612	31 (5.1)	31 (5.1)	[3.5; 7.1]	< 0.001	CA > FBOp
Hungary	344	41 (11.9)	14 (4.1)	[2.2; 6.7]	_	-	_	_	_	_
Ireland	164	96 (58.5)	10 (6.1)	[3; 10.9]	1,031	379 (36.8)	75 (7.3)	[5.8; 9.0]	NS	
Italy	1,233	639 (51.8)	310 (25.1)	[22.7; 27.7]	5,591	NA	466 (8.3)	[7.6; 9.1]	< 0.001	CA > FBOp
Latvia	100	4 (4.0)	0 (0)	[0; 3.6] <sup>(a)</sup>	434	90 (20.7)	24 (5.5)	[3.6; 8.1]	0.01	CA < FBOp
Netherlands	333	79 (23.7)	10 (3.0)	[1.4; 5.5]	3,336	201 (6.0)	201 (6.0)	[5.2; 6.9]	0.0239	CA < FBOp
Poland	885	287 (32.4)	174 (19.7)	[17.1; 22.4]	1,365	112 (8.2)	109 (8.0)	[6.6; 9.6]	< 0.001	CA > FBOp
Portugal	_	_	_	_	3,528	1,006 (28.5)	521 (14.8)	[13.6; 16.0]	_	_
Romania	1,399	521 (37.2)	84 (6.0)	[4.8; 7.4]	1,450	491 (33.9)	6 (0.41)	[0.15; 0.90]	< 0.001	CA > FBOp
Slovakia	_	_	_	_	1,075	20 (1.9)	0	[0; 0.34] <sup>(a)</sup>	_	_
Slovenia	_	_	_	_	804	595 (74.0)	333 (41.4)	[38; 44.9]	_	_
Spain	757	584 (77.1)	370 (48.9)	[45.3; 52.5]	635	139 (21.9)	139 (21.9)	[18.7; 25.3]	< 0.001	CA > FBOp
Sweden	_	_	_	_	1,046	15 (1.4)	15 (1.4)	[0.8; 2.4]	_	_
EU Total (27 + XI)	8,063	3,119 (42.1) <sup>(d)</sup>	1,486 (18.4)	[17.6; 19.3]	53,351	5,817 (27.2) <sup>(e)</sup>	8,759 (16.4)	[16.1; 16.7]	< 0.001	CA > FBOp



		Competent Authority (CA)				ood busines	ss operator (Fl	3Op)		
Country	N samples Tested	N (%) samples positive	N (%) samples above 1,000 CFU/g	CI <sub>95</sub> samples above 1,000 CFU/g	N samples Tested	N (%) samples positive	N (%) samples above 1,000 CFU/g	CI <sub>95</sub> samples above 1,000 CFU/g	p-value <sup>(b),(c)</sup>	Interpretation <sup>(c)</sup>
EU Total (27 + XI) providing CA and FBOp data	5,619	2,263 (45.6) <sup>(d)</sup>	1,090 (19.4)	[18.4; 20.5]	23,739	1,443 (15.8) <sup>(e)</sup>	1,733 (7.3)	[7.0; 7.6]	< 0.001	CA > FBOp

NA: not available.

(a): One-sided, 97.5% confidence interval.

(b): p-value: NS, not significant.

(c): Related to the percentage of positive samples above 1,000 CFU/g.

(d): Belgium and Germany did not report Campylobacter-positive samples below 1,000 CFU/g from test results of the Competent Authority.

(e): Austria, Belgium, France, Germany and Italy did not report Campylobacter-positive samples below 1,000 CFU/g from test results of food business operators.

(f): Reporting error. Finland indicated, during the last phase of the production of this report, that the number of Campylobacter-positive samples below 1,000 CFU/g from the FBOp was unknown.

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903j.efsa.2022.7666 by Istituo Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

## Other food monitoring data

Table 7 summarises the reported occurrence of *Campylobacter* in the main food categories in 2021 and over the 4-year period of 2017–2020 within the EU. A distinction is made between RTE and non-RTE food, and fresh meat.

The proportion of *Campylobacter*-positive samples in the RTE and non-RTE categories was 0.31% and 10.9%, respectively. In fresh meat, 11.7% of sampling units were positive.

In 2021, most of the results from the 3,220 RTE food sampling units reported by 12 MSs came from 'fruit, vegetables and juices' (37.7%), followed by 'milk and milk products' (28.2%) and 'meat and meat products' (13.1%). In total, *Campylobacter* was detected in 10 RTE food samples: one from 'raw milk', one from 'meat products', one from 'ready-to-eat salads', three from meat-based dishes reported as 'other processed food products and prepared dishes' and four from oysters reported as 'other foods'. During the period 2017–2020, the percentage of *Campylobacter*-positive sampling units in RTE food was low, at below 1% for all categories. Over the entire period for the main RTE categories, the highest percentage of *Campylobacter*-positive units was for 'raw milk': 10 positives out of 1,036 (0.97%) sample units tested.

The results reported in 2021 by 16 MSs for non-RTE food show that 'meat and meat products' was the most contaminated food category, followed by 'other food' and 'milk and milk products'. During 2017–2020, 'meat and meat products' was the most contaminated food category, followed by 'milk and milk products'.

Thirteen MSs reported results for fresh meat categories. The percentage of *Campylobacter*-positive units was similar for meat from turkey (12.9%) and broilers (11.5%), and higher for 'other fresh meat' (17.7%). The percentage for fresh meat from pigs and bovine animals remained relatively low; 2.5% and 0.52%, respectively.

In 2021, a substantial decrease in the number of positive units was observed for non-RTE food and fresh meat, compared with the period 2017–2020, except for 'other fresh meat', for which the positive percentage was comparable.

**Table 7:** Occurrence of *Campylobacter* in the main food categories, EU, 2021 and 2017–2020

		2021		2	2017–2020 <sup>(a</sup>	)
Food	N reporting MSs	N sampled units	Positive N (%)	N reporting MSs	N sampled units	Positive N (%)
RTE food						
All	12	3,220	10 (0.31)	15	11,429	19 (0.17)
Meat and meat products	7	421	1 (0.24)	10	1,301	3 (0.23)
Meat and meat products from broilers	2	6	1 (16.7)	3	39	0
Meat and meat products from turkeys	0	0	0	2	15	1 (6.7)
Other meat and meat products	5	415	0	9	1,247	2 (0.16)
Milk and milk products	7	909	1 (0.11)	6	3,029	10 (0.33)
Milk	3	361	1 (0.28)	6	1,055	10 (0.95)
Raw milk <sup>(b)</sup>	2	212	1 (0.47)	6	1,036	10 (0.97)
Cheese	5	546	0	7	1,959	0
Dairy products excluding cheeses, butter, cream, ice cream, whey, yoghurt and fermented dairy products	1	2	0	3	15	0
Fruits, vegetables and juices	5	1,215	0	5	2,530	1 (0.04)
Ready-to-eat salads	2	367	1 (0.27)	5	640	1 (0.16)
Other processed food products and prepared dishes	4	239	3 (1.3)	6	3,275	4 (0.12)
Other foods	5	69	4 (5.8)	6	654	0



		2021		2	2017–2020 <sup>(a</sup>	a)
Food	N reporting MSs	N sampled units	Positive N (%)	N reporting MSs	N sampled units	Positive N (%)
Non-RTE food						
All	16	14,158	1,543 (10.9)	22	73,990	17,431 (23.6)
Meat and meat products	15	12,704	1,511 (11.9)	22	66,947	17,353 (25.9)
Meat and meat products from broilers	14	10,287	1,219 (11.8)	20	38,220	12,666 (33.1)
Meat and meat products from turkeys	9	610	85 (13.9)	10	3,876	1,058 (27.3)
Other meat and meat products	11	1,807	207 (11.5)	18	24,851	3,629 (14.6)
Milk and milk products	5	394	1 (0.25)	8	2,556	51 (2)
Fruits, vegetables and juices	4	337	0	6	1,500	2 (0.13)
Other food	6	723	31 (4.3)	8	2,987	25 (0.84)
Fresh meat <sup>(c)</sup>						
All	13	11,783	1,381 (11.7)	22	61,805	16,061 (26.0)
Fresh meat from broilers	13	9,845	1,135 (11.5)	20	35,858	11,857 (33.1)
Fresh meat from turkeys	8	583	75 (12.9)	10	3,419	957 (28.0)
Fresh meat from pigs	2	239	6 (2.5)	10	2,571	110 (4.3)
Fresh meat from bovine animals	3	192	1 (0.52)	12	4,174	40 (0.96)
Other fresh meat	8	924	164 (17.7)	13	15,783	3,097 (19.6)

RTE: ready-to-eat.

For a further interactive look at *Campylobacter* monitoring results, dashboards have been implemented (*different filters can be applied to query the data*) (link).

## 1.4.4. *Campylobacter* in animals

Table 8 shows the number of positive *Campylobacter* spp. samples detected during 2021 in the five main animal species, as well as in the 'other animals' category containing more than 30 different animal groups. In total, 16 MSs and three non-MSs reported data, primarily relating to broilers (39.6%, N = 12,733), followed by cattle (24.6%, N = 7,923) and pigs (15.2%, N = 4,877). The overall proportion of positive units in EU was highest in pigs (41.3%) followed by cattle (13.5%), cats and dogs (12.3%) and finally broilers (10.5%). Although fewer samples were tested for 'other animals', a considerable proportion of positive units were detected in samples from birds collected by MS (25.1%, N = 131).

**Table 8:** Summary of *Campylobacter* statistics related to major animal species, reporting EU MSs and non-MS countries, 2021

		EU MSs			P	Non-MS countrie	ries			
Animals	N reporting countries	N tested <sup>(a)</sup> sampling units	Positive sampling units				N reporting countries	N tested <sup>(a)</sup> sampling units	nits unit	pling
	,pg		N	%			N	%		
Gallus gallus (broilers)	6	10,162	1,065	10.5	2	2,571	117	4.6		
Bovine animals <sup>(b)</sup>	11	7,529	1,015	13.5	2	394	168	42.6		

Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.c

<sup>(</sup>a): Data from the United Kingdom are taken into account for 2017–2019, because the United Kingdom was an EU MS, but it became a third country on 1 February 2020.

<sup>(</sup>b): Raw RTE milk sampling units are a subset of RTE milk.

<sup>(</sup>c): Fresh meat sampling units are a subset of the two main categories above.

		EU MSs			ı	Non-MS countrie	s	
	N reporting countries	N tested <sup>(a)</sup> sampling units	Positive sampling units		N reporting countries	N tested <sup>(a)</sup> sampling units	Positive sampling units	
			N	%			N	%
Pigs	13	4,428	1,827	41.3	2	449	338	75.3
Small ruminants	6	2,934	109	3.7	1	21	5	23.8
Cats and dogs	3	913	112	12.3	2	1,581	95	6.0
Other animals <sup>(c)</sup>	4	876	143	16.3	2	284	12	4.2
Total	16	26,842	4,271	15.9	3	5,300	735	13.9

MS: Member State.

- (a): Summary statistics were obtained by totalling all sampling units (single samples, batch samples, animals, slaughter animal batches and herds or flocks).
- (b): Animals from the sampling stage 'Artificial insemination stations' were not included.
- (c): Other animals include Alpacas, Antelopes, Badgers, Barbary sheep, Bears, Birds, Budgerigars, Camels, Canary, Cantabrian chamois, Chinchillas, Deer, Elephants, Falcons, Ferrets, Foxes, Geese, Guinea pigs, Hedgehogs, Kangaroos, Land game mammals, Leporidae, Lions, Llamas, Martens, Monkeys, Oscine birds, Other animals, Other carnivores, Parrots, Pigeons, Raccoons, Rats, Reindeer, Reptiles, Rodents, Sea lions, Snakes, Solipeds, domestic Steinbock, Turkeys, Turtles, Water buffalos, Wild boars, Wild cats (Felis silvestris), Wild ducks, Zoo animals, all.

## 1.5. Discussion

Campylobacteriosis has been the most frequently reported zoonosis in humans across the EU since 2007. Despite comprehensive surveillance and national coverage in most MSs, the number of reported cases is underestimated in the EU (Teunis et al., 2013). In 2019, in two-thirds of EU MSs, the number of confirmed campylobacteriosis cases decreased. A decrease in the number of cases was also observed in 2020, probably due to the withdrawal of the United Kingdom from the EU and the COVID-19 pandemic, which had an impact on surveillance, including diagnosis and reporting. In 2021, although the COVID-19 pandemic was still ongoing, a slight increase was observed in the number of confirmed campylobacteriosis cases compared with 2020. The gradual scaling down of national public response measures to COVID-19 in the EU in 2021, <sup>12</sup> along with the return of mass gatherings, sporting and recreational/social events and the reopening of bars, restaurants and catering facilities (i.e. schools, workplaces) may explain this slight increase. However, the overall campylobacteriosis trend in 2017–2021 showed no statistically significant increase or decrease.

Compared with 2020, no difference in travel-associated campylobacteriosis cases was observed. The number of travel-associated cases remained below pre-pandemic level. European and Mediterranean countries were the main places of origin for travel-associated campylobacteriosis. *Campylobacter* has a characteristic seasonality with cases increasing sharply in the summer. Campylobacteriosis cases have been positively associated with temperature and, to a lesser degree, precipitation (Lake et al., 2019). Recent studies have even evidenced a possible association between campylobacteriosis and global climate change (Kuhn et al., 2020).

A smaller but distinct winter peak in the EU has become apparent in the past 10 years, including in 2021. Disease onsets concerning cases that were notified during the winter peaks occurred predominantly in January. This points to an exposure around the Christmas/New Year period. Reports indicate that meat fondues or table-top grilling, which are popular during the festive season, could promote the transmission of *Campylobacter* in some countries, causing the winter peak (Bless et al., 2017; Rosner et al., 2021).

In 2021, more than 10,000 cases of campylobacteriosis were hospitalised, making it the second most common foodborne agent associated with the number of hospitalisations, after salmonellosis. The proportion of hospitalised campylobacteriosis cases was higher than expected in some MSs, where all or most of the confirmed cases were hospitalised. These MSs also reported the lowest notification rates, indicating that surveillance focuses primarily on hospitalised (i.e. severe) cases. This can lead to an

\_

18314732, 2022, 12, Downdoaded from https://disa.onlinelibitrary.wiley.com/doi/10/2033/gfsa2022/7666 by Istitute Zoperinented ell'Umbia e delle Marche (ZSUM), Wiley Online Library on [1812/2023], See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10/2033/gfsa2022/7666 by Istitute Zoperinented by the applicable Cereive Common

<sup>&</sup>lt;sup>12</sup> More information is available at: https://www.ecdc.europa.eu/en/publications-data/download-data-response-measures-covid-19



overestimation of the proportion of hospitalised cases in some countries. As in previous years, *C. jejuni* and *C. coli* were the main species notified by MSs in 2021. Unfortunately, there was still a high percentage (34.9%) of campylobacteriosis cases in which the *Campylobacter* species was not determined. Although the proportion of characterised isolates at species level has increased over the last 5 years, further improvement is still necessary. A One Health approach, using whole genome sequencing (WGS) for typing *Campylobacter* isolates in humans, food and animals, will allow for better species characterisation and will enhance the monitoring of zoonotic transmission, improving public health surveillance.

In 2021, as part of a food control strategy, 24 MSs submitted Campylobacter PHC data, with 11 reporting both official and own-check results, four only official results and nine only own-check results. Official control and FBOp monitoring data from 2021 showed that about one in five and one in six samples exceeded the limit of 1,000 CFU/q, respectively. For the MSs that submitted data from both samplers, the results exceeding the limit concerned one in five units for the CA and one in 14 for the FBOp, respectively. The results reported by the CA were always significantly higher than those reported by FBOp. Moreover, the CA also reported considerably higher percentages of Campylobacter positive units. This discrepancy, observed for the second year, deserves more thorough investigation in order to identify the critical parameters and factors explaining these differences. Monitoring Campylobacter for the purposes of improving biosecurity measures on farms is of paramount importance (Newell et al., 2011). The aim here is to stimulate action to lower Campylobacter counts on broiler carcases and to reduce the number of human campylobacteriosis cases caused by the consumption or handling of contaminated chicken/broiler meat. It was recently reported that a 3-log10 reduction in broiler caecal concentrations would reduce the relative risk within the EU of human campylobacteriosis attributable to broiler meat by a substantial 58% (EFSA BIOHAZ Panel et al., 2020a). Overall, the comparison with 2020 percentages of Campylobacter samples exceeding the limit revealed a slight increase and a slight decrease, respectively, for CA and FBOp. The number of samples reported and the number of reporting MSs increased compared to previous years, which can be explained by the need for MSs to comply with the EU regulation.

Monitoring of other food in the EU showed overall percentages of *Campylobacter*-positive units in RTE and non-RTE foods of 0.31% and 10.9%, respectively. Although the presence of *Campylobacter* in RTE was low and has remained stable over the years, these findings are of concern given that contaminated RTE products directly expose consumers to infection. The RTE foods found to be contaminated with *Campylobacter* were 'raw milk', 'meat and meat products', 'fruit, vegetables and juices' along with 'other food', including meat-based dishes. This confirms the importance of those categories over the period 2017–2020. Moreover, positive results were observed for four 'other foods', including oysters. Given the risk of contracting campylobacteriosis from raw shellfish (Teunis et al., 1997), efforts must be encouraged to increase the sampling frequency of these food products, eaten raw in some regions. Overall, a small number of sampling units could have resulted in an imprecise estimation of the prevalence of RTE food samples contaminated by *Campylobacter*.

Monitoring data for non-RTE food showed positive results for 1 in 10 'meat and meat products', and 1 in 400 'milk and milk products'. Although the *Campylobacter* contamination observed in fresh meat categories was moderate in 2021, a decrease in the number of positive units was observed compared with the period 2017–2020. Nevertheless, the data continue to underline the key role of these products in campylobacteriosis epidemiology, either through direct handling or through cross-contamination of other foods.

In 2021, 16 MSs and three non-MSs reported data from several animal groups. *Campylobacter* spp. was detected in all the major animal categories: broilers, pigs, cattle, small ruminants, cats and dogs. Broilers were tested most frequently and accounted for 24.5% of test results, followed by cattle and pigs. The highest percentage of positive units, however, was observed for pigs. Despite an increase in the number of tested units, only about half of MSs reported broiler data in 2021, thereby hampering a fair comparison of data and probably indicating the fluctuation in positive results as being associated with the testing of a different epidemiological scenario.




18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilattico

dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms


# 2. Salmonella

Summary data substantiating this chapter, as well as additional information on related projects and Internet sources are published for this report on the EFSA Knowledge Junction on Zenodo here. Summary statistics on *Salmonella* food and animal monitoring data and on salmonellosis foodborne outbreaks reported in the framework of Directive 2003/99/EC, with downloadable files, are retrievable using the EFSA *Salmonella* dashboard and the EFSA foodborne outbreaks dashboard, respectively available here and here. Summary statistics on human surveillance data with downloadable files are retrievable using the ECDC Surveillance Atlas of Infectious Diseases here.


# **Salmonella** in the EU, 2021



# **Human cases in foodborne outbreaks**



# Foodborne outbreaks





# 2.1. Key facts

- Salmonellosis was the second most commonly reported foodborne gastrointestinal infection in humans after campylobacteriosis and was a major cause of foodborne outbreaks in EU MSs and non-MS countries.
- In 2021, the number of confirmed cases of human salmonellosis was 60,050, corresponding to an EU notification rate of 15.7 per 100,000 population. This was an increase of 14.3% compared with the rate in 2020.
- In 2020, ECDC recorded the lowest number of human *Salmonella* cases since 2007, when salmonellosis surveillance started. The number of cases was impacted by the COVID-19 pandemic and the United Kingdom's withdrawal from the EU.
- Compared with the rate before the COVID-19 pandemic (2017–2019 annual mean), the 2021 EU notification rate decreased by 19.6% and 23.1%, with and without the data from the United Kingdom, respectively.
- Notwithstanding, the overall trend for salmonellosis in 2017–2021 did not show any statistically significant increase or decrease.
- The proportion of hospitalised cases was 38.1%, which was higher than in 2020, with an EU case fatality rate of 0.18%, which is similar to 2020.
- The top five *Salmonella* serovars involved in human infections overall were distributed as follows: S. Enteritidis (54.6%), S. Typhimurium (11.4%), monophasic S. Typhimurium (1,4, [5],12:i:-) (8.8%), S. Infantis (2.0%) and S. Derby (0.93%).
- Sampling to verify compliance with process hygiene criteria on carcases at the slaughterhouse found the highest proportions of positive samples among those collected by the competent authorities for broilers (14%), turkeys (7.4%), pigs (1.7%) and sheep (1.2%). Conversely, for goats, the highest proportion of positive samples was for those collected by food business operators (2.1%).
- In 2021, 73,238 'ready-to-eat' food sampling units were collected with a very low proportion of *Salmonella*-positive units (0.23%), overall. The highest proportions of positives were found for 'meat and meat products from pigs' (0.82%) and 'spices and herbs' (0.72%). For 'non-ready-to-eat' food, 466,290 sampling units were collected and the proportion of positive samples was low (2.1%). 'Meat and meat products' (2.2%) and especially those from broilers (4.4%) and turkeys (3.6%) were the food categories with the highest proportions of positive units.
- Sixteen MSs and the United Kingdom (Northern Ireland) reporting on *Salmonella* control programmes met the reduction targets for all poultry populations. The number of MSs that did not meet the reduction targets was seven for laying hens, five for breeding *Gallus gallus*, three for broilers, one for breeding turkeys and one for fattening turkeys.
- For flocks of broilers and of breeding and fattening turkeys, for which results of *Salmonella* control programmes are required to be reported separately by sampler, the prevalence reported by food business operators was significantly lower than that reported by competent authorities.
- A significant increase in the estimated breeding turkey flock prevalence of Salmonella was noted in 2021 compared with 2016, when the estimated prevalence reached the lowest value seen in the entire study period (2010–2021). Flock prevalence trends for target Salmonella serovars have, in contrast, been fairly stable over the last few years for all poultry populations.
- Considering the top 20 EU ranking of *Salmonella* serovars by food-animal source, focusing on those most frequently isolated from humans, *S.* Enteritidis was the most commonly reported serovar in laying hens and the second most commonly reported one in broilers. Amongst the top three for pig and bovine isolates were *S.* Typhimurium and its monophasic variant. Some other very common serovars were *S.* Infantis for broilers (one out of two isolates from this source belonged to this serovar) and *S.* Derby for pigs.
- With the present report, EFSA has also published two new interactive communication tools on *Salmonella*: the EFSA story map (available here) and the dashboard (available here).



# 2.2. Surveillance and monitoring of Salmonella in the EU

#### EFSA story map on Salmonella

The EFSA story map on *Salmonella* is a new interactive communication tool developed by EFSA in 2022, available online (here) and geared to the general public. This story map provides general information on the pathogen and its epidemiology, including information on where the pathogen can be found, how people and animals get infected, the occurrence of this pathogen in different sources, the disease it causes and how to prevent infection. In addition, this story map also illustrates the monitoring activities implemented in the EU and the role of EFSA with respect to these activities. Users can easily display and explore the content of the different sections in the story map, browsing the dynamic maps, images, text and multimedia features.

#### **2.2.1.** Humans

In 2021, 27 EU MSs reported information on non-typhoidal salmonellosis infections in humans. The notification of salmonellosis is mandatory in 24 EU MSs, whereas it is voluntary in three MSs (Belgium, France and the Netherlands). The EU case definition was used by 23 MSs, while four MSs (Denmark, France, Germany and Italy) reported using other case definitions. All countries except the Netherlands had a comprehensive surveillance system. The surveillance systems for salmonellosis covered the whole population in all MSs except three: France, the Netherlands and Spain. The estimated coverage of the surveillance system was 48% in France and 64% in the Netherlands. These estimated proportions of population coverage were used in the calculation of notification rates for these two MSs. No estimated population coverage in Spain was provided, so notification rates were not calculated. In 2020–2021, Spain did not receive data from all the regions that normally report, due to COVID-19, so the case numbers may therefore not be complete. All countries reported case-based data except Bulgaria, which reported aggregated data.

### 2.2.2. Food, animals and feed

# Salmonella data in the context of Regulation (EC) No 2073/2005

Regulation (EC) No 2073/2005 lays down microbiological criteria, intended as food safety criteria (FSC) and process hygiene criteria (PHC), for Salmonella in specific food categories. Compliance with these criteria must be legally verified by the individual food business operator (FBOp) as part of their own HACCP programme. In addition, the competent authority (CA), through official sampling or oversight of data, should ensure that the FBOp complies with these regulatory requirements. The Salmonella FSC require that the pathogen not be detected in different products when they are on the market, during their shelf-life. Moreover, in fresh poultry meat (from species covered by national control programmes), the FSC require the absence of target serovars (S. Enteritidis and S. Typhimurium including monophasic S. Typhimurium  $(1,4,[5],12:i:-)^{14}$ ). The Salmonella PHC are regulated for carcases of pigs, cattle, sheep, goats, horses, broilers and turkeys, as sampled by the FBOp. Moreover, in accordance with Regulation (EU) No 2019/627<sup>15</sup>, the CA has to verify whether the FBOp correctly implements and checks these PHC for carcases by choosing between different approaches: (i) implementing official sampling, (ii) collecting all information on the total number and the number of Salmonella-positive samples from own checks by the FBOp and/or (iii) collecting information on the total number and the number of Salmonella-positive samples as part of national control programmes in the MSs with special guarantees (Regulation (EC) No 853/2004<sup>16</sup>).

EFSA Journal 2022;20(12):7666

<sup>&</sup>lt;sup>13</sup> Regulation (EC) No 852/2004 of the European Parliament and of the Council of 29 April 2004 on the hygiene of foodstuffs. OJ L 139, 30.4.2004, pp. 1–54.

<sup>&</sup>lt;sup>14</sup> Commission Regulation (EU) No 1086/2011 of 27 October 2011 amending Annex II to Regulation (EC) No 2160/2003 of the European Parliament and of the Council and Annex I to Commission Regulation (EC) No 2073/2005 as regards salmonella in fresh poultry meat Text with EEA relevance. OJ L 281, 28.10.2011, pp. 7–11.

<sup>&</sup>lt;sup>15</sup> Commission Implementing Regulation (EU) 2019/627 of 15 March 2019 laying down uniform practical arrangements for the performance of official controls on products of animal origin intended for human consumption in accordance with Regulation (EU) 2017/625 of the European Parliament and of the Council and amending Commission Regulation (EC) No 2074/2005 as regards official controls. OJ L 131, 17.5.2019, pp. 51–100.

Regulation (EC) No 853/2004 of the European Parliament and of the Council of 29 April 2004 laying down specific hygiene rules for food of animal origin. OJ L 139, 30.4.2004, pp. 55–205.



# Data for compliance with national control programmes for Salmonella in poultry populations

In accordance with Regulation (EC) No 2160/2003<sup>17</sup> and its subsequent amendments, MSs have to set up national control programmes (NCPs) for *Salmonella* aimed at reducing the prevalence of *Salmonella* serovars that are considered relevant for public health (from this point forward, termed 'target serovars'). Currently, prevalence targets have been defined for breeding flocks of *Gallus gallus*, laying hens, broilers and breeding and fattening turkeys and correspond to the maximum annual percentage of flocks positive for *S*. Enteritidis and *S*. Typhimurium, including its monophasic variant, except for breeding flocks of *Gallus gallus*, where *S*. Infantis, *S*. Virchow and *S*. Hadar are considered to be relevant as well. The prevalence target is equal to 1% or less for breeding flocks of *Gallus gallus*, <sup>18</sup> broilers<sup>19</sup> and breeding and fattening turkeys<sup>20</sup>; it is 2% for laying hens.<sup>21</sup> Every year, MSs must report results for their *Salmonella* NCP and, for broiler flocks and breeding and fattening turkey flocks, results for sampling conducted by the CA and FBOp must also be reported separately.

# Other monitoring data for food, animals and feed including serovars

Food, animal and feed data other than those described above are not collected in a harmonised way, because there are no specific legal requirements. The reported occurrence of *Salmonella* in the main food categories was descriptively summarised, with a distinction being made between 'ready-to-eat (RTE) and non-RTE food'. Data sets were extracted with 'objective sampling' being specified as the sampling strategy, which means that samples were representative of the population being analysed and were collected according to a planned strategy.

The occurrence of *Salmonella* in animal populations was descriptively summarised considering all data collected in different sampling contexts and reported as different sampling units (e.g. 'holding', 'herd/flock', 'animal' and 'slaughter animal batch'), with the exception of data related to poultry populations covered by NCPs, which have been discussed separately.

The reported data on *Salmonella* serovars were also descriptively summarised. MSs are required to report the target serovars as part of their NCPs for poultry, whereas for the samples collected in different contexts, serotyping is not mandatory. Also, for the food sector, the FSC is the absence of *Salmonella*, except for fresh poultry meat, for which the criterion is the absence of target serovars. The compulsory reporting of target serovars in some contexts (NCPs for poultry and FSC for fresh poultry meat) guarantees the consistency of such data over the years and among MSs but could result in these target serovars being overestimated compared with the other serovars.

# 2.3. Data analyses

# 2.3.1. Comparison between competent authority and food business operator sampling results

CA and FBOp Salmonella results in the context of NCPs for those poultry populations requiring separate reporting (i.e. broilers and fattening and breeding turkeys) were compared, as were Salmonella PHC monitoring data from carcases (of pigs, cattle, goats, sheep, horses, broilers and turkeys). The significance of differences was verified by the one-tailed Fisher's exact probability test, in cases where the expected values in any of the cells of a contingency table were below five; otherwise, the z-statistic one-tailed test was performed. CA control sampling results and the own-check results of the FBOp were expressed as prevalence and exact binomial confidence interval (95% level). A p-value

EFSA Journal 2022;20(12):7666

<sup>&</sup>lt;sup>17</sup> Regulation (EC) No 2160/2003 of the European Parliament and of the Council of 17 November 2003 on the control of salmonella and other specified foodborne zoonotic agents. OJ L 325, 12.12.2003, pp. 1–15.

Commission Regulation (EU) No 200/2010 of 10 March 2010 implementing Regulation (EC) No 2160/2003 of the European Parliament and of the Council as regards a Union target for the reduction of the prevalence of Salmonella serotypes in adult breeding flocks of Gallus (Text with EEA relevance). OJ L 61, 11.3.2010, pp. 1–9.

<sup>19</sup> Commission Regulation (EU) No 200/2012 of 8 March 2012 concerning a Union target for the reduction of Salmonella Enteritidis and Salmonella Typhimurium in flocks of broilers, as provided for in Regulation (EC) No 2160/2003 of the European Parliament and of the Council Text with EEA relevance. OJ L 71, 9.3.2012, pp. 31–36.

Commission Regulation (EU) No 1190/2012 of 12 December 2012 concerning a Union target for the reduction of Salmonella Enteritidis and Salmonella Typhimurium in flocks of turkeys, as provided for in Regulation (EC) No 2160/2003 of the European Parliament and of the Council Text with EEA relevance. OJ L 340, 13.12.2012, p. 29–34.

<sup>&</sup>lt;sup>21</sup> Commission Regulation (EU) No 517/2011 of 25 May 2011 implementing Regulation (EC) No 2160/2003 of the European Parliament and of the Council as regards a Union target for the reduction of the prevalence of certain *Salmonella* serotypes in laying hens of *Gallus* and amending Regulation (EC) No 2160/2003 and Commission Regulation (EU) No 200/2010. OJ L 138, 26.5.2011, pp. 45–51.

< 0.10 (Clayton and Hills, 2013) was considered significant to include all possible evidence of differences between data collected by the FBOp and CA.

R software (www.r-project.org, version 4.2.1) was used to conduct the above-mentioned analyses.

# 2.3.2. Statistical trend analyses for poultry monitoring data

Statistical trend analyses were carried out with the objective of evaluating the significance of temporal variations in the EU-level flock prevalence of *Salmonella* and target *Salmonella* serovars in poultry since the start of NCP implementation.

The tested flocks were either positive or negative for target serovars and *Salmonella*, and so the status of the flocks is a dichotomous outcome variable. Therefore, the binomial probability distribution for the response variable was assumed and the logit link function was computed in the model for the trend analysis. The logit is defined as the logarithm of p/(1-p), where p/(1-p) is the odds of being positive for *Salmonella*.

According to temporal flock prevalence trends in the MSs, B-spline basic models for the logit of the probability of flocks being positive were fitted for the different poultry populations over the entire period of NCP implementation. Moreover, attention was paid to the period after achievement of the minimum prevalence reported to date, to capture any evidence of a significant increase in *Salmonella* prevalence. Marginal and conditional generalised linear models for repeated measures were used to perform these trend analyses (EFSA, 2009a, 2011). Details about the estimated parameters of the models, odds ratios, prevalence rates and graphical analyses (conditional and marginal) are published in the file '*Salmonella* poultry outcome trend analyses' on the EFSA Knowledge Junction on Zenodo here.

To investigate EU-level prevalence, considering the relevant heterogeneity among MSs for the flock prevalence of *Salmonella* and target serovars over time, the results obtained using the conditional generalised mixed model for longitudinal binary data were summarised and discussed in the report, for all poultry populations covered by NCPs. To take into account the different levels (baselines) of probability of MSs having positive flocks, yet with similar patterns over time, a random MS-specific intercept effect was included in the model. To evaluate the trend over time, the 'time' variable was included in the model as a fixed effect. The correlation between repeated observations in the same MS in subsequent years was considered using a first autoregressive or exchangeable structure of the correlation matrix for the residuals. To evaluate the significance of the overall effect of fixed factors specified in the model, Type III F-tests were applied, whereas the receiver operating characteristic (ROC) curve was used to assess the goodness of fit of the model. A p-value < 0.10 was considered to be significant for both random and fixed effects.

GLIMMIX and SGPLOT procedures in SAS 9.4 software were used to fit the models and to produce the graphical outputs, respectively.

# 2.3.3. Descriptive analyses of *Salmonella* serovars

With the aims of evaluating the distribution of *Salmonella* serovars across the food chain and identifying potential sources of human infections, a Sankey diagram was provided to link the food and food-producing animal sources to the five most commonly reported *Salmonella* serovars from human cases acquired within the EU (domestically or during travel within the EU). For animal categories covered by NCPs, only serovar data reported in the context of these programmes were presented. For cattle, meat-producing animals were considered, whereas for pigs, data from fattening animals were used. In addition to possible reporting biases as regards serovars, reporting for animal or food categories may also have been unbalanced and specific sources (e.g. cattle) may have been underrepresented.

18314732, 2022, 12, Downloaded from https://fs.aonlinelibrary.wiley.com/doi/10.9293/j-fs.a.2022.7566 by Istituto Zoporofilatios Sperimenta dell'Umbria e delle Marche (ZSUM), Wiley Online Library on [1812023], See the Terms and Conditions (https://onlinelibrary.wiley.com/rerms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Cerawise Common.

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilattico

Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

#### EFSA dashboard on Salmonella

The EFSA dashboard on *Salmonella* (available online here) is a graphical user interface for searching and querying the large amount of data collected each year by EFSA from EU MSs and other reporting countries based on Zoonoses Directive 2003/99/EC. The *Salmonella* dashboard shows summary statistics for the monitoring results of the pathogen with regard to major food categories and animal species, *Salmonella*-positive official samples in the context of FSC and HPC, the occurrence of *Salmonella* in major food categories and the achievement of *Salmonella* reduction targets in poultry populations. The *Salmonella* data and related statistics can be displayed interactively using charts, graphs and maps in the online EFSA dashboard. In this tool, the main statistics can also be viewed and downloaded in tabular format. Detailed information on the use and features of the *Salmonella* dashboard can be found in the user guide available on Zenodo here and can also be downloaded from the online tool. Links to the dashboard are available in the relevant sections of this chapter.

# 2.4. Results

# 2.4.1. Overview of key statistics, EU, 2017–2021

#### **Humans**

In total, the number of reported human salmonellosis cases and the notification rate in 2021 were higher than in 2020 (Table 9). The number of reported human salmonellosis cases acquired in the EU (i.e. by domestic infection and through travel within the EU), the number of outbreak-related cases and the total number of foodborne salmonellosis outbreaks were higher in 2021 than in 2020, but lower than in previous years.

**Table 9:** Summary of *Salmonella* statistics related to humans, major food categories and the main animal species, EU, 2017–2021

	2021 <sup>(a)</sup>	2020	2019 <sup>(b)</sup>	2018 <sup>(b)</sup>	2017 <sup>(b)</sup>	Data source
Humans						
Total number of confirmed cases	60,050	52,690	87,908	91,858	91,587	ECDC
Total number of confirmed cases/100,000 population (notification rate)	15.7	13.7	19.5	19.6	19.4	ECDC
Number of reporting MSs	27	27	28	28	28	ECDC
Infection acquired in the EU	43,720	38,247	58,157	59,763	59,642	ECDC
Infection acquired outside the EU	925	973	6,343	6,376	6,001	ECDC
Unknown travel status or unknown country of infection	15,405	13,470	23,408	25,719	25,944	ECDC
Number of outbreak-related cases	6,755	3,686	10,240	11,631	9,607	EFSA
Total number of outbreaks	773	694	1,284	1,588	1,241	EFSA
Food						
Meat and meat products						
Number of sampling units	977,434	557,341	552,590	433,197	380,000	EFSA
Number of reporting MSs	28	26	28	28	28	EFSA
Milk and milk products						
Number of sampling units	43,902	38,492	46,797	44,078	30,796	EFSA
Number of reporting MSs	25	24	25	24	24	EFSA
Fish and fishery products						
Number of sampling units	14,882	16,486	13,974	17,075	13,486	EFSA
Number of reporting MSs	25	23	24	22	22	EFSA
Eggs and egg products						
Number of sampling units	14,696	11,579	12,093	10,611	15,435	EFSA
Number of reporting MSs	22	18	21	21	23	EFSA



18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (thps://onlinelibrary.wiley.com/terms

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

	2021 <sup>(a)</sup>	2020	2019 <sup>(b)</sup>	2018 <sup>(b)</sup>	2017 <sup>(b)</sup>	Data source
Fruit and vegetables (including juice)						
Number of sampling units	12,483	17,222	17,068	10,888	7,579	EFSA
Number of reporting MSs	23	23	22	22	25	EFSA
Animals						
Gallus gallus (chickens)						
Number of sampling units <sup>(c)</sup>	812,163	620,141	752,172	720,717	736,534	EFSA
Number of reporting MSs	28	26	27	27	28	EFSA
Turkeys						
Number of sampling units <sup>(c)</sup>	70,869	63,473	65,950	68,009	74,739	EFSA
Number of reporting MSs	25	22	23	24	26	EFSA
Ducks and geese						
Number of sampling units <sup>(c)</sup>	3,751	412	8,700	9,846	5,743	EFSA
Number of reporting MSs	10	6	9	6	8	EFSA
Pigs						
Number of sampling units <sup>(c)</sup>	16,689	17,234	18,619	17,868	19,239	EFSA
Number of reporting MSs	15	10	14	14	17	EFSA
Cattle (bovine animals)						
Number of sampling units <sup>(c)</sup>	26,061	28,363	86,871	30,302	654,593	EFSA
Number of reporting MSs	14	11	14	14	15	EFSA

ECDC: European Centre for Disease Prevention and Control; EFSA: European Food Safety Authority; MSs: Member States.

- (a): Data on food and animal samples from the United Kingdom (Northern Ireland) are taken into account for 2021. In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/Northern Ireland, the EU requirements on data sampling are also applicable to Northern Ireland.
- (b): Data from the United Kingdom are taken into account for 2017–2019, because the United Kingdom was an EU MS, but it became a third country on 1 February 2020.
- (c): Summary statistics referring to MSs were obtained by totalling all sampling units (single animals, slaughter animal batches and herds or flocks).

#### **Food categories**

The number of sampling units reported in 2021 for three of the five food categories was higher than in 2020, and for all food categories, there was an increase in the number of reporting MSs except for 'fruit and vegetables (including juice)'. The largest increase in the number of reported sampling units was for 'meat and meat products'. Conversely, for 'fruit and vegetables (including juice)' and 'fish and fishery products', there was a decrease in the number of sampled units reported in 2021 compared to 2020.

# **Animal categories**

For all animal categories, there was an increase in the number of reporting MSs. In general, there was an increase in the number of sampling units between 2020 and 2021, except for 'cattle' and 'pigs', for which the number was fairly stable between the 2 years.

More detailed descriptions of the above statistics are provided in the below subsections and in the chapter on foodborne outbreaks.

For a further interactive look at *Salmonella* monitoring results, dashboards have been created (different filters can be applied to query the data) (link).

#### 2.4.2. Human salmonellosis

In total, 60,050 human salmonellosis cases were reported by 27 MSs in 2021, corresponding to an EU notification rate of 15.7 cases per 100,000 population (Table 10). This was an increase of 14.3% compared with the rate in 2020 and a decrease of 19.6% and 23.1% compared with the rates in prepandemic years (2021 vs. 2017–2019) with and without the data from the United Kingdom, respectively.

As in the previous year, the highest notification rates in 2021 were reported by Czechia (93.7 cases per 100,000 population) and Slovakia (81.3 cases per 100,000 population), while the lowest rates were

reported by Bulgaria, Cyprus, Greece, Ireland, Romania and Portugal ( $\leq$  4.6 cases per 100,000 population).

The proportion of domestic vs. travel-associated cases varied markedly between countries, but most of the confirmed salmonellosis cases were acquired in the EU (72.8%), whereas 1.5% reported travel outside the EU and 25.7% of infections were of unknown origin (Table 9). Considering all cases in EU MSs, the highest proportions of domestic cases (100%) were reported by Croatia, Hungary, Malta, Poland, Romania, Slovakia and Spain. The highest proportions of travel-associated cases were reported by four countries: France (21%), Sweden (20%), Slovenia (19%) and Ireland (16%). Of 1,443 travel-associated cases with known information on the probable country of infection, 64.1% involved travel outside the EU. Turkey, Egypt, Morocco and Kosovo were the most frequently reported travel destinations outside the EU (21.2%, 7.1%, 6.9% and 3.7%, respectively). For travel-associated cases in the EU, the most common countries of infection were Spain and Italy.

**Table 10:** Reported confirmed human cases of salmonellosis and notification rates per 100,000 population in EU MSs and non-MS countries, by country and year, 2017–2021

		2021			202	0	201	.9	201	.8	201	7
Country	National coverage <sup>(a)</sup>	Data format <sup>(a)</sup>	Confir cases rate	and	Confir cases rate	and	Confir cases rate	and	Confirmed cases and rates		Confirmed cases and rates	
			Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate
Austria	Υ	С	993	11.1	817	9.2	1,866	21.1	1,538	17.4	1,667	19.0
Belgium	Υ	С	2,084	18.0	1,595	13.8	2,527	22.1	2,958	26.0	2,298	20.2
Bulgaria	Υ	Α	241	3.5	187	2.7	594	8.5	586	8.3	796	11.2
Croatia	Υ	С	605	15.0	786	19.4	1,308	32.1	1,323	32.2	1,242	29.9
Cyprus	Υ	С	41	4.6	70	7.9	62	7.1	44	5.1	59	6.9
Czechia	Υ	С	10,032	93.7	10,516	98.3	13,009	122.2	10,901	102.7	11,473	108.5
Denmark	Υ	С	692	11.8	614	10.5	1119	19.3	1168	20.2	1067	18.6
Estonia	Υ	С	112	8.4	91	6.8	150	11.3	314	23.8	265	20.1
Finland	Υ	С	474	8.6	516	9.3	1,175	21.3	1,431	26.0	1,535	27.9
France <sup>(b)</sup>	N	С	9,315	28.7	7,071	21.9	8,935	27.7	8,936	27.8	7,993	24.9
Germany	Υ	С	8,144	9.8	8,664	10.4	13,495	16.3	13,293	16.1	14,051	17.0
Greece	Υ	С	284	2.7	381	3.6	643	6.0	640	6.0	672	6.2
Hungary	Υ	С	3,298	33.9	2,964	30.3	4,452	45.6	4,161	42.6	3,922	40.0
Ireland	Υ	С	173	3.5	214	4.3	347	7.1	352	7.3	379	7.9
Italy	Υ	С	3768	6.4	2713	4.5	3256	5.4	3635	6.0	3347	5.5
Latvia	Υ	С	218	11.5	296	15.5	438	22.8	409	21.1	225	11.5
Lithuania	Υ	С	281	10.1	419	15.0	736	26.3	779	27.7	1,005	35.3
Luxembourg	Υ	С	133	21.0	93	14.9	131	21.3	135	22.4	118	20.0
Malta	Υ	С	249	48.2	176	34.2	131	26.5	116	24.4	107	23.2
Netherlands <sup>(c)</sup>	N	С	862	7.7	695	6.2	1,197	10.8	1,061	9.6	954	8.7
Poland	Υ	С	7,702	20.4	5,192	13.7	8,373	22.0	9,064	23.9	8,921	23.5
Portugal	Υ	С	361	3.5	262	2.5	432	4.2	302	2.9	462	4.5
Romania	Υ	С	518	2.7	408	2.1	1,383	7.1	1,410	7.2	1,154	5.9
Slovakia	Υ	С	4,439	81.3	3,385	62.0	4,992	91.6	6,791	124.8	5,789	106.5
Slovenia	Υ	С	185	8.8	214	10.2	362	17.4	274	13.3	275	13.3
Spain <sup>(d),(e)</sup>	N	С	3,913	-	3,526	_	5,087	_	8,730	_	9,426	-
Sweden	Υ	С	933	9.0	825	8.0	1,990	19.5	2,041	20.2	2,280	22.8
EU Total 27			60,050	15.7	52,690	13.7	78,190	20.4	82,392	20.5	81,482	20.1
United Kingdom	-	-	-	_	-	_	9,718	14.6	9466	14.3	10,105	15.3
EU Total <sup>(f)</sup>			60,050	15.7	52,690	13.7	87,908	19.5	91,858	19.6	91,587	19.4
Iceland	Υ	С	54	14.6	32	8.8	50	14.0	63	18.1	64	18.9
Norway	Υ	С	389	7.2	441	8.2	1,092	20.5	961	18.1	992	18.9




'		2021					201	2019		.8	2017	
Country	National coverage <sup>(a)</sup>	Data format <sup>(a)</sup>	Confirmed cases and rates									
	_		Cases	Rate								
Liechtenstein	Υ	С	7	17.9	1,260	14.6	1,538	17.9	1,468	17.2	1,835	21.7
Switzerland <sup>(g)</sup>	Υ	С	1,480	17.1								

<sup>-:</sup> Data not reported.

A seasonal trend was observed for confirmed salmonellosis cases in the EU in 2012–2021, with more cases reported during summer months (Figure 3). In 2021, a slight increase in notified human cases was registered compared with 2020. Notwithstanding, the overall trend for salmonellosis in 2017–2021 did not show any statistically significant increase or decrease.

Denmark, Estonia, Finland, Ireland, Romania and Sweden reported a significantly decreasing trend (p < 0.05) in the last 5 years (2017–2021). Only Malta reported a significantly increasing trend (p < 0.05) in 2017–2021.



Source: Austria, Belgium, Cyprus, Czechia, Denmark, Estonia, Germany, Greece, Finland, France, Hungary, Ireland, Italy, Luxembourg, Latvia, Malta, the Netherlands, Poland, Portugal, Romania, Sweden, Slovenia and Slovakia.

**Figure 3:** Trend in reported confirmed human cases of non-typhoidal salmonellosis in the EU by month, 2017–2021

18314732, 2022, 12, Downloaded from https://efs.aonlinelibray.wiley.com/doi/10/2033/jfs/a2022/7666 by Istituto Zoproficitieto's Speriment ad ell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023], See the Terms and Conditions (https://onlinelibray.wiley.com/ems-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons.

<sup>(</sup>a): Y: yes; N: no; A: aggregated data; C: case-based data.

<sup>(</sup>b): Sentinel system; notification rates calculated with an estimated population coverage of 48%.

<sup>(</sup>c): Sentinel system; notification rates calculated with an estimated population coverage of 64%.

<sup>(</sup>d): The notification rate was not calculated since information on estimated coverage was not available.

<sup>(</sup>e): Data not complete in 2020 and 2021: rate not estimated.

<sup>(</sup>f): Cases reported by the United Kingdom for the period 2017–2019 were also considered for this estimation (EU-28). When the United Kingdom data were collected for the period 2017–2019, the United Kingdom was an EU MS, but it became a third country on 1 February 2020.

<sup>(</sup>g): Switzerland provided data directly to EFSA. The human data for Switzerland include data from Liechtenstein for the years 2017–2020.

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (thps://onlinelibrary.wiley.com/terms

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons

The most affected age groups were 0-4 years (27.7%), 5-9 years (13.8%) and over 65 years (16.5%). It is important to underline that for 74.4% of the samples, information about the type of specimen was present; 69.9% of specimens were faeces and the rest consisted of 1.8% other, 1.6% blood, 1.1% urine and 0.03% cerebrospinal fluid and pus.

Sixteen MSs provided information on hospitalisation for 30,951 cases (51.5%) at the EU level. Among these, the proportion of hospitalised cases was 38.1%, which was higher than in 2020. The highest proportions of hospitalised cases were reported, as in previous years, in Cyprus, Greece and Lithuania. Two of these countries also reported the lowest notification rates for salmonellosis, which may suggest that the surveillance systems in these countries primarily capture the most severe cases. Within different classes of specimens, considering cases with information on hospitalisation, higher rates of cases were reported from blood (82.4%), pus (62.5%), urine (39.1%) and faeces (36.5%).

Sixteen EU MSs provided data on the outcome of salmonellosis, accounting for 64.4% of confirmed cases. Among these, 10 reported 71 fatal cases, resulting in an EU case fatality rate of 0.18%.

#### 2.4.3. Salmonella in food

#### Data collected in the context of Regulation (EC) No 2073/2005

#### Food safety criteria

The number of official single samples collected at manufacturing (N = 32,212 samples, notified by 14 MSs) was higher than that at distribution (N = 22,614 samples, notified by 14 MSs), whereas the proportions of positive samples were similar at the two stages (2.81% at manufacturing and 2.53% at distribution) (Table 11).

At manufacturing level, the highest percentages of *Salmonella*-positive samples were reported from 'mechanically separated meat (MSM)' (13.3%), 'meat products made from poultry meat intended to be eaten cooked' (10%), 'minced meat and meat preparations made from poultry meat intended to be eaten cooked' (5.2%) and 'fresh poultry meat' (3.1%). Next, there were 'dried follow-on formulae' (1.1%, one positive out of 90 tested samples) and 'minced meat and meat preparations made from other species than poultry intended to be eaten cooked' (1.1%). For the other food matrices, the percentage of positive samples was lower than 0.30%, or no *Salmonella*-positive samples were reported.

At distribution level, the highest proportions of *Salmonella*-positive samples were reported in 'minced meat and meat preparations made from poultry meat intended to be eaten cooked' (8.7%), 'fresh poultry meat' (6.5%), 'egg products, excluding products where the manufacturing process or the composition of the product will eliminate the *Salmonella* risk' (3.1%) and 'minced meat and meat preparations made from other species than poultry intended to be eaten cooked' (1.3%). For the other food matrices covered by Regulation (EC) No 2073/2005, the percentage of positive samples was consistently lower than 0.70% and for the majority of them, no *Salmonella*-positive samples were reported.

**Table 11:** Proportion (%) of *Salmonella*-positive samples from official sampling as part of the verification of *Salmonella* FSC in accordance with Regulation (EC) No 2073/2005, by stage in the food chain, EU, 2021

		nufacturin cluding pro		Distribution stage (including retail)			
Food matrices	N MSs	N tested samples	N (%) testing positive	N MSs	N tested samples	N (%) testing positive	
Cheeses, butter and cream made from raw milk or milk that has undergone a lower heat treatment than pasteurisation	9	1,718	4 (0.23)	7	2,614	6 (0.23)	
Cooked crustaceans and molluscan shellfish	5	390	0	7	846	6 (0.71)	
Dried follow-on formulae	1	90	1 (1.1)	4	259	0	
Dried infant formulae and dried dietary foods for special medical purposes intended for infants below 6 months of age	_	_	_	5	449	1 (0.22)	



18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wley.com/doi/10.2903j.gfsa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/erms

		nufacturin cluding pro			istribution including r	
Food matrices	N MSs	N tested samples	N (%) testing positive	N MSs	N tested samples	N (%) testing positive
Egg products, excluding products where the manufacturing process or the composition of the product will eliminate the <i>Salmonella</i> risk	5	203	0	4	292	9 (3.1)
Fresh poultry meat	9	16,322	501 (3.1)	10	4,889	319 (6.5)
Gelatine and collagen	2	45	0	7	114	0
Ice cream, excluding products where the manufacturing process or the composition of the product will eliminate the <i>Salmonella</i> risk	5	352	1 (0.28)	7	483	0
Live bivalve molluscs and live echinoderms, tunicates and gastropods	2	535	0	4	371	2 (0.54)
Meat products intended to be eaten raw, excluding products where the manufacturing process or the composition of the product will eliminate the <i>Salmonella</i> risk	9	1,130	3 (0.27)	7	968	4 (0.41)
Meat products made from poultry meat intended to be eaten cooked	2	30	3 (10.0)	1	40	0
Mechanically separated meat (MSM)	7	188	25 (13.3)	2	30	0
Milk powder and whey powder	8	184	0	4	143	0
Minced meat and meat preparations intended to be eaten raw	4	118	0	4	194	0
Minced meat and meat preparations made from other species than poultry intended to be eaten cooked	11	3,879	41 (1.1)	12	5,935	80 (1.3)
Minced meat and meat preparations made from poultry meat intended to be eaten cooked	9	6,236	325 (5.2)	9	1,654	144 (8.7)
Precut fruit and vegetables (ready-to-eat)	5	185	0	9	1,915	2 (0.1)
Ready-to-eat foods containing raw egg, excluding products where the manufacturing process or the composition of the product will eliminate the <i>Salmonella</i> risk	1	285	0	2	50	0
Sprouted seeds (ready-to-eat)	4	67	0	6	383	0
Unpasteurised fruit and vegetable juices (ready- to-eat)	4	255	0	7	985	0
EU Total <sup>(a)</sup>	14	32,212	904 (2.8)	14	22,614	573 (2.5)

MSs: Member States; FSC: food safety criteria.

For a further interactive look at *Salmonella* monitoring results, dashboards have been created (*different filters can be applied to query the data*) (*link*).

# Process hygiene criteria

Considering data on *Salmonella* from carcases of different species, irrespective of the sampler, the prevalence values found for neck skin samples from broilers and turkeys were much higher than those reported for carcase surfaces of ruminants (cattle, sheep and goats) and horses.

# Carcases of pigs

Considering all PHC monitoring data from pig carcases collected at the slaughterhouse after dressing but before chilling sent by a total of 23 MSs, the overall proportion of Salmonella-positive samples based on official controls was 1.7% (N = 24,802) and was significantly higher than that based

<sup>(</sup>a): Data from the United Kingdom (Northern Ireland) are taken into account for 2021. In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/Northern Ireland, the EU requirements on data sampling are also applicable to Northern Ireland.



18314732, 2022, 12, Downloaded from https://dfsa.onlinelibrary.wiley.com/doi/10.2903j.dfsa.2022.7666 by Istituo Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons

on own checks (1.4%, N = 103,270) (Table 12). The same finding was true considering overall data from those MSs (nine) that reported data collected by both the CA (2.2%, N = 15,617) and the FBOp (0.85%, N = 55,699) and specifically for six MSs (Belgium, Estonia, Germany, Ireland, Italy and Spain).

Finland, Sweden and Norway, which are countries with special guarantees in relation to *Salmonella* on pig carcases (in accordance with Regulation (EU) No 853/2004), reported the following monitoring results: one positive out of 2,369 own-check samples taken by the FBOp (0.04%) in Finland, one positive out of 2,833 official samples (0.03%) in Norway and one positive out of 6,426 official samples (0.01%) in Sweden. Moreover, Switzerland reported zero positive out of 1,117 and Montenegro zero positive out of nine own-check samples collected by the FBOp.

**Table 12:** Comparisons of proportions (%) of *Salmonella*-positive single samples from pig carcases after dressing, but before chilling, by sampler and reporting MS, EU, 2021

	Compe	tent autho	ority (CA)	Food bus	siness oper	ator (FBOp)		
Country	N tested samples	N (%) positive samples	CI <sub>95</sub>	N tested samples	N (%) positive samples	CI <sub>95</sub>	p-value <sup>(b)</sup>	Interpretation
Austria	_	-	_	4,056	2 (0.05)	[0.01; 0.18]	-	-
Belgium	955	59 (6.2)	[4.7; 7.9]	4,108	62 (1.5)	[1.2; 1.9]	< 0.001	CA > FBOp
Bulgaria	5,049	0	[0; 0.07] <sup>(a)</sup>	305	0	[0; 1.2] <sup>(a)</sup>	NS	
Croatia	1,630	4 (0.24)	[0.07; 0.63]	_	_	_	_	_
Cyprus	2	0	[-]	_	_	_	-	-
Czechia	4,508	35 (0.78)	[0.54; 1.1]	_	_	_	_	_
Denmark	_	_	_	10,773	80 (0.74)	[0.59; 0.92]	_	-
Estonia	357	6 (1.7)	[0.62; 3.6]	1,581	12 (0.76)	[0.39; 1.3]	0.096	CA > FBOp
France	-	-	_	13,662	628 (4.6)	[4.3; 5.0]	-	_
Germany	200	4 (2.0)	[0.55; 5.0]	24,885	156 (0.63)	[0.53; 0.73]	0.039	CA > FBOp
Hungary	3,045	44 (1.4)	[1.1; 1.9]	_	_	_	_	_
Ireland	391	19 (4.9)	[3.0; 7.5]	2,064	25 (1.2)	[0.78; 1.8]	< 0.001	CA > FBOp
Italy	5,147	174 (3.4)	[2.9; 3.9]	11,494	107 (0.93)	[0.76; 1.1]	< 0.001	CA > FBOp
Latvia	_	_	_	579	3 (0.52)	[0.11; 1.5]	_	_
Luxembourg	_	_	_	345	2 (0.58)	[0.07; 2.1]	-	_
Malta	_	_	_	120	13 (10.8)	[5.9; 17.8]	_	_
Netherlands	_	-	_	5,635	120 (2.1)	[1.8; 2.5]	_	_
Poland	273	0	[0; 1.3] <sup>(a)</sup>	5,576	3 (0.05)	[0.01; 0.16]	NS	
Portugal	-	-	_	8,222	66 (0.8)	[0.62; 1.0]	-	_
Romania	2,540	0	[0; 0.14] <sup>(a)</sup>	2,881	0	[0; 0.13] <sup>(a)</sup>	NS	
Slovakia	_	_	_	3,209	0	[0; 0.12] <sup>(a)</sup>	_	_
Slovenia	_	_	_	970	13 (1.3)	[0.72; 2.3]	_	_
Spain	705	82 (11.6)	[9.4; 14.2]	2,805	109 (3.9)	[3.2; 4.7]	< 0.001	CA > FBOp
EU Total (27 + XI)	24,802	427 (1.7)	[1.6; 1.9]	103,270	1401 (1.4)	[1.3; 1.4]	< 0.001	CA > FBOp
EU Total (27 + XI) providing CA and FBOp data	15,617	344 (2.2)	[2.0; 2.4]	55,699	474 (0.85)	[0.78; 0.93]	< 0.001	CA > FBOp

<sup>-:</sup> Data not reported.

<sup>[-]:</sup> The confidence interval is not provided because of the small sample size.

<sup>(</sup>a): One-sided, 97.5% confidence interval.

<sup>(</sup>b): p-value: NS, not significant.

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico

dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

#### Carcases of broilers

The overall proportion of *Salmonella*-positive PHC neck skin samples collected at the slaughterhouse from broiler carcases after chilling based on official controls was 14.0% (N = 6,544), which was significantly higher than that based on own checks (3.2%, N = 56,532) (Table 13). Similarly, for the five MSs that reported data collected by both samplers, the overall proportion detected in samples collected by the CA (12.8%, N = 3,285) was significantly higher than that reported by the FBOp (4.1%, N = 10,955), and this finding was specifically confirmed for four MSs (Belgium, Italy, Romania and Spain).

Finland, Sweden and Norway, which are countries with special guarantees in relation to *Salmonella* on broiler carcases (in accordance with Regulation (EU) No 853/2004), reported the following monitoring results: Finland (N = 1,271 collected by the FBOp) and Sweden (N = 1,846 collected by the CA) did not report any positive samples, whereas Norway did not report any data for broiler carcases. Moreover, Switzerland reported 13 positive out of 845 (1.5%) and Montenegro two positive out of 50 tested broiler carcase samples collected by the FBOp.

**Table 13:** Comparisons of proportions (%) of *Salmonella*-positive single samples from broiler carcases (neck skin samples) after chilling, by sampler and reporting MS, EU, 2021

•	Comp	etent autho	ority (CA)	Food bu	siness oper	ator (FBOp)		
Country	N tested samples	N (%) positive samples	CI <sub>95</sub>	N tested samples	N (%) positive samples	CI <sub>95</sub>	p-value <sup>(b)</sup>	Interpretation
Austria	_	-	-	1,029	147 (14.3)	[12.2; 16.6]	-	_
Belgium	631	62 (9.8)	[7.6; 12.4]	2,634	128 (4.9)	[4.1; 5.8]	< 0.001	CA > FBOp
Croatia	1,290	270 (20.9)	[18.7; 23.3]	_	_	_	_	_
Cyprus	220	65 (29.5)	[23.6; 36]	-	_	_	_	_
Czechia	1,040	65 (6.2)	[4.9; 7.9]	_	_	_	_	_
Denmark	_	_	_	263	0	[0; 1.4] <sup>(a)</sup>	_	_
Estonia	_	_	_	260	0	[0; 1.4] <sup>(a)</sup>	_	_
France	_	_	_	13,739	319 (2.3)	[2.1; 2.6]	_	_
Germany	_	_	_	19,509	504 (2.6)	[2.4; 2.8]	_	_
Greece	_	_	_	1,991	0	[0; 0.18] <sup>(a)</sup>	_	_
Hungary	709	94 (13.3)	[10.8; 16.0]	-	_	_	_	_
Ireland	279	3 (1.1)	[0.22; 3.1]	1,190	11 (0.92)	[0.46; 1.6]	NS	
Italy	1,125	277 (24.6)	[22.1; 27.2]	4,605	300 (6.5)	[5.8; 7.3]	< 0.001	CA > FBOp
Latvia	_	_	_	599	23 (3.8)	[2.4; 5.7]	_	_
Netherlands	-	_	_	3,536	273 (7.7)	[6.9; 8.7]	_	_
Portugal	_	_	_	2,862	5 (0.17)	[0.06; 0.41]	_	_
Romania	800	59 (7.4)	[5.7; 9.4]	1,891	9 (0.48)	[0.22; 0.90]	< 0.001	CA > FBOp
Slovakia	_	_	_	945	0	[0; 0.39] <sup>(a)</sup>	_	_
Slovenia	_	_	_	844	59 (7.0)	[5.4; 8.9]	_	_
Spain	450	19 (4.2)	[2.6; 6.5]	635	4 (0.63)	[0.17; 1.6]	< 0.001	CA > FBOp
EU Total (27 + XI)	6,544	914 (14.0)	[13.1; 14.8]	56,532	1782 (3.2)	[3.0; 3.3]	< 0.001	CA > FBOp
EU Total (27 + XI) providing CA and FBOp data		420 (12.8)	[11.7; 14.0]	10,955	452 (4.1)	[3.8; 4.5]	< 0.001	CA > FBOp

<sup>-:</sup> Data not reported.

<sup>(</sup>a): One-sided, 97.5% confidence interval.

<sup>(</sup>b): p-value: NS, not significant.

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico

dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

#### Carcases of turkeys

The overall percentage of *Salmonella*-positive PHC neck skin samples collected at the slaughterhouse from turkey carcases after chilling based on official controls was 7.4% (N = 1,321) and was significantly higher than the percentage based on own-check samples collected by the FBOp (3.2%, N = 7,941) (Table 14). The same finding was true considering the overall proportion of positive samples for the four MSs that reported data from both samplers, whereas looking at the data of the single MSs, this finding was confirmed only for Italy.

Finland, Sweden and Norway are countries with special guarantees in relation to Salmonella on turkey carcases (in accordance with Regulation (EU) No 853/2004). Finland (N = 271 collected by the FBOp) and Sweden (N = 124 collected by the CA) did not report any positive samples, whereas Norway did not report any data on turkey carcases. Moreover, Switzerland reported zero positive out of 135 tested turkey samples collected by the FBOp.

**Table 14:** Comparisons of proportions (%) of *Salmonella*-positive single samples from turkey carcases (neck skin samples) after chilling, by sampler and reporting MS, EU, 2021

	carcases (neck skin samples) after chilling, by sampler and reporting MS, EU, 2021										
	Compet	tent auth	ority (CA)	Food busi	ness operat	or (FBOp)					
Country	N tested samples	N (%) positive samples	CI <sub>95</sub>	N tested samples	N (%) positive samples	CI <sub>95</sub>	p-value <sup>(b)</sup>	Interpretation			
Austria	_	_	-	135	5 (3.7)	[1.2; 8.4]	-	-			
Belgium	49	0	[0; 7.3] <sup>(a)</sup>	135	0	[0; 2.7] <sup>(a)</sup>	NS				
Czechia	300	3 (1.0)	[0.21; 2.9]	_	_	_	_	_			
France	_	_	-	2,948	35 (1.2)	[0.83; 1.6]	_	_			
Germany	_	_	_	1,611	102 (6.3)	[5.2; 7.6]	_	_			
Greece	_	_	-	116	0	[0; 3.1] <sup>(a)</sup>	-	_			
Hungary	651	32 (4.9)	[3.4; 6.9]	_	_	_	_	_			
Ireland	47	0	[0; 7.5] <sup>(a)</sup>	538	0	[0; 0.68] <sup>(a)</sup>	NS				
Italy	144	29 (20.1)	[13.9; 27.6]	1,079	91 (8.4)	[6.8; 10.3]	< 0.001	CA > FBOp			
Latvia	_	_	_	5	0	[-]	_	_			
Poland	_	_	-	371	7 (1.9)	[0.76; 3.8]	_	_			
Portugal	_	_	-	752	6 (0.80)	[0.29; 1.7]	_	_			
Romania	30	0	[0; 11.6] <sup>(a)</sup>	25	0	[0; 13.7] <sup>(a)</sup>	NS				
Slovakia	_	_	-	60	0	[0; 6.0] <sup>(a)</sup>	_	_			
Slovenia	_	_	-	166	5 (3.0)	[0.98; 6.9]	_	_			
Spain	100	34 (34.0)	[24.8; 44.2]	_	_	_	_	_			
EU Total (27 + XI)	1,321	98 (7.4)	[6.1; 9.0]	7,941	251 (3.2)	[2.8; 3.6]	< 0.001	CA > FBOp			
EU Total (27 + XI) providing CA and FBOp data	270	29 (10.7)	[7.3; 15.1]	1,777	91 (5.1)	[4.1; 6.2]	< 0.001	CA > FBOp			

<sup>-:</sup> Data not reported.

#### Carcases of cattle

The overall percentage of *Salmonella*-positive PHC samples from bovine carcases collected at the slaughterhouse after dressing but before chilling based on official controls was 0.81% (N = 13,017) and was significantly higher than that based on own checks conducted by the FBOp (0.24%, N = 69,412) (Table 15). The same finding was true considering the overall proportion of positive samples for the five MSs that reported data from both samplers (1.1%, N = 5,784 collected by the CA and 0.42%, N = 15,828 collected by the FBOp) and specifically for Italy and Spain. Only Belgium

<sup>(</sup>a): One-sided, 97.5% confidence interval.

<sup>(</sup>b): p-value: NS, not significant.



Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

reported a higher percentage of *Salmonella*-positive samples based on own checks (0.34%, N = 2,667) than for samples based on official controls (0%, N = 1,164).

Finland, Sweden and Norway are countries with special guarantees in relation to *Salmonella* on bovine carcases (in accordance with Regulation (EU) No 853/2004). Finland (N = 2,177 collected by the FBOp) and Norway (N = 3,172 collected by the CA) did not report any positive samples, whereas Sweden reported two positive out of 3,831 tested samples collected by the CA (0.05%). Moreover, Switzerland reported two positive samples out of 1,017 (0.2%), Iceland zero positive out of 300 and Montenegro zero positive out of 66 tested samples collected by the FBOp.

**Table 15:** Comparisons of proportions (%) of *Salmonella*-positive single samples from bovine carcases after dressing but before chilling, by sampler and reporting MS, EU, 2021

	Compet	ent autho	ority (CA)	Food bus	siness opera	ator (FBOp)		
Country	N tested samples	N (%) positive samples	CI <sub>95</sub>	N tested samples	N (%) positive samples	CI <sub>95</sub>	p-value <sup>(b)</sup>	Interpretation
Austria	-	-	-	2,607	0	[0; 0.14] <sup>(a)</sup>	-	_
Belgium	1,164	0	[0; 0.32] <sup>(a)</sup>	2,667	9 (0.34)	[0.15; 0.64]	0.038	CA < FBOp
Bulgaria	44	0	[0; 8.0] <sup>(a)</sup>	_	-	-	-	_
Croatia	2,819	32 (1.1)	[0.78; 1.6]	_	_	_	_	_
Czechia	3,951	9 (0.23)	[0.1; 0.43]	_	_	_	-	_
Denmark	_	_	_	4,971	3 (0.06)	[0.01; 0.18]	_	_
Estonia	222	3 (1.4)	[0.28; 3.9]	_	_	_	-	_
France	_	_	_	18,088	38 (0.21)	[0.15; 0.29]	_	_
Germany	_	_	_	11,283	7 (0.06)	[0.03; 0.13]	_	_
Greece	_	_	_	167	0	[0; 2.2] <sup>(a)</sup>	_	_
Hungary	197	1 (0.51)	[0.01; 2.8]	_	_	_	-	_
Ireland	_	_	_	6,203	1 (0.02)	[0; 0.09]	_	_
Italy	2,409	38 (1.6)	[1.1; 2.2]	10,009	58 (0.58)	[0.44; 0.75]	< 0.001	CA > FBOp
Latvia	_	_	_	539	0	[0; 0.68] <sup>(a)</sup>	_	_
Luxembourg	_	_	_	255	0	[0; 1.4] <sup>(a)</sup>	_	_
Malta	_	_	_	120	26 (21.7)	[14.7; 30.1]	_	_
Netherlands	_	_	_	2,572	9 (0.35)	[0.16; 0.66]	_	_
Poland	105	0	[0; 3.5] <sup>(a)</sup>	108	0	[0; 3.4] <sup>(a)</sup>	NS	
Portugal	_	_	_	3,120	14 (0.45)	[0.25; 0.75]	_	_
Romania	1,471	0	[0; 0.25] <sup>(a)</sup>	2,194	0	[0; 0.17] <sup>(a)</sup>	NS	
Slovakia	_	_	_	2,107	0	[0; 0.17] <sup>(a)</sup>	_	_
Slovenia	_	_	_	1,552	0	[0; 0.24] <sup>(a)</sup>	_	_
Spain	635	23 (3.6)	[2.3; 5.4]	850	0	[0; 0.43] <sup>(a)</sup>	< 0.001	CA > FBOp
EU Total (27 + XI)	13,017	106 (0.81)	[0.67; 0.98]	69,412	165 (0.24)	[0.2; 0.28]	< 0.001	CA > FBOp
EU Total (27 + XI) providing CA and FBOp data	5,784	61 (1.1)	[0.81; 1.4]	15,828	67 (0.42)	[0.33; 0.54]	< 0.001	CA > FBOp

<sup>-:</sup> Data not reported.

### Carcases of sheep

The overall percentage of Salmonella-positive PHC samples from sheep carcases collected at the slaughterhouse after dressing but before chilling based on official controls was 1.2% (N = 3,693) and was significantly higher than that based on own checks (0.49%, N = 16,683) (Table 16). The same finding was true considering overall data sent by the four MSs providing both CA (0.81%, N = 1,238)

<sup>(</sup>a): One-sided, 97.5% confidence interval.

<sup>(</sup>b): p-value: NS, not significant.



18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (thps://onlinelibrary.wiley.com/terms

-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons

and FBOp (0.11%, N = 4,673) data and specifically for Belgium and Italy. Moreover, Switzerland reported zero positive out of 367, Iceland zero positive out of 132 and Montenegro zero positive out of 12 tested samples collected by the FBOp. There are no special guarantees in relation to *Salmonella* in sheep carcases.

**Table 16:** Comparisons of proportions (%) of *Salmonella*-positive single samples from sheep carcases after dressing but before chilling, by sampler and reporting MS, EU, 2021

,	Compete	ent autho	rity (CA)	Food bu	siness ope	erator (FBOp)		
Country	N tested samples	N (%) positive samples	CI <sub>95</sub>	N tested samples	N (%) positive samples	CI <sub>95</sub>	p-value <sup>(b)</sup>	Interpretation
Austria	_	_	_	229	0	[0; 1.6] <sup>(a)</sup>	_	_
Belgium	338	4 (1.2)	[0.32; 3.0]	641	1 (0.16)	[0; 0.87]	0.051	CA > FBOp
Bulgaria	120	0	[0; 3.0] <sup>(a)</sup>	_	_	_	_	_
Croatia	2,034	33 (1.6)	[1.1; 2.3]	_	_	_	_	_
Cyprus	10	0	[0; 30.9] <sup>(a)</sup>	_	_	_	_	_
Czechia	291	0	[0; 1.3] <sup>(a)</sup>	_	_	_	_	_
Finland	_	_	_	57	0	[0; 6.3] <sup>(a)</sup>	_	_
France	_	_	_	6,265	51 (0.81)	[0.61; 1.1]	_	_
Germany	_	_	_	907	3 (0.33)	[0.07; 0.96]	-	_
Ireland	_	_	_	1,402	2 (0.14)	[0.02; 0.51]	_	_
Italy	346	6 (1.7)	[0.64; 3.7]	3,135	4 (0.13)	[0.04; 0.33]	< 0.001	CA > FBOp
Latvia	_	_	_	143	0	[0; 2.5] <sup>(a)</sup>	_	_
Luxembourg	_	_	_	78	0	[0; 4.6] <sup>(a)</sup>	-	_
Malta	_	_	_	96	4 (4.2)	[1.1; 10.3]	_	_
Netherlands	_	_	_	288	0	[0; 1.3] <sup>(a)</sup>	-	_
Portugal	_	_	_	2,101	17 (0.81)	[0.47; 1.3]	_	_
Romania	464	0	[0; 0.79] <sup>(a)</sup>	652	0	[0; 0.56] <sup>(a)</sup>	NS	
Slovakia	_	_	_	352	0	[0; 1.0] <sup>(a)</sup>	_	_
Slovenia	_	_	_	92	0	[0; 3.9] <sup>(a)</sup>	-	_
Spain	90	0	[0; 4.0] <sup>(a)</sup>	245	0	[0; 1.5] <sup>(a)</sup>	NS	
EU Total (27 + XI)	3,693	43 (1.2)	[0.84; 1.6]	16,683	82 (0.49)	[0.39; 0.61]	< 0.001	CA > FBOp
EU Total (27 + XI) providing CA and FBOp data	1,238	10 (0.81)	[0.39; 1.5]	4,673	5 (0.11)	[0.04; 0.25]	< 0.001	CA > FBOp

<sup>-:</sup> Data not reported.

#### Carcases of goats

The overall percentage of *Salmonella*-positive PHC samples from goat carcases collected at the slaughterhouse after dressing but before chilling based on own checks was 2.1% (N = 1,211) and was significantly higher than that based on official controls (0.51%, N = 585). This finding was markedly influenced by the results of two MSs (France and Portugal) that reported the large majority of the positive samples from goat carcases notified by the FBOp (Table 17). There are no special guarantees in relation to *Salmonella* in goat carcases.

<sup>(</sup>a): One-sided, 97.5% confidence interval.

<sup>(</sup>b): p-value: NS, not significant.



**Table 17:** Comparisons of proportions (%) of *Salmonella*-positive single samples from goat carcases after dressing but before chilling, by sampler and reporting MS, EU, 2021

	Compe	tent auth	ority (CA)	Food bus	siness oper	ator (FBOp)		
Country	N tested	N (%) positive samples	CI <sub>95</sub>	N tested samples	N (%) positive	CI <sub>95</sub>	p-value <sup>(b)</sup>	Interpretation
Austria	_	_	-	13	0	[0; 24.7] <sup>(a)</sup>	-	_
Belgium	113	2 (1.8)	[0.22; 6.2]	55	0	[0; 6.5] <sup>(a)</sup>	NS	
Bulgaria	5	0	[-]	-	-	_	_	_
Croatia	314	1 (0.32)	[0.01; 1.8]	_	_	_	_	_
Czechia	70	0	[0; 5.1] <sup>(a)</sup>	-	_	_	_	_
France	_	_	_	171	15 (8.8)	[5.0; 14.1]	_	_
Germany	_	_	_	36	0	[0; 9.7] <sup>(a)</sup>	_	_
Greece	_	_	_	203	0	[0; 1.8] <sup>(a)</sup>	_	_
Italy	53	0	[0; 6.7] <sup>(a)</sup>	114	1 (0.88)	[0.02; 4.8]	NS	
Malta	_	_	_	24	1 (4.2)	[0.10; 21.1]	_	_
Netherlands	_	_	_	42	0	[0; 8.4] <sup>(a)</sup>	_	_
Portugal	_	_	_	531	9 (1.7)	[0.78; 3.2]	_	_
Slovakia	_	_	-	7	0	[-]	_	_
Slovenia	_	_	_	15	0	[0; 21.8] <sup>(a)</sup>	_	_
Spain	30	0	[0; 11.6] <sup>(a)</sup>	_	_	_	_	_
EU Total (27 + XI)	585	3 (0.51)	[0.11; 1.5]	1,211	26 (2.1)	[1.4; 3.1]	0.005	CA < FBOp
EU Total (27 + XI) providing CA and FBOp data	166	2 (1.2)	[0.15; 4.3]	169	1 (0.59)	[0.01; 3.3]	NS	

<sup>-:</sup> Data not reported.

# Carcases of horses

The overall percentage of *Salmonella*-positive PHC samples from horse carcases collected at the slaughterhouse after dressing but before chilling based on official controls was 0.19% (N = 517) and was not significantly higher than that based on FBOp own checks (0.15%, N = 1,360) (Table 18). There are no special guarantees in relation to *Salmonella* in horse carcases.

**Table 18:** Comparisons of proportions (%) of *Salmonella*-positive single samples from horse carcases after dressing but before chilling, by sampler and reporting MS, EU, 2021

Country	Compet	Competent authority (CA)			Food business operator (FBOp)			
	N tested samples	DOCITIVA.	CI <sub>95</sub>	N tested samples		CI <sub>95</sub>	p-value <sup>(b)</sup>	Interpretation
Austria	_	_	_	6	0	[-]	_	_
Belgium	69	0	[0; 5.2] <sup>(a)</sup>	114	0	[0; 3.2] <sup>(a)</sup>	NS	
Croatia	49	0	[0; 7.3] <sup>(a)</sup>	_	_	_	_	_
Czechia	7	0	[-]	_	_	_	_	_
Finland	_	_	_	1	0	-	_	_
France	_	_	_	70	0	[0; 5.1] <sup>(a)</sup>	_	_
Germany	_	_	_	21	0	[0; 16.1] <sup>(a)</sup>	_	_
Ireland	_	_	_	62	0	[0; 5.8] <sup>(a)</sup>	_	_

18314732, 2022, 12, Downloaded from https://efs.aonlinelibrary.wiley.com/doi/10.2903j.d/sa.2022.72666 by Istituto Cooperinents of the applicable Cereive Common.

<sup>(</sup>a): One-sided, 97.5% confidence interval.

<sup>(</sup>b): p-value: NS, not significant.

onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico

dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

	Compet	Competent authority (CA)			business (FBOp	operator )		
Country	N tested samples	N (%) positive samples	CI <sub>95</sub>	N tested samples		CI <sub>95</sub>	p-value <sup>(b)</sup>	Interpretation
Italy	177	1 (0.56)	[0.01; 3.1]	820	1 (0.12)	[0; 0.68]	NS	
Latvia	_	_	_	3	0	[-]	_	_
Poland	5	0	[-]	_	_	_	-	_
Portugal	_	_	_	52	1 (1.9)	[0.05; 10.3]	_	_
Romania	145	0	[0; 2.5] <sup>(a)</sup>	186	0	[0; 2.0] <sup>(a)</sup>	NS	
Slovenia	_	_	_	25	0	[0; 13.7] <sup>(a)</sup>	_	_
Spain	65	0	[0; 5.5] <sup>(a)</sup>	_	_	_	_	_
EU Total (27 + XI)	517	1 (0.19)	[0; 1.1]	1,360	2 (0.15)	[0.02; 0.53]	NS	
EU Total (27 + XI) providing CA and FBOp data	391	1 (0.26)	[0.01; 1.4]	1,120	1 (0.09)	[0; 0.5]	NS	

<sup>-:</sup> Data not reported.

For a further interactive look at *Salmonella* monitoring results, dashboards have been created (*different filters can be applied to query the data*) (link).

#### Occurrence in food

Monitoring data reported for food samples, which do not fit with the criteria described in the previous paragraphs, were described by merging investigations from all sampling stages (primary production, manufacturing, distribution, other and unspecified), all samplers except 'HACCP and own checks' and 'private sampling' and all sampling units (single, batch and slaughter animal batch). Only samples collected through 'objective sampling' were considered in this context.

#### RTE food and non-RTE food

In 2021, 73,238 RTE and 466,290 non-RTE food sampling units were reported from 23 and 26 MSs with very low (0.23%) and low (2.1%) proportions of positive samples, respectively (Table 19).

Within the category of RTE food, all matrices with *Salmonella* isolates had very low proportions of positive samples and the highest percentages were from 'meat and meat products from pigs' (0.82%), 'spices and herbs' (0.72%), 'fish and fishery products' (0.39%) and 'other meat and meat products' (0.32%).

Within the category of non-RTE food, the highest percentages of positive samples were reported for 'meat and meat products from broilers' (4.4%), 'meat and meat products from turkeys' (3.6%), 'other meat and meat products' (2.2%), 'meat and meat products from pigs' (1.5%), 'mixed meat and meat products' (1.1%) and 'other processed food products and prepared dishes' (1.0%). Some *Salmonella* isolates were also reported for 'eggs and egg products' (0.82%) and 'fish and fishery products' (0.65%).

Comparing the results for the year 2021 and the 4-year period of 2017–2020, the overall percentage of *Salmonella*-positive RTE food samples remained rather stable. The greatest percentage reductions in *Salmonella*-positive samples were found for 'meat and meat products from broilers', 'meat and meat products from turkeys', 'infant formulae and follow-on formulae–RTE', 'other meat and meat products' and 'salads'. In contrast, a small increase was reported for 'meat and meat products from pigs' and for 'fish and fishery products'.

Similarly, also for non-RTE food, the overall percentage of *Salmonella*-positive samples reported in 2021 was very close to the average reported over the period 2017–2020. The greatest reductions in proportions of *Salmonella*-positive samples were found for 'meat and meat products from broilers', 'meat and meat products from turkeys', 'fruits, vegetables and juices' and 'cereals and dried seeds'.

<sup>(</sup>a): One-sided, 97.5% confidence interval.

<sup>(</sup>b): p-value: NS, not significant.



Conversely, an increase was reported in the last year for 'other processed food products and prepared dishes', 'other meat and meat products' and 'eggs and egg products'.

#### Fresh meat

For fresh meat, in 2021, 2.1% of sampling units (N = 387,152) were positive for *Salmonella*. Within this category, the highest percentages of positive samples were reported for 'fresh meat from broilers' (4.4%) and for 'fresh meat from turkeys' (3.5%), with a general reduction vs. the proportions of positive samples recorded in the previous years.

**Table 19:** Occurrence of *Salmonella* in the main food categories, EU, 2021 and 2017–2020

		2021 <sup>(a)</sup>			2017–2020	(b)
Food	N reporting MSs	N sampled units	N positive (%)	N reporting MSs	N sampled units	N positive (%)
RTE food		,				
All	23	73,238	169 (0.23)	25	279,268	746 (0.27)
Meat and meat products	21	22,589	99 (0.44)	21	66,284	320 (0.48)
Meat and meat products from broilers	12	1,706	0	16	5,128	25 (0.49)
Meat and meat products from turkeys	9	213	0	14	1,593	7 (0.44)
Meat and meat products from pigs	18	8,014	66 (0.82)	19	27,874	110 (0.39)
Meat and meat products from bovine animals	16	1,219	2 (0.16)	19	4,243	14 (0.33)
Mixed	10	2,956	4 (0.14)	17	5,289	18 (0.34)
Other meat and meat products	13	8,481	27 (0.32)	17	22,157	146 (0.66)
Milk and milk products	21	21,583	22 (0.1)	24	93,802	111 (0.12)
Milk	12	634	1 (0.16)	15	2,828	1 (0.04)
Raw milk <sup>(c)</sup>	5	211	0	7	1,812	0
Cheese	21	15,422	19 (0.12)	23	48,587	74 (0.15)
Dairy products excluding cheeses (butter, cream, ice cream, whey, yoghurt and fermented dairy products)	20	5,527	2 (0.04)	21	42,387	36 (0.08)
Fruits and vegetables and juices	18	6,261	3 (0.05)	20	20,591	22 (0.11)
Fish and fishery products	22	2,809	11 (0.39)	22	12,909	10 (0.08)
Spices and herbs	16	1,529	11 (0.72)	19	5,636	41 (0.73)
Bakery products	17	2,940	0 (0)	18	16,110	34 (0.21)
Salads	9	2,194	1 (0.05)	15	12,866	47 (0.37)
Other processed food products and prepared dishes	17	8,891	16 (0.18)	16	30,362	122 (0.4)
Eggs and egg products	3	46	0	8	244	0
Sprouts (sprouted seeds)	9	512	0	11	1,234	2 (0.16)
Cereals and nuts	9	859	1 (0.12)	15	3,008	3 (0.1)
Confections	4	73	0	7	3,740	3 (0.08)
Infant formulae and follow-on formulae_RTE	12	1,127	1 (0.09)	16	4,933	22 (0.45)
Foodstuffs intended for special nutritional uses	10	588	1 (0.17)	14	1,905	1 (0.05)

1831/472, 2022, 12, Downloaded from https://efs.aonininlabrary.wiley.com/oid/10.2003j.efs.a.2022/2666by Isitiuz Oxported by the applicable Ceawive Common. 1812/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/erem-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Ceawive Common.



18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [1.8/12/2023]. See the Terms

		2021 <sup>(a)</sup>		2017–2020 <sup>(b)</sup>			
Food	N reporting MSs	N sampled units	N positive (%)	N reporting MSs	N sampled units	N positive (%)	
Non-RTE food							
All	26	466,290	9,764 (2.1)	28	1,034,606	22,086 (2.1)	
Meat and meat products	26	443,056	9,593 (2.2)	28	945,084	21,747 (2.3)	
Meat and meat products from broilers	25	109,342	4,763 (4.4)	26	148,380	9,521 (6.4)	
Meat and meat products from turkeys	22	13,049	472 (3.6)	25	21,513	1,097 (5.1)	
Meat and meat products from pigs	26	136,975	2,050 (1.5)	28	390,797	6,789 (1.7)	
Meat and meat products from bovine animals	25	86,507	237 (0.27)	27	135,711	532 (0.39)	
Mixed	14	6,102	66 (1.1)	19	25,656	218 (0.85)	
Other meat and meat products	25	91,081	2,005 (2.2)	25	223,027	3,590 (1.6)	
Milk and milk products	11	996	0 (0)	13	4,902	1 (0.02)	
Fruits, vegetables and juices	13	1,795	1 (0.06)	21	9,078	54 (0.59)	
Fish and fishery products	18	7,093	46 (0.65)	21	27,074	147 (0.54)	
Eggs and egg products	18	6,501	53 (0.82)	20	27,281	92 (0.34)	
Sprouts (sprouted seeds)	5	85	0	8	1,337	5 (0.37)	
Infant formulae	4	175	0	4	244	0	
Foodstuffs intended for special nutritional uses	5	317	0	7	789	2 (0.25)	
Cereals, dried seeds	9	409	1 (0.24)	18	2,700	21 (0.78)	
Other processed food products and prepared dishes	10	3,779	38 (1)	18	10,240	12 (0.12)	
Fresh meat <sup>(d)</sup>							
All	26	387,152	8,219 (2.1)	26	122,337	18,812 (2.4)	
Fresh meat from broilers	25	99,370	4,382 (4.4)	25	17,843	8,586 (7.0)	
Fresh meat from turkeys	22	12,600	441 (3.5)	28	348,156	989 (5.5)	
Fresh meat from pigs	26	124,467	1,902 (1.5)	27	129,721	6,205 (1.8)	
Fresh meat from bovine animals	25	83,703	214 (0.26)	24	154,371	481 (0.37)	
Other fresh meat	25	67,012	1,280 (1.9)	26	122,337	2,551 (1.7)	

MSs: Member States; RTE: ready-to-eat.

For a further interactive look at *Salmonella* monitoring results, dashboards have been created (different filters can be applied to query the data) (link).

#### 2.4.4. Salmonella in animals

For a further interactive look at *Salmonella* monitoring results, dashboards have been created (different filters can be applied to query the data) (link).

<sup>(</sup>a): Data from the United Kingdom (Northern Ireland) are taken into account for 2021. In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/Northern Ireland, the EU requirements on data sampling are also applicable to Northern Ireland.

<sup>(</sup>b): Data from the United Kingdom are taken into account for 2017–2019, because the United Kingdom was an EU MS, but it became a third country on 1 February 2020.

<sup>(</sup>c): Raw RTE milk sampling units are a subset of RTE milk.

<sup>(</sup>d): Fresh meat sampling units are a subset of the two main categories above.




18314732, 2022, 12, Downlo

com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilattico

dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

# Poultry monitoring data according to the national control programmes for *Salmonella Achievement of Salmonella reduction targets*



Percentage (%) of positive flocks

Vertical bars indicate the target to be reached, which was set at 1% for all poultry populations with the exception of laying hens, for which it was 2%.

**Figure 4:** Prevalence of poultry flocks (breeding flocks of *Gallus gallus*, laying hens, broilers, breeding turkeys and fattening turkeys) positive for target *Salmonella* serovars, EU MSs and non-MS countries, 2021

# Breeding flocks of Gallus gallus

In total, 25 MSs, the United Kingdom (Northern Ireland) and three non-MSs reported *Salmonella* NCP data for breeding flocks of *Gallus gallus*. Luxembourg and Malta do not have such flocks. In the EU in



18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico

Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

2021, considering merged data from the CA and FBOp, *Salmonella* was found in 348 (2.5%) of the 13,983 flocks tested, compared with 2.0% and 2.3% for 2020 and 2019, respectively. The prevalence of flocks that were positive for any of the five target serovars was 0.58% in 2021, while it was 0.52% in 2020 and 0.62% in 2019. In 2020, Poland did not report any data for *Gallus gallus* breeders. All reporting countries, except Austria, Belgium, Greece, Ireland and Poland, met the flock prevalence target of 1% maximum (Figure 4). The most frequently reported target serovar was S. Enteritidis (EU flock prevalence of 0.39%, 55 positive flocks), with 44 flocks (80.0%) reported by Poland. The total number of S. Enteritidis-positive breeding flocks (55) increased compared to 2020 (29 positive flocks) and 2019 (53 positive flocks). S. Typhimurium (including the monophasic variant) was the second most commonly reported target serovar (with 15 positive flocks). The third most commonly reported serovar was S. Infantis (with six positive flocks). With regard to the other target serovars, three flocks tested positive for S. Virchow (0.02%) and two flocks tested positive for S. Hadar (0.01%) (Table 20).

**Table 20:** Salmonella in breeding flocks of Gallus gallus during the production period (all types of breeding flocks, flock-based data) in countries running control programmes in accordance with Regulation (EC) No 2160/2003, 2021

		N (%)	N (%)		N (%) positive	e sample:	s for	
Country	N tested flocks	positive for Salmonella spp.	positive for target serovars	S. Enteritidis	<i>S</i> . Typhimurium	<i>S</i> . Hadar	S. Infantis	S. Virchow
Austria	184	6 (3.3)	2 (1.1)	1 (0.54)	1 (0.54)	0	0	0
Belgium	556	12 (2.2)	6 (1.1)	1 (0.18)	4 (0.72)	0	1 (0.18)	0
Bulgaria	608	3 (0.49)	1 (0.16)	0	0	0	1 (0.16)	0
Croatia	95	0	0	_	_	_	_	_
Cyprus	34	0	0	_	_	_	-	_
Czechia	641	0	0	_	_	_	_	_
Denmark	308	1 (0.32)	1 (0.32)	0	1 (0.32)	0	0	0
Estonia	7	0	0	_	_	_	_	_
Finland	145	0	0	_	_	_	-	_
France	1,555	22 (1.4)	6 (0.39)	2 (0.13)	3 (0.19)	0	1 (0.06)	0
Germany	882	42 (4.8)	1 (0.11)	1 (0.11)	0	0	0	0
Greece	224	30 (13.4)	4 (1.8)	1 (0.45)	0	0	0	3 (1.3)
Hungary	554	4 (0.72)	4 (0.72)	1 (0.18)	1 (0.18)	1 (0.18)	1 (0.18)	0
Ireland	175	3 (1.7)	2 (1.1)	2 (1.1)	0	0	0	0
Italy	1,176	60 (5.1)	1 (0.09)	0	1 (0.09)	0	0	0
Latvia	38	0	0	_	_	_	_	_
Lithuania	47	0	0	_	_	_	_	_
Netherlands	1,540	12 (0.78)	6 (0.39)	2 (0.13)	3 (0.20)	0	1 (0.06)	0
Poland	1,991	53 (2.7)	45 (2.3)	44 (2.2)	0	0	1 (0.05)	0
Portugal	476	3 (0.63)	0	_	_	_	_	_
Romania	457	7 (1.5)	0	_	_	_	-	_
Slovakia	115	0	0	_	_	_	_	_
Slovenia	131	0	0	_	_	-	-	_
Spain	1,620	90 (5.6)	2 (0.12)	0	1 (0.06)	1 (0.06)	0	0
Sweden	156	0	0	_	_	_	-	_
United Kingdom (Northern Ireland)	268	0	0	_	-	-	-	-
EU Total (27 + XI)	13,983	348 (2.5)	81 (0.58)	55 (0.39)	15 (0.11)	2 (0.01)	6 (0.04)	3 (0.02)
Iceland	43	0	0	_	_	_	_	_
Norway	230	0	0	_	-	_	_	_
Switzerland	122	0	0	_	_	_	_	_

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilattico

Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

-and--conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

#### Flocks of laying hens

All MSs, the United Kingdom (Northern Ireland) and three non-MSs reported *Salmonella* NCP data for laying hen flocks. Considering merged data from the CA and FBOp, *Salmonella* was found in 1,323 flocks (3.3%), compared with 1,389 (4.0%) in 2020. The EU prevalence of laying hen flocks that were positive for either of the two target serovars was 1.3%, which was stable compared with 2020, when 1.3% of tested flocks were positive for target serovars.

Seven MSs (Belgium, Cyprus, Czechia, France, Malta, Poland and Spain) did not meet the reduction target of 2% (Figure 4). The most frequently reported target serovar was *S*. Enteritidis (EU flock prevalence of 1.0%), with 81.3% of 407 *S*. Enteritidis-positive flocks reported by six MSs. France alone accounted for 28.0% (114 positive flocks) of the *S*. Enteritidis notified; this situation was similar to that of the previous year (29.2%). For *S*. Typhimurium (including the monophasic variant), 126 positive flocks were reported (EU flock prevalence of 0.32%) and the majority (36.5%; 46 positive flocks) were reported by France (Table 21).

**Table 21:** Salmonella in laying hen flocks of Gallus gallus during the production period (flock-based data) in countries running control programmes in accordance with Regulation (EC) No 2160/2003, 2021

	N tested	N (%) positive	N (%) positive	N (%) positiv	e samples for
Country	flocks	for <i>Salmonella</i> spp.	for target serovars	S. Enteritidis	S. Typhimurium
Austria	3,318	25 (0.75)	10 (0.3)	10 (0.3)	0
Belgium	642	35 (5.5)	14 (2.2)	11 (1.7)	3 (0.47)
Bulgaria	1,194	6 (0.50)	2 (0.17)	2 (0.17)	0
Croatia	364	18 (4.9)	4 (1.1)	2 (0.55)	2 (0.55)
Cyprus	126	18 (14.3)	10 (7.9)	7 (5.6)	3 (2.4)
Czechia	491	22 (4.5)	19 (3.9)	13 (2.6)	6 (1.2)
Denmark	429	4 (0.93)	2 (0.47)	2 (0.47)	0
Estonia	35	1 (2.9)	1 (2.9)	1 (2.9)	0
Finland	651	4 (0.61)	4 (0.61)	2 (0.31)	2 (0.31)
France	6,996	300 (4.3)	160 (2.3)	114 (1.6)	46 (0.66)
Germany	7,070	73 (1.0)	48 (0.68)	37 (0.52)	11 (0.16)
Greece	728	45 (6.2)	10 (1.4)	6 (0.82)	4 (0.55)
Hungary	958	11 (1.1)	11 (1.1)	7 (0.73)	4 (0.42)
Ireland	383	1 (0.26)	1 (0.26)	0	1 (0.26)
Italy	4,300	243 (5.7)	35 (0.81)	26 (0.6)	9 (0.21)
Latvia	71	0	0	_	_
Lithuania	47	3 (6.4)	0	_	_
Luxembourg	46	0	0	_	_
Malta	109	25 (22.9)	3 (2.8)	2 (1.8)	1 (0.92)
Netherlands	2,744	51 (1.9)	33 (1.2)	32 (1.2)	1 (0.04)
Poland	2,242	89 (4.0)	72 (3.2)	69 (3.1)	3 (0.13)
Portugal	375	18 (4.8)	4 (1.1)	4 (1.1)	0
Romania	873	49 (5.6)	6 (0.69)	6 (0.69)	0
Slovakia	288	0	0	_	_
Slovenia	263	3 (1.1)	0	_	-
Spain	3,201	268 (8.4)	80 (2.5)	53 (1.7)	27 (0.84)
Sweden	891	3 (0.34)	3 (0.34)	1 (0.11)	2 (0.22)
United Kingdom (Northern Ireland)	711	8 (1.1)	1 (0.14)	0	1 (0.14)
EU Total (27 + XI)	39,546	1,323 (3.3)	533 (1.3)	407 (1.0)	126 (0.32)
Iceland	55	0	0	-	_
Norway	910	1 (0.11)	0	_	_
Switzerland	677	4 (0.59)	4 (0.59)	4 (0.59)	0

Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

#### Broiler flocks

All MSs, the United Kingdom (Northern Ireland) and three non-MSs reported *Salmonella* NCP data for broiler flocks. Considering merged data from the CA and FBOp, *Salmonella* was found in 3.8% (N = 12,040) of the tested flocks, compared with 3.9% in 2020 and 3.6% in 2019. The EU prevalence of broiler flocks positive for either of the two target *Salmonella* serovars was 0.28% (corresponding to 901 flocks), which was similar to the prevalence in previous years (0.25% in 2020 and 0.20% in 2019). Three MSs (Czechia, Luxembourg and Malta) did not meet the target of 1% or less of broiler flocks positive for *S*. Enteritidis and/or *S*. Typhimurium, like in the previous year, when the same three countries did not meet the target. As already reported in 2019 and 2020, the EU prevalence was very similar for the two target serovars: in 2021, *S*. Enteritidis accounted for 51.4% of flocks positive for target serovars, whereas S. Typhimurium accounted for 48.6%. France and Poland accounted for 65.0% of all the EU flocks positive for *S*. Enteritidis, and France alone accounted for 64.6% of all the EU broiler flocks positive for *S*. Typhimurium (Table 22).

**Table 22:** Salmonella in broiler flocks of Gallus gallus before slaughter (flock-based data) in countries running control programmes in accordance with Regulation (EC) No 2160/2003, 2021

_	N tested		N (%) positive	N (%) positiv	e samples for
Country	flocks	for <i>Salmonella</i> spp.	for target serovars	S. Enteritidis	S. Typhimurium
Austria	6,153	167 (2.7)	9 (0.15)	3 (0.05)	6 (0.10)
Belgium	11,194	413 (3.7)	30 (0.27)	9 (0.08)	21 (0.19)
Bulgaria	8,482	2 (0.02)	0	_	-
Croatia	3,120	134 (4.3)	7 (0.22)	3 (0.10)	4 (0.13)
Cyprus	1,033	5 (0.48)	0	0	0
Czechia	4,945	97 (2.0)	70 (1.4)	65 (1.3)	5 (0.10)
Denmark	3,758	6 (0.16)	4 (0.11)	0	4 (0.11)
Estonia	669	0	0	_	_
Finland	4,081	0	0	_	-
France	65,469	1,955 (3.0)	414 (0.63)	131 (0.20)	283 (0.43)
Germany	25,929	254 (0.98)	6 (0.02)	1 (< 0.01)	5 (0.02)
Greece	7,743	17 (0.22)	3 (0.04)	0	3 (0.04)
Hungary	8,864	3 (0.03)	3 (0.03)	2 (0.02)	1 (0.01)
Ireland	3,416	29 (0.85)	8 (0.23)	7 (0.20)	1 (0.03)
Italy	29,281	6,341 (21.7)	7 (0.02)	2 (0.01)	5 (0.02)
Latvia	794	13 (1.6)	0	_	_
Lithuania	143	3 (2.1)	0	_	_
Luxembourg	8	1 (12.5)	1 (12.5)	0	1 (12.5)
Malta	413	43 (10.4)	20 (4.8)	1 (0.24)	19 (4.6)
Netherlands	15,587	567 (3.6)	22 (0.14)	6 (0.04)	16 (0.10)
Poland	37,701	373 (0.99)	175 (0.46)	170 (0.45)	5 (0.01)
Portugal	11,074	10 (0.09)	6 (0.05)	2 (0.02)	4 (0.04)
Romania	13,520	338 (2.5)	26 (0.19)	24 (0.18)	2 (0.01)
Slovakia	2,824	13 (0.46)	5 (0.18)	5 (0.18)	0
Slovenia	2,480	363 (14.6)	6 (0.24)	2 (0.08)	4 (0.16)
Spain	38,465	876 (2.3)	70 (0.18)	21 (0.06)	49 (0.13)
Sweden	4,077	11 (0.27)	9 (0.22)	9 (0.22)	0
United Kingdom (Northern Ireland)	7,571	6 (0.08)	0	_	-
EU Total (27 + XI)	318,794	12,040 (3.8)	901 (0.28)	463 (0.14)	438 (0.14)
Iceland	687	9 (1.3)	4 (0.58)	0	4 (0.58)
Norway	4,674	0	0	_	_
Switzerland	612	0	0	_	_



18314732, 2022, 12, Downloaded from https://dfsa.onlinelibrary.wiley.com/doi/10.2903/j.cfsa.2022.7666 by Istituo Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM). Wiley Online Library on [18/12/2023]. See the Terms

Regulation (EU) No 200/2012 requires that MSs separately report the results obtained by the FBOp and by the CA for broiler flocks. Most MSs (23) and the United Kingdom (Northern Ireland) reported both the overall merged results collected as part of the NCP and separate results from the CA and FBOp investigations, for their broiler flocks. One MS (Hungary) reported only data collected by the CA. Three MSs (Croatia, Lithuania and the Netherlands) did not comply. Considering all the data sent by the MSs providing data from both the CA and the FBOp, the EU flock prevalence of target Salmonella serovars based on CA sampling was 0.74% (N = 49,950), which was significantly higher than that based on FBOp sampling (0.24%, N = 286,741). The flock prevalence of target Salmonella serovars in broilers obtained by the CA was also significantly higher for Czechia, France, Germany, Greece, Italy, Poland and Spain. For the remaining reporting MSs, the differences between the results of the two types of samplers were not significant, or the sample sizes for one or both samplers were too small to be analysed (Table 23).

**Table 23:** Comparisons of the prevalence of target *Salmonella* serovar-positive broiler flocks, by sampler and by reporting MS, EU, 2021

	Compe	tent authoi	rity (CA)	Food	business ( (FBOp)			
Country	N tested flocks	N (%) flocks positive for target serovars	CI <sub>95</sub>	N tested flocks	N (%) flocks positive for target serovars	CI <sub>95</sub>	p-value <sup>(b)</sup>	Interpretation
Austria	115	0	[0; 3.2] <sup>(a)</sup>	6,041	9 (0.15)	[0.07; 0.28]	NS	
Belgium	89	1 (1.1)	[0.03; 6.1]	11,190	30 (0.27)	[0.18; 0.38]	NS	
Bulgaria	260	0	[0; 1.4] <sup>(a)</sup>	8,222	0	[0; 0.04] <sup>(a)</sup>	NS	
Cyprus	9	0	_	1,033	0	[0; 0.36] <sup>(a)</sup>	_	_
Czechia	41	8 (19.5)	[8.8; 34.9]	4,906	62 (1.3)	[0.97; 1.6]	< 0.001	CA > FBOp
Denmark	276	0	[0; 1.3] <sup>(a)</sup>	3,758	4 (0.11)	[0.03; 0.27]	NS	
Estonia	642	0	[0; 0.57] <sup>(a)</sup>	254	0	[0; 1.4] <sup>(a)</sup>	NS	
Finland	498	0	[0; 0.74] <sup>(a)</sup>	3,583	0	[0; 0.10] <sup>(a)</sup>	NS	
France	559	14 (2.5)	[1.4; 4.2]	65,212	400 (0.61)	[0.56; 0.68]	< 0.001	CA > FBOp
Germany	282	5 (1.8)	[0.58; 4.1]	25,929	3 (0.01)	[0; 0.03]	< 0.001	CA > FBOp
Greece	87	3 (3.4)	[0.72; 9.7]	7,743	0	[0; 0.05] <sup>(a)</sup>	< 0.001	CA > FBOp
Hungary <sup>(c)</sup>	8,864	0	[0; 0.04] <sup>(a)</sup>	_	_	_	_	_
Ireland	82	0	[0; 4.4] <sup>(a)</sup>	3,416	8 (0.23)	[0.10; 0.46]	NS	
Italy	419	2 (0.48)	[0.06; 1.7]	29,228	5 (0.02)	[0.01; 0.04]	0.004	CA > FBOp
Latvia	3	0	[-]	791	0	[0; 0.47] <sup>(a)</sup>	-	_
Luxembourg	3	1 (33.3)	[-]	8	0	[-]	_	_
Malta	6	1 (16.7)	[-]	407	19 (4.7)	[2.8; 7.2]	_	_
Poland	41,409	296 (0.72)	[0.64; 0.80]	35,535	34 (0.10)	[0.07; 0.13]	< 0.001	CA > FBOp
Portugal	111	0	[0; 3.3] <sup>(a)</sup>	11,060	6 (0.05)	[0.02; 0.12]	NS	
Romania	355	1 (0.28)	[0.01; 1.6]	13,165	25 (0.19)	[0.12; 0.28]	NS	
Slovakia	36	0	[0; 9.7] <sup>(a)</sup>	2,807	5 (0.18)	[0.06; 0.42]	NS	
Slovenia	41	0	[0; 8.6] <sup>(a)</sup>	2,439	6 (0.25)	[0.09; 0.54]	NS	
Spain	449	8 (1.8)	[0.77; 3.5]	38,405	64 (0.17)	[0.13; 0.21]	< 0.001	CA > FBOp
Sweden	139	0	[0; 2.6] <sup>(a)</sup>	4,077	9 (0.22)	[0.10; 0.42]	NS	
United Kingdom (Northern Ireland)	39	0	[0; 9.0] <sup>(a)</sup>	7,532	0	[0; 0.05] <sup>(a)</sup>	NS	



Country	Compe	Competent authority (CA)			Food business operator (FBOp)			
	N tested flocks	N (%) flocks positive for target serovars	CI <sub>95</sub>	N tested flocks	N (%) flocks positive for target serovars	CI <sub>95</sub>	p-value <sup>(b</sup>	Interpretation
EU Total (27 + XI)	54,814	340 (0.62)	[0.56; 0.69]	286,741	689 (0.24)	[0.22; 0.26]	< 0.001	CA > FBOp
EU Total (27 + XI) providing CA and FBOp data	49,950	340 (0.74)	[0.66; 0.82]	286,741	689 (0.24)	[0.22; 0.26]	< 0.001	CA > FBOp

<sup>-:</sup> Data not reported.

# Breeding flocks of turkeys

For breeding turkeys, 12 MSs and two non-MSs reported *Salmonella* NCP data. Hungary did not provide data to assess achievement of the reduction target. Considering merged data from the CA and FBOp, *Salmonella* was found in 48 of the 1,219 flocks tested (3.9%), compared with 5.1% in 2020 and 5.2% in 2019. In 2021, the prevalence of flocks positive for either of the two target *Salmonella* serovars was 0.49% (six positive flocks), compared with 0.48% and 0.30% in 2020 and 2019, respectively. Five of these positive flocks were notified by France and one by Germany (Table 24). France did not meet the reduction target of 1% or less of breeding flocks of turkeys positive for *S*. Enteritidis and/or *S*. Typhimurium (Table 24).

**Table 24:** Salmonella in breeding flocks of turkeys during the production period (flock-based data) in countries running control programmes in accordance with Regulation (EC) No 2160/2003, 2021

	N tested	N (%) positive for	N (%) positive for	N (%) pos	itive samples for
Country	flocks	Salmonella spp.	target serovars	S. Enteritidis	S. Typhimurium
Bulgaria	2	0	0	-	_
Croatia	1	0	0	_	_
Finland	7	0	0	_	-
France	481	18 (3.7)	5 (1.0)	2 (0.42)	3 (0.62)
Germany	86	1 (1.2)	1 (1.2)	0	1 (1.2)
Greece	9	1 (11.1)	0	_	_
Ireland	4	2 (50.0)	0	_	_
Italy	293	23 (7.8)	0	_	_
Poland	178	0	0	_	_
Slovakia	68	0	0	_	_
Spain	86	3 (3.5)	0	_	_
Sweden	4	0	0	_	_
EU Total (27 + XI)	1,219	48 (3.9)	6 (0.49)	2 (0.16)	4 (0.33)
Iceland	5	0	0	_	_
Norway	15	0	0	_	-

<sup>[-]:</sup> The confidence interval is not provided because of the small sample size.

<sup>(</sup>a): One-sided, 97.5% confidence interval.

<sup>(</sup>b): p-value: NS, not significant.

<sup>(</sup>c): Considering merged CA and FBOp data, Hungary notified three flocks positive for target serovars (Table 22), but in the context of separate reporting, data from the FBOp were not provided.

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903j.efsa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons

In accordance with Regulation (EC) No 1190/2012, Salmonella NCP monitoring data for breeding turkey flocks must be reported separately for sampling performed by the CA and the FBOp, in addition to the overall merged data. Eleven MSs complied with this requirement, whereas Croatia did not report separate results from the CA and the FBOp and Hungary only reported results from the CA. Considering all the data sent by those MSs that provided data from both samplers (CA and FBOp), the EU prevalence of target Salmonella serovar-positive flocks based on CA sampling was 0.70% (N = 711), which was significantly higher than that based on FBOp sampling (0.08%, N = 1,210). This finding was markedly influenced by the data reported by one single MS, France (Table 25), which was the only MS with a reported prevalence of target Salmonella serovar-positive flocks by CA significantly higher than that reported by FBOp.

**Table 25:** Comparisons of the prevalence of target *Salmonella* serovar-positive flocks of breeding turkeys, by sampler and by reporting MS, EU, 2021

	Compe	etent autho	ority (CA)	Food	business o <sub>l</sub> (FBOp)	perator		
Country	N tested flocks	N (%) flocks positive for target serovars	CI <sub>95</sub>	N tested flocks	N (%) flocks positive for target serovars	CI <sub>95</sub>	p-value <sup>(b)</sup>	Interpretation
Bulgaria	1	0	[-]	1	0	[-]	_	_
Finland	7	0	[-]	7	0	[-]	_	_
France	188	4 (2.1)	[0.58; 5.4]	481	1 (0.21)	[0; 1.2]	0.023	CA > FBOp
Germany	74	1 (1.4)	[0.03; 7.3]	83	0	[0; 4.3] <sup>(a)</sup>	NS	
Greece	5	0	[-]	9	0	[-]	_	_
Hungary	105	0	[0; 3.5] <sup>(a)</sup>	_	_	_	_	_
Ireland	4	0	[-]	4	0	[-]	_	_
Italy	163	0	[0; 2.2] <sup>(a)</sup>	293	0	[0; 1.3] <sup>(a)</sup>	NS	
Poland	154	0	[0; 2.4] <sup>(a)</sup>	178	0	[0; 2.1] <sup>(a)</sup>	NS	
Slovakia	61	0	[0; 5.9] <sup>(a)</sup>	68	0	[0; 5.3] <sup>(a)</sup>	NS	
Spain	50	0	[0; 7.1] <sup>(a)</sup>	82	0	[0; 4.4] <sup>(a)</sup>	NS	
Sweden	4	0	[-]	4	0	[-]	_	_
EU Total (27 + XI)	816	5 (0.61)	[0.20; 1.4]	1,210	1 (0.08)	[0; 0.46]	0.042	CA > FBOp
EU Total (27 + XI) providing CA and FBOp data	711	5 (0.70)	[0.23; 1.6]	1,210	1 (0.08)	[0; 0.46]	0.029	CA > FBOp

<sup>-:</sup> Data not reported.

#### Flocks of fattening turkeys

For fattening turkey flocks, 22 MSs, the United Kingdom (Northern Ireland) and three non-MSs provided data. In the EU in 2021, considering merged data from the CA and FBOp, Salmonella was found in 3,012 (9.1%) fattening turkey flocks, compared with 8.8% and 5.8% in 2020 and 2019, respectively. The EU prevalence of flocks positive for either of the two target Salmonella serovars was 0.31%, compared with 0.38% in 2020 and 0.24% in 2019. Belgium did not meet the reduction target of 1% (Figure 4), as in previous years. The EU flock prevalence was higher for S. Typhimurium (0.22%, 73 flocks) than for S. Enteritidis (0.09%, 28 flocks), with 63.0% and 53.6% of the flocks positive for S. Typhimurium and S. Enteritidis being reported by France, which was similar to the situation in previous years (Table 26).

<sup>[-]:</sup> The confidence interval is not provided because of the small sample size.

<sup>(</sup>a): One-sided, 97.5% confidence interval.

<sup>(</sup>b): p-value: NS, not significant.



18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilattico

Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [1812/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common.

**Table 26:** Salmonella in fattening flocks of turkeys before slaughter (flock-based data) in countries running control programmes, 2021

Carreton	N tested	N (%) positive for	N (%) positive for	N (%) positi	ve samples for
Country	flocks	Salmonella spp.	target serovars	S. Enteritidis	S. Typhimurium
Austria	470	14 (3.0)	3 (0.64)	3 (0.64)	0
Belgium	188	8 (4.3)	6 (3.2)	1 (0.53)	5 (2.7)
Bulgaria	4	0	0	_	_
Croatia	439	53 (12.1)	4 (0.91)	2 (0.46)	2 (0.46)
Cyprus	7	3 (42.9)	0	_	_
Czechia	260	2 (0.77)	1 (0.38)	1 (0.38)	0
Denmark	115	0	0	_	_
Finland	302	0	0	_	_
France	6,219	262 (4.2)	61 (0.98)	15 (0.24)	46 (0.74)
Germany	4,538	16 (0.35)	10 (0.22)	2 (0.04)	8 (0.18)
Greece	50	2 (4.0)	0	_	_
Hungary	1,874	5 (0.27)	5 (0.27)	1 (0.05)	4 (0.21)
Ireland	553	1 (0.18)	0	_	_
Italy	5,467	1,881 (34.4)	1 (0.02)	0	1 (0.02)
Netherlands	567	11 (1.9)	1 (0.18)	1 (0.18)	0
Poland	5,980	14 (0.23)	4 (0.07)	1 (0.02)	3 (0.05)
Portugal	1,191	5 (0.42)	1 (0.08)	0	1 (0.08)
Romania	274	1 (0.36)	0	_	_
Slovakia	98	1 (1.0)	0	_	_
Slovenia	104	9 (8.7)	1 (0.96)	0	1 (0.96)
Spain	4,041	724 (17.9)	3 (0.07)	1 (0.03)	2 (0.05)
Sweden	175	0	0	_	-
United Kingdom (Northern Ireland)	55	0	0	_	_
EU Total (27 + XI)	32,971	3,012 (9.1)	101 (0.31)	28 (0.09)	73 (0.22)
Iceland	31	0	0	_	_
Norway	235	0	0 –		_
Switzerland	38	0	0	_	_

Salmonella NCP monitoring data for fattening turkey flocks must be reported separately for sampling performed by the CA and the FBOp, in addition to the overall merged results, as defined in Regulation (EU) No 1190/2012. Nineteen MSs and the United Kingdom (Northern Ireland) complied with the requirement, whereas the two MSs Croatia and the Netherlands did not report separate results from the CA and the FBOp and the MS Hungary only reported results from the CA. Considering all the data sent by those MSs that provided data from both samplers (CA and FBOp), the EU prevalence of target Salmonella serovar-positive flocks based on CA sampling was 1.3% (N = 1,161), which was significantly higher than that based on FBOp sampling (0.26%, N = 29,840). The same finding was also seen for data provided by Germany, Italy, Poland and Spain (Table 27).



onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico

Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

**Table 27:** Comparisons of the prevalence of target *Salmonella* serovar-positive flocks of fattening turkeys, by sampler and by reporting MS, EU, 2021

turkeys, by sampler and by reporting MS, EU, 2021									
	Com	petent aut	hority (CA)	Food bu	usiness ope	rator (FBOp)			
Country	N tested flocks	N (%) flocks positive for target serovars	CI <sub>95</sub>	N tested flocks	N (%) flocks positive for target serovars	CI <sub>95</sub>	p-value <sup>(b)</sup>	Interpretation	
Austria	22	0	[0; 15.4] <sup>(a)</sup>	462	3 (0.65)	[0.13; 1.9]	NS		
Belgium	4	1 (25.0)	_	188	5 (2.7)	[0.87; 6.1]	_	_	
Bulgaria	1	0	[-]	3	0	[-]	_	_	
Cyprus	3	0	[-]	6	0	[-]	_	_	
Czechia	16	0	[0; 20.6] <sup>(a)</sup>	250	1 (0.40)	[0.01; 2.2]	NS		
Denmark	106	0	[0; 3.4] <sup>(a)</sup>	115	0	[0; 3.2] <sup>(a)</sup>	NS		
Finland	40	0	[0; 8.8] <sup>(a)</sup>	262	0	[0; 1.4] <sup>(a)</sup>	NS		
France	150	2 (1.3)	[0.16; 4.7]	6,171	59 (0.96)	[0.73; 1.2]	NS		
Germany	160	6 (3.8)	[1.4; 8.0]	4,538	5 (0.11)	[0.04; 0.26]	< 0.001	CA > FBOp	
Greece	7	0	[-]	49	0	[0; 7.3] <sup>(a)</sup>	_	_	
Hungary <sup>(c)</sup>	1,874	0	[0; 0.20] <sup>(a)</sup>	_	_	-	_	_	
Ireland	16	0	[0; 20.6] <sup>(a)</sup>	553	0	[0; 0.66] <sup>(a)</sup>	NS		
Italy	111	1 (0.90)	[0.02; 4.9]	5,462	0	[0; 0.07] <sup>(a)</sup>	0.020	CA > FBOp	
Poland	350	3 (0.86)	[0.18; 2.5]	5,891	1 (0.02)	[0; 0.10]	< 0.001	CA > FBOp	
Portugal	14	0	[0; 23.2] <sup>(a)</sup>	1,191	1 (0.08)	[0; 0.47]	NS		
Romania	28	0	[0; 12.3] <sup>(a)</sup>	246	0	[0; 1.5] <sup>(a)</sup>	NS		
Slovakia	7	0	[-]	98	0	[0; 3.7] <sup>(a)</sup>	_	_	
Slovenia	6	0	[-]	98	1 (1.0)	[0.03; 5.6]	_	_	
Spain	77	2 (2.6)	[0.32; 9.1]	4,031	2 (0.05)	[0.01; 0.18]	0.002	CA > FBOp	
Sweden	39	0	[0; 9.0] <sup>(a)</sup>	175	0	[0; 2.1] <sup>(a)</sup>	NS		
United Kingdom (Northern Ireland)	4	0	[-]	51	0	[0; 7.0] <sup>(a)</sup>	_	-	
<b>EU Total</b>	3,035	35 (0.49)	[0.28; 0.81]	29,840	78 (0.26)	[0.21; 0.33]	0.011	CA > FBOp	
(27 + XI)									
EU Total (27 + XI) providing CA and FBOp data	·	15 (1.3)	[0.72; 2.1]	29,840	78 (0.26)	[0.21; 0.33]	< 0.001	CA > FBOp	

<sup>-:</sup> Data not reported.

### Salmonella prevalence trends in poultry flocks

Trends in the estimated EU prevalence of poultry flocks positive for *Salmonella* spp. and target *Salmonella* serovars, for different poultry populations, since the implementation of the EU-wide 2007–2021 NCP, are displayed in (Figure 5). In 2020, data provided by the United Kingdom were not considered, whereas in 2021, data provided by the United Kingdom (Northern Ireland) were taken into account.

In the file 'Salmonella poultry outcome trend analyses' published on the EFSA Knowledge Junction on Zenodo here, the EU percentages of positive flocks for Salmonella, target and non-target Salmonella serovars and S. Enteritidis over time are shown and compared for each poultry population

<sup>[-]:</sup> The confidence interval is not provided because of the small sample size.

<sup>(</sup>a): One-sided, 97.5% confidence interval.

<sup>(</sup>b): p-value: NS, not significant.

<sup>(</sup>c): Considering merged CA and FBOp results, Hungary notified five flocks positive for target serovars (Table 26), but in the context of separate reporting, data from the FBOp were not provided.

covered by the NCP. Moreover, figures show the modelling of prevalence trends for *Salmonella* and target *Salmonella* serovars in poultry flocks. Detailed outputs of trend analyses (at subject level and population level) are reported.

The apparent discrepancy between the proportion of positive flocks (both for target *Salmonella* serovars and for *Salmonella*, as described in the previous paragraphs) and the estimated prevalence shown below is due to the fact that the first value is the ratio of all positive to all tested flocks, whereas the estimated prevalence is obtained by modelling the ratio of positive to all tested flocks in each reporting country, taking into account inter-country variability and the correlation between years.

Breeding flocks of Gallus gallus: 2007-2021

Since the beginning of the NCP, there has been an overall decreasing trend for the prevalence of breeding *Gallus gallus* flocks positive for target serovars (Figure 5). The prevalence estimated by modelling decreased from 1% CI<sub>95</sub> [0.57; 1.8] in 2007 to 0.38% CI<sub>95</sub> [0.27; 0.52] in 2016, when the estimated prevalence reached its lowest value. Over the next 5 years, the estimated prevalence slightly increased, reaching 0.41% CI<sub>95</sub> [0.22; 0.76] in 2021, but this increase was not statistically significant.

The estimated EU prevalence of *Salmonella*-positive breeding flocks was 1.6%  $CI_{95}$  [0.92; 2.7] in 2007 and then decreased, reaching the minimum value of 0.98%  $CI_{95}$  [0.65; 1.5] in 2016. During the following years, it increased slightly to reach 1.4%  $CI_{95}$  [0.85; 2.3] in 2021, but this increase was not statistically significant.

Flocks of laying hens: 2008–2021

Since the beginning of the NCP, there has been an overall decreasing trend for the prevalence of flocks positive for target serovars (Figure 5). The prevalence estimated by modelling was 3.6%  $\rm CI_{95}$  [2.4; 5.4] in 2008 and decreased to reach the lowest value of 0.90%  $\rm CI_{95}$  [0.65; 1.2] in 2013, with a steep downturn. From 2014 onwards, it increased slightly and stabilised at 1.1%  $\rm CI_{95}$  [0.71; 1.7] in 2021. This prevalence was not significantly different from that of the previous 2 years or compared with the lowest prevalence estimated in 2013.

The estimated EU prevalence of *Salmonella* spp. in laying hen flocks was 7.0%  $CI_{95}$  [4.3; 11.3] in 2008 and decreased to 2.1%  $CI_{95}$  [1.4; 3.2] in 2014, with a steep downturn. During the following years, it increased and reached 2.6%  $CI_{95}$  [1.7; 4.0] in 2021. In 2021, the estimated *Salmonella* prevalence in laying hen flocks was not significantly different from that in the previous 2 years or compared with the lowest prevalence estimated in 2014.

Broiler flocks: 2009–2021

From the beginning of the NCP, the flock prevalence of target serovars estimated by the model steeply decreased in the first time interval (until 2011) and then further decreased (Figure 5). The estimated prevalence was 0.44%  $CI_{95}$  [0.22; 0.87] in 2009 and decreased to 0.18%  $CI_{95}$  [0.11; 0.30] in 2021. This prevalence was not significantly different from that during the previous 2 years.

The EU prevalence of *Salmonella* spp.-positive broiler flocks estimated by modelling decreased from 2.6% CI<sub>95</sub> [1.3; 5.3] in 2009 to 1.1% CI<sub>95</sub> [0.59; 2.0] in 2015 and then increased to 1.5% CI<sub>95</sub> [0.68; 3.3] in 2021. Nevertheless, the estimated EU prevalence of *Salmonella*-positive broiler flocks in 2021 was not significantly different to that of the previous 2 years or that of 2015, when the estimated prevalence reached its lowest value.

Breeding turkey flocks: 2010–2021

From the beginning of the NCP, the prevalence of target Salmonella serovar-positive breeding turkey flocks fluctuated between 0.26%  $CI_{95}$  [0.08; 0.78] in 2010 and 0.51%  $CI_{95}$  [0.22; 1.2] (Figure 5) in 2021. This trend may have been affected by the low number of MSs with breeding turkey flocks positive for target Salmonella serovars.

With regard to EU-level Salmonella spp.-positive breeding turkey flocks, after an initial fluctuation in the EU prevalence from 7.8%  $CI_{95}$  [3.5; 16.4] in 2010 to 1.3%  $CI_{95}$  [0.69; 2.6] in 2016, when this estimated prevalence reached the lowest value seen in the entire study period, the estimated prevalence increased over time to reach 3.6%  $CI_{95}$  [2.0; 6.4] in 2021. This estimated prevalence in 2021 was not significantly different from that of the previous 2 years, but it was significantly higher than the estimated prevalence in 2016 (p-value = 0.0261).

18314732, 2022, 12, Downloaded from https://fs.aonlinelibrary.wiley.com/continelibrary.wiley.com

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico


Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Comm

#### Fattening turkey flocks: 2010–2021

The estimated flock prevalence of target serovars was 0.38%  $CI_{95}$  [0.23; 0.62] in 2010; it decreased to 0.25%  $CI_{95}$  [0.17; 0.35] in 2014 and then increased to 0.28%  $CI_{95}$  [0.13; 0.59] in 2021, after some small temporal fluctuations (Figure 5). Nevertheless, there were no significant differences in the estimated prevalence of the target *Salmonella* serovars in EU fattening turkey flocks in the last 2 years or compared with the lowest prevalence estimated in 2014.

For this poultry category, after an initial fluctuation in the EU prevalence of *Salmonella* spp.-positive flocks from 6.1%  $CI_{95}$  [3.6; 10.0] in 2010 to 2.0%  $CI_{95}$  [0.96; 4.3] in 2016, when the estimated prevalence reached its lowest value, the prevalence increased to 4.1%  $CI_{95}$  [1.6; 10.0] in 2021. Nevertheless, the prevalence in 2021 was not significantly different from that in the previous 2 years or from the lowest estimated prevalence in 2016.



**Figure 5:** Trend in the estimated prevalence of poultry flocks positive for *Salmonella* spp. and target *Salmonella* serovars, at EU level for different poultry populations, 2007–2021

#### Salmonella data for other animals

Considering all the collected data on the presence of *Salmonella* in different categories of animal species in the EU with the exception of data collected in the framework of NCPs for poultry, 70,326 samples collected from animals of various species were reported by 20 MSs (Table 28). The overall prevalence of *Salmonella*-positive samples was 4.0% (N = 2,843). The highest number of samples was from cattle (bovine animals) (N = 26,412 notified by 14 MSs) and 3.5% were reported as being positive for *Salmonella*. The highest prevalence of positive samples were notified for small ruminants (9.0% for nine MSs) and wild boar (6.6% for three MSs). For pigs, based on data reported by 17 MSs, the prevalence of positive samples was 2.9% (520 positive samples,

N = 17,927). For solipeds, 2.9% of samples were positive for *Salmonella* (18 positive samples) and were notified by seven MSs.

**Table 28:** Summary of *Salmonella* statistics related to major animal species, reporting MSs and non-MS countries, 2021

		EU <sup>(a)</sup> MSs		Non-MS countries				
Animals	N reporting	N tested	Positive animals		N reporting	N tested	Positive animals	
	countries	animals	N	%	countries	animals	N	%
Birds <sup>(b)</sup>	12	13,150	480	3.7	3	336	21	6.3
Cats	5	1,072	50	4.7	2	559	20	3.6
Cattle (bovine animals)	14	26,412	920	3.5	4	5,390	242	4.5
Dogs	8	1,995	53	2.7	2	1,082	46	4.3
Pigs	17	17,927	520	2.9	3	4,517	151	3.3
Small ruminants <sup>(c)</sup>	9	5,622	505	9.0	3	262	28	10.7
Solipeds	7	616	18	2.9	2	364	11	3.0
Wild boar	3	1,175	78	6.6	1	287	13	4.5
Wild ungulates <sup>(d)</sup>	3	388	2	0.52	1	1	0	0
Others/Not specified <sup>(e)</sup>	13	1,969	217	11.0	2	422	68	16.1
Total	20	70,326	2,843	4.0	5	13,220	600	4.5

MSs: Member States.

- (c): Included in this category: sheep and goats.
- (d): Included in this category: deer and Cantabrian chamois.
- (e): Included in this category: shellfish, quails, fish, monkeys, pigeons, badgers, foxes, other animals, Leporidae, alpacas, antelopes, Barbary sheep, bats, bears, bison, camels, capybaras, chinchillas, dolphins, dromedaries, ferrets, guinea pigs, hedgehogs, jays, kangaroos, leopards, lions, lynxes, marine mammals, martens, mice, minks, mouflon, other carnivores, otter, passerines, unspecified, Psittacidae, raccoon dogs, rats, reptiles, rodents, seals, snakes, squirrels, Alpine ibex, turtles, water buffalo, wildcats (*Felis silvestris*), wolves, zoo animals, all, beavers, raccoons, ratites (ostriches, emus, nandus), other ruminants, penguins, moose, reindeer.

For a further interactive look at *Salmonella* monitoring results, dashboards have been created (*different filters can be applied to query the data*) (*link*).

#### 2.4.5. Salmonella in feed

In 2021, the overall EU-level occurrence of *Salmonella*-positive samples in any 'animal and vegetable-derived feed' was 0.55% (N = 71,965). In compound feed (finished feed for animals), the prevalence of *Salmonella*-positive units was 0.40% (N = 15,463) for samples from poultry, 0.58% for samples from cattle (N = 2,909) and 0.36% for samples from pigs (N = 4,123). There were no noticeable isolates in 2021, except for fish, where in over 134 tested units, the prevalence of *Salmonella*-positive samples was 1.5%, and for rabbits, where in 55 tested units, the prevalence was 1.8%. Regarding compound feedingstuffs for fur animals, *Salmonella* was reported in 76 out of 176 tested units (43.2%) with all the positive samples having been detected in Poland. Lastly, the prevalence of *Salmonella*-positive sampling units for pet food was 1.0% (N = 3,196). For non-specified compound feedingstuffs, *Salmonella* was reported in 0.23% of over 8,222 tested units, mainly by Sweden.

18314732, 2022, 12, Downloaded from https://fs.aonlinelibrary.wiley.com/doi/10.9293/j-fs.a.2022.7566 by Istituto Zoporofilatios Sperimenta dell'Umbria e delle Marche (ZSUM), Wiley Online Library on [1812023], See the Terms and Conditions (https://onlinelibrary.wiley.com/rerms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Cerawise Common.

<sup>(</sup>a): Data from the United Kingdom (Northern Ireland) are taken into account for 2021. In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/Northern Ireland, the EU requirements on data sampling are also applicable to Northern Ireland.

<sup>(</sup>b): Included in this category: geese, canaries, ducks, parrots, pheasants, *Gallus gallus* (fowl), owls, partridges, turkeys, doves, falcons, gulls, magpies, night herons, ostriches, peafowl, starlings, swans, poultry (unspecified), guinea fowl, budgerigars, songbirds.



#### 2.4.6. Salmonella serovars in humans, food and animals

#### **Humans**

Serovars among all confirmed salmonellosis cases

For humans, information on *Salmonella* serovars was available for 84.6% of the total number of confirmed cases (50,817 cases out of 60,050) for 25 MSs (Bulgaria, Luxembourg and Spain did not report serovar data). Data included all cases reported with serovar information regardless of travel status. The proportion of cases of *Salmonella* with serovar data available decreased compared to 2020 (87.9%). As in previous years, the three most commonly reported *Salmonella* serovars in 2021 were S. Enteritidis (54.6%), S. Typhimurium (11.4%) and monophasic S. Typhimurium (1,4,[5],12:i:-) (8.8%), representing 74.8% of the 50,817 confirmed human cases. S. Enteritidis increased by 15.5%, when considering the absolute number of cases of this serovar, but it increased by only 2.8% compared with 2020 with respect to the total number of isolates. Monophasic S. Typhimurium (1,4,[5],12:i:-) apparently decreased by 4.3% compared with 2020, when considering the absolute number of cases of this serovar in these years, but the proportion decreased by 1.3% compared with 2020 with respect to the total number of isolates in the relative years.

The proportions of these three serovars, mainly driven by S. Enteritidis, increased during the last 3 years: from 70.5% in 2019 to 72.2% in 2020 and to 74.8% in 2021. The fourth and fifth serovars, S. Infantis and S. Derby, were at the same levels as in 2020 and 2019 (Table 29).

In 2021, the proportion of cases of the S. Coeln serovar increased by 0.21% and 0.35% compared with 2020 and 2019, respectively; it increased by 42.9% compared with 2020 while it decreased by 4.9% compared with 2019, when considering the absolute number of cases of this serovar in these years. For the first time since 2019, four serovars, S. Braenderup, S. Montevideo, S. Oranienburg and S. Rissen, entered the top 20 list of the most frequent serovars in 2021. In particular, the proportion of cases of S. Braenderup increased by 0.53% and 0.36% compared with 2020 and 2019, respectively; the absolute number of cases increased by 301% and 27.7% compared with 2020 and 2019, respectively. The proportion of S. Montevideo cases increased by 0.21% and 0.12% compared with 2020 and 2019, respectively, but the absolute number of cases increased by 114.7% compared with 2020 and decreased by 10.2% compared with 2019. Furthermore, S. Virchow entered the top 20 list of the most frequent serovars in 2019.

**Table 29:** Distribution of reported confirmed cases of human salmonellosis in the EU, 2019–2021, for the 20 most frequent *Salmonella* serovars in 2021

	2	2021		2	2020		2	2019	
Serovar	Cases	MSs	%	Cases	MSs	%	Cases	MSs	%
Enteritidis	27,734	24	54.6	24,008	25	51.8	39,451	27	50.4
Typhimurium	5,781	24	11.4	5,337	25	11.5	9,288	27	11.9
Monophasic Typhimurium 1.4.[5] 0.12:i:-	4,495	14	8.8	4,697	16	10.1	6,432	18	8.2
Infantis	1,019	24	2.0	1,064	23	2.30	1,912	26	2.4
Derby	474	17	0.93	525	20	1.13	719	23	0.92
Coeln	463	15	0.91	324	19	0.70	441	18	0.56
Braenderup	373	15	0.73	93	12	0.20	292	18	0.37
Napoli	352	12	0.69	412	12	0.89	493	18	0.63
Chester	316	12	0.62	129	13	0.28	340	17	0.43
Newport	311	20	0.61	336	21	0.73	846	24	1.08
Montevideo	219	11	0.43	102	14	0.22	244	19	0.31
Brandenburg	213	15	0.42	309	16	0.67	288	17	0.37
Bovismorbificans	205	14	0.40	337	15	0.73	452	19	0.58
Oranienburg	193	14	0.38	90	14	0.19	236	19	0.30
Stanley	190	16	0.37	208	21	0.45	509	19	0.65
Virchow	170	16	0.33	121	16	0.26	469	21	0.60
Dublin	148	10	0.29	196	9	0.42	207	13	0.26
Anatum	143	11	0.28	57	7	0.12	134	15	0.17

Communication	2	2021		2	2020		2019		
Serovar	Cases	MSs	%	Cases	MSs	%	Cases	MSs	%
Rissen	139	14	0.27	114	13	0.25	252	19	0.32
Agona	137	16	0.27	157	18	0.34	490	20	0.63
Other	7,742		15.2	7,725		16.7	14,787		18.9
Total <sup>(a),(b)</sup>	50,817	24	100	46,341	25	100	78,282	27	100

MSs: Member States.

- (a): Source(s): 2021 25 MSs: Austria, Belgium, Cyprus, Czechia, Germany, Denmark, Estonia, Greece, Finland, France, Croatia, Hungary, Ireland, Italy, Lithuania, Latvia, Malta, Netherlands, Poland, Portugal, Romania, Sweden, Slovenia, Slovakia. 2020 25 MSs: Austria, Belgium, Cyprus, Czechia, Germany, Denmark, Estonia, Greece, Finland, France, Croatia, Hungary, Ireland, Italy, Latvia, Luxembourg, Malta, Netherlands, Poland, Portugal, Romania, Spain, Sweden, Slovenia, Slovakia. 2019 27 MSs: Austria, Belgium, Cyprus, Czechia, Germany, Denmark, Estonia, Greece, Finland, France, Croatia, Hungary, Ireland, Italy, Lithuania, Latvia, Luxembourg, Malta, Netherlands, Poland, Portugal, Romania, Spain, Sweden, Slovenia, Slovakia the United Kingdom.The total number of confirmed cases with information on Salmonella serovars includes cases classified as S. enterica, S. SubspI, S. SubspIIIb, S. SubspIIIa, S. SubspIV, S. SubspIV, S. SubspV, S. SubspVI and other.
- (b): Data from the United Kingdom are taken into account for the year 2019, because the United Kingdom was an EU MS, but it became a third country on 1 February 2020.

#### Serovars acquired in the EU

To estimate the impact of *Salmonella* infections acquired at the EU level, serovar data were analysed for domestic and travel-associated cases in which the probable country of infection was an EU MS. Information on *Salmonella* serovars with travel data was available from 23 MSs, representing 72.0% of cases with known serovar data in 2021. Most cases (97.7%) with a known serovar and with travel data were infected within the EU. For the travel-associated cases, the most frequently reported travel destinations in the EU were Spain (16.6%), Italy (15%), Greece (11.4%) and Poland (10.2%). For the reported cases of human salmonellosis acquired in the EU, *S.* Enteritidis dominated and 64.6% of these reported cases were infected with this serovar. *S.* Enteritidis, *S.* Typhimurium and monophasic *S.* Typhimurium (1,4,[5],12:i:-) together represented 79.1% of the confirmed human cases acquired in the EU in 2021 (Table 30). *S.* Enteritidis cases were predominantly (85.2%) infected within the EU. The proportions of cases of *S.* Enteritidis, *S.* Typhimurium and its monophasic variant (1,4,[5],12:i:-) were at the same level as in 2019–2020, when considering the number of these serovars vs. the total number of reported serovars. Also, *S.* Infantis and *S.* Derby remained at the same level as in 2020.

**Table 30:** Distribution of reported confirmed cases of human salmonellosis acquired in the EU, 2019–2021, for the six most frequent *Salmonella* serovars in 2021

_	2	2021		2020			2019		
Serovar	Cases	MSs	%	Cases	MSs	%	Cases	MSs	%
Enteritidis	23,634	23	64.6	21,203	23	63.1	32,010	24	61.6
Typhimurium	4,027	23	11.0	3,702	22	11.0	6,044	24	11.6
Monophasic Typhimurium 1.4.[5] 0.12:I:-	1,269	14	3.5	1,530	16	4.6	2,688	17	5.2
Infantis	633	23	1.7	716	21	2.1	1,215	24	2.3
Derby	239	16	0.65	260	17	0.77	396	20	0.76
Coeln	315	14	0.86	201	17	0.60	270	15	0.52
Other	6,462	_	17.7	6,009	_	17.9	9,378	_	18.0
Total <sup>(a)</sup>	36,579	23	100	33,621	23	100	52,001	24	100

MSs: Member States.


(a): Data from the United Kingdom are taken into account for the year 2019, because the United Kingdom was an EU MS, but it became a third country on 1 February 2020.

A seasonal trend was observed for confirmed S. Enteritidis infections acquired in the EU in 2017–2021, with more cases reported during summer months. A decrease in cases was observed in 2020, probably due to the COVID-19 pandemic. Notwithstanding, the overall trend for S. Enteritidis in 2017–2021 did not show any statistically significant increase or decrease (Figure 6). Belgium, Estonia and Greece showed a significantly decreasing (p < 0.05) trend in S. Enteritidis infections within

18314732, 2022, 12, Downdoaded from https://disa.onlinelibitrary.wiley.com/doi/10/2033/gfsa2022/7666 by Istitute Zoperinented ell'Umbia e delle Marche (ZSUM), Wiley Online Library on [1812/2023], See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10/2033/gfsa2022/7666 by Istitute Zoperinented by the applicable Cereive Common



the EU over the last 5 years (2017–2021). A significant increasing trend (p < 0.05) was observed in France and Malta.



Source: Austria, Belgium, Czechia, Germany, Denmark, Estonia, Greece, Finland, Hungary, Ireland, Italy, Latvia, Malta, the Netherlands, Portugal, Sweden and Slovakia.

**Figure 6:** Trend in reported confirmed human cases of *S*. Enteritidis infections acquired in the EU, by month, 2017–2021

#### Food and animals

Descriptive analyses were undertaken using serotyped isolates from food and animals. In this context, only isolates related to the most common food-producing animal species and food matrices thereof were considered and were aggregated into the following categories for further analysis: 'broiler flocks – broiler meat', 'laying hen flocks – eggs', 'fattening turkey flocks – turkey meat', 'fattening pigs – pig meat' and 'cattle – bovine meat'. Overall, a selection of 20,020 serotyped *Salmonella* isolates meeting the mentioned inclusion criteria were obtained (Table 31).

**Table 31:** Distribution of *Salmonella* isolates (number and percentage of positive sampling units) with and without serotype identification among the different selected sources (food and animals), EU, 2021

Source	Salmonella pos units without se					
	N	%	N	%		
Broilers	973	14.0	11,143	55.7		
Broiler meat	2,905	41.7	3,206	16		
Cattle	30	0.43	124	0.62		
Cattle meat	138	2.0	261	1.3		
Pigs	71	1.0	394	2.0		
Pig meat	1,545	22.2	1,114	5.6		
Turkeys	663	9.5	2,373	11.9		

18314732, 2022, 12, Downloaded from https://eis.an.onlinelibrary.wiley.com/lenbra/wiley.com/doi/10.2033/jefsa.2022.7666 by. Istituto Zooporfollatios Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023], See the Terms and Conditions (https://onlinelibrary.wiley.com/lenms-and-conditions) on Wiley Online Library for rules of use; OA atticles are goverend by the applicable Cerative Commons.



18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprefilatios Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions


Source		sitive-sampling erotyped isolates	Salmonella positive-sampling units with serotyped isolates			
	N	%	N	%		
Turkey meat	395	5.7	203	1.0		
Layers	198	2.8	1,166	5.8		
Eggs and egg products	43	0.62	36	0.18		
Total	6,961	100	20,020	100		

The large majority of the serotyped isolates were from 'broilers' (both animals (55.7%) and food (16.0%)). 'Turkey' sources (animals and food) accounted for 12.9% of the serotyped isolates, while 'pig' and 'laying hen' sources represented 7.6% and 6.0% of the serotyped isolates, respectively. Serotyped isolates from 'cattle' sources made up about 1.9% of the total.

Isolates belonging to the five most frequently reported *Salmonella* serovars involved in cases of human salmonellosis acquired in the EU in 2021 were considered for further analysis, which were S. Enteritidis, S. Typhimurium, monophasic S. Typhimurium (1,4,[5],12:i:-), S. Infantis and S. Derby. From the above-mentioned food-animal sources a total of 20,020 serotyped isolates were reported, from which S. Infantis accounted for 33.9%, S. Enteritidis for 8.2%, S. Typhimurium for 3.8%, monophasic S. Typhimurium (1,4,[5],12:i:-) for 3.2% and S. Derby for 2.2%.

A Sankey diagram (Figure 7) illustrates how these top five EU-level *Salmonella* serovars involved in human salmonellosis cases acquired in the EU were linked with the major animal species.

S. Enteritidis was primarily related to 'broiler' sources (70.0% of the S. Enteritidis isolates were from broiler flocks and meat) and also to 'layers and eggs' (26.0%). S. Typhimurium isolates were distributed among the different sources, although they were mainly related to 'broiler' and 'pig' sources (43.2% and 29.7% of the isolates were from these sources respectively), followed by 'laying hen' and 'turkey' sources (15.0% and 9.1%). Monophasic S. Typhimurium (1,4,[5],12:i:-) was related mainly to 'pig' (65.4%) and secondly to 'broiler' (21.2%) sources. S. Infantis was strictly related to 'broiler' sources (95.2%). S. Derby was primarily related to 'pig' (75.3%) and secondly to 'turkey' (13.5%) sources. To interpret these data, it is important to be aware that the distribution of the serotyped isolates among the different sources is very unbalanced in terms of the number of isolates per source, and the large majority of the serotyped isolates for the subsets considered were from poultry populations covered by NCPs, especially broilers.



The left side of the diagram shows the five most commonly reported *Salmonella* serovars involved in human salmonellosis cases acquired in the EU: *S.* Enteritidis (blue), *S.* Infantis (green), *S.* Typhimurium (orange), monophasic *S.* Typhimurium (1,4,[5],12:i:-) (violet) and *S.* Derby (magenta). Animal and food data from the same source were merged: 'broiler' includes isolates from broiler flocks and broiler meat, 'bovine' includes isolates from bovines for meat production and from bovine meat, 'pig' includes isolates from fattening pigs and pig meat, 'turkey' includes isolates from fattening turkey flocks and turkey meat and 'layers' includes isolates from laying hen flocks and eggs. The right side shows the five sources considered (broilers, bovines, pigs, turkeys and layers). The width of the coloured bands linking the sources and serovars is proportional to the percentage of isolates of each serovar from each source.

**Figure 7:** Sankey diagram of the distribution of the top five EU-level *Salmonella* serovars involved in human salmonellosis cases acquired in the EU, reported from specified food–animal categories, by food–animal source, EU, 2021

Table 32 shows the top 20 serovars notified, considering all serotyped isolates (including those from both food and animals) from the following species: laying hens, broilers, turkeys, pigs and bovines. For laying hens, 44.9% of the isolates (N = 1,198) belonged to two target serovars (S. Enteritidis (35.5%) and S. Typhimurium (9.4%)). The other most common serovars from this source were S. Kentucky (10.3%) and S. Infantis (7.1%). For broilers (N = 14,348), four serovars (S. Infantis, S. Enteritidis, S. Livingstone and S. Mbandaka) represented 65.9% of the serotyped isolates, with S. Infantis including more than 45% of all strains. S. Typhimurium and its monophasic variant accounted for 2.9% of the isolates from this species. Also, for turkey sources (N = 2,639), two serovars (S. Anatum (29.8%) and S. Agona (23.5%)) were by far the most common ones, followed by S. Infantis (7.2%). Overall, S. Typhimurium, S. Enteritidis and the monophasic variant of S. Typhimurium accounted for 5.6% of the isolates from this species. For pigs, the monophasic variant of S. Typhimurium (28.2%) as well as S. Derby (22.3%) represented half of the serotyped strains (N = 1,559) from this source, with S. Typhimurium (15.3%) and S. Rissen (6.6%) being two other common serovars. Considering serotyped isolates from bovine sources (N = 891), two serovars were by far the most common ones: S. Dublin and S. Typhimurium represented 31.5% and 29.7% of the serotyped isolates from this source respectively.

1831/4372, 2022, 1,2 Downloaded from https://ef.son.inlinibitary.wiley.com/oid/10/2003/jefssa2022/7666 by Islituic Copyor/Inlinib Sperimental dell'Umbria e delle Marche (ISUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://oillneibbrary.wiley.com/ems-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

Laying hens



**Table 32:** Distribution of the top 20 *Salmonella* serovars by food–animal source (laying hens, broilers, turkeys, pigs and bovines), EU, 2021 **Poultry population** 

Serovar	N (%) positive samples	N MSs
Enteritidis	425 (35.5)	22
Kentucky	124 (10.3)	5
Typhimurium	113 (9.4)	15
Enterica, subspecies enterica	97 (8.1)	3
Infantis	85 (7.1)	12
Braenderup	24 (2.0)	8
Mbandaka	20 (1.7)	8
Anatum	19 (1.6)	5
Corvallis	17 (1.4)	6
Montevideo	16 (1.3)	6
Ohio	16 (1.3)	4
Thompson	15 (1.2)	6
Newport	13 (1.1)	5
Senftenberg	13 (1.1)	6
Livingstone	11 (0.92)	5
Mikawasima	11 (0.92)	2
Agona	10 (0.83)	4
Hadar	10 (0.83)	3
Mkamba	10 (0.83)	1
Monophasic Typhimurium	9 (0.75)	1
Other	140 (11.7)	26
Total	1,198	

Serovar	N (%) positive samples	N MSs
Infantis	6,468 (45.1)	23
Enteritidis	1,147 (8.0)	21
Livingstone	998 (7.0)	7
Mbandaka	838 (5.8)	10
Java	439 (3.1)	5
Enterica, subspecies enterica	436 (3.0)	4
Thompson	374 (2.6)	7
Typhimurium	330 (2.3)	17
Montevideo	301 (2.1)	5
Group O:7	281 (2.0)	3
Senftenberg	273 (1.9)	8
Newport	171 (1.2)	15
Agona	170 (1.2)	13
Kedougou	156 (1.1)	7
enterica	150 (1.0)	3
Virchow	99 (0.69)	5
Kentucky	96 (0.67)	9
Mkamba	92 (0.64)	1
Napoli	81 (0.56)	2
Monophasic Typhimurium	81 (0.56)	2
Other	1,367 (9.5)	27
Total	14,348	

Serovar	N (%) positive samples	N MSs
Anatum	785 (29.8)	7
Agona	620 (23.5)	9
Infantis	189 (7.2)	9
Newport	94 (3.6)	9
Senftenberg	90 (3.4)	4
Enterica, subspecies enterica	81 (3.1)	1
Typhimurium	71 (2.7)	9
Derby	61 (2.3)	6
Bredeney	55 (2.1)	5
Saintpaul	55 (2.1)	4
Hadar	52 (2.0)	4
Lagos	40 (1.5)	1
Monophasic Typhimurium	40 (1.5)	2
Kedougou	38 (1.4)	3
Enteritidis	37 (1.4)	11
Coeln	32 (1.2)	5
Napoli	27 (1.0)	3
Kentucky	24 (0.91)	7
Thompson	18 (0.68)	1
Haifa	15 (0.57)	1
Other	215 (8.2)	19
Total	2,639	



## Pigs and cattle (bovine animals)

Serovar	N (%) positive samples	N MSs	
Monophasic Typhimurium	440 (28.2)	16	
Derby	347 (22.3)	21	
Typhimurium	239 (15.3)	19	
Rissen	103 (6.6)	8	
enterica	81 (5.2)	3	
Brandenburg	44 (2.8)	8	
Infantis	39 (2.5)	10	
Group B	24 (1.5)	4	
Group O:4	22 (1.4)	1	
Enterica, subspecies enterica	19 (1.2)	4	
London	19 (1.2)	7	
4,12:a:-	14 (0.9)	1	
Goldcoast	12 (0.77)	3	
Afula	10 (0.64)	1	
Enteritidis	10 (0.64)	5	
Livingstone	10 (0.64)	3	
Agona	9 (0.58)	2	
Bovismorbificans	6 (0.38)	2	
Choleraesuis	6 (0.38)	2	
Choleraesuis var. Kunzendorf	6 (0.38)	1	
Other	99 (6.3)	26	
Total	1,559		

Serovar	N (%)	N
Seloval	positive samples	MSs
Dublin	281 (31.5)	9
Typhimurium	265 (29.7)	15
Monophasic Typhimurium	80 (9.0)	7
Havana	57 (6.4)	2
enterica	28 (3.1)	3
Group B	19 (2.1)	3
Derby	18 (2.0)	8
Enteritidis	18 (2.0)	6
Infantis	14 (1.6)	7
Altona	9 (1.0)	1
Agona	6 (0.67)	2
Group C	6 (0.67)	1
Montevideo	6 (0.67)	1
Group C1	5 (0.56)	1
Mbandaka	5 (0.56)	2
Kentucky	4 (0.45)	2
Muenster	4 (0.45)	1
Newport	4 (0.45)	2
Give	3 (0.34)	1
Rissen	3 (0.34)	3
Other	56 (6.3)	21
Total	891	

MSs: Member States.

Pigs



#### 2.5. Discussion

Salmonellosis remained the second most common foodborne zoonosis in humans in the EU in 2021 after campylobacteriosis. The previous decreasing trend for confirmed cases has stabilised since 2014. In 2020, the number of reported confirmed human cases and the EU notification rate reached the lowest levels observed since the beginning of *Salmonella* surveillance at the EU level in 2007. This was probably due to the COVID-19 pandemic and the withdrawal of the United Kingdom from the EU. In 2021, a slight increase in notified human cases was registered compared with 2020, caused in part by the lower impact of the COVID-19 pandemic compared with the previous year. The slight increase in human cases of salmonellosis in 2021 could be associated with the reduction of COVID-19 pandemic restriction measures and with the slow recovery of daily activities (social events, doctor's visits, travel). The numbers of human salmonellosis cases acquired in the EU, outbreak-related cases and foodborne salmonellosis outbreaks were higher in 2021 than in 2020 but lower than in previous years, and may have been consequences of reduced COVID-19 restrictions. Notwithstanding, the overall EU trend for salmonellosis in 2017–2021 did not show any statistically significant increase or decrease. Conversely, over the period 2017–2021, Denmark, Estonia, Finland, Ireland, Romania and Sweden reported a decreasing trend.

In addition, notification rates for salmonellosis in humans varied between MSs, reflecting potential variations, for example, in the quality, coverage and disease-severity focus of the surveillance systems, sampling and testing practices, disease prevalence in the food-producing animal population and food and animal trade between MSs.

The hospitalisation rate varied from 23.9% to 91.9%. Countries reporting the lowest notification rates for salmonellosis had the highest proportions of hospitalisation, suggesting that the surveillance systems in these countries are focused on the most severe cases and underlining the variability of national surveillance systems. It is important to note that a higher hospitalisation rate was reported for patients with specimens from blood (82.4%), suggesting that people with systemic infection are more likely to be hospitalised.

In 2021, a decrease in the proportion of reported data for *Salmonella* serovars was also observed compared with 2020. This was probably due to a lack of serovar data from Spain and Luxembourg. For Spain, serovar information was not available when *Salmonella* serovar data were analysed for this report (this information will be shown in the ECDC Surveillance Atlas). In 2021, the proportion of monophasic *S.* Typhimurium (1,4,[5],12:i-) decreased and the proportion of *S.* Enteritidis increased compared with previous years. Concerning the increased proportion of *S.* Enteritidis cases, it is worth noting that Poland provided serovar data in 2021, whereas it had not provided such data in 2020; it is to be remembered that Poland had been involved, during previous years, in a long-lasting multicountry outbreak associated with contaminated eggs from Polish farms.

Regarding the cases acquired in the EU, the ranking of the five most common serovars was stable, but the proportion of S. Enteritidis was much higher than in relation to total cases. Overall, the three most commonly reported human serovars, S. Enteritidis, S. Typhimurium and its monophasic variant (1,4,[5],12:i:-) continued to account for over 70% of human cases acquired in the EU, as has been observed since 2014, reaching 79.1% in 2021. S. Infantis has consistently been the fourth most frequently reported serovar involved in domestically acquired and travel-associated human infections. After S. Infantis, S. Derby was the fifth most frequently reported serovar in 2021, while S. Napoli slid to eighth position, from sixth in 2020, preceded by S. Braenderup, probably due to a multi-country outbreak of S. Braenderup ST22 that was presumed to be linked to imported melons (ECDC and EFSA, 2021a).

S. Enteritidis continued to be the serovar most frequently reported in human cases and was associated with different sources, especially broilers and laying hens. Eggs and egg products were confirmed as the primary source of salmonellosis outbreaks in 2021, as had been the case during previous years (Chanamé Pinedo et al., 2022). The importance of eggs as a source of human salmonellosis was underlined by the recurrent high number (seven) of MSs that did not meet the Salmonella reduction target in laying hens under their NCPs, even though that target was set at 2% instead of 1% as for other poultry populations. It is also of note that for breeding flocks of Gallus gallus, a number (five) of MSs did not meet the reduction target, which had also been the case in the last few years. Moreover, trend analyses confirmed that during the last few years, there was no significant reduction in target Salmonella serovar flock prevalence in any of the poultry populations covered by NCPs. This indicates that Salmonella elimination and eradication, also in relation to those

18314732, 2022, 12, Downloaded from https://fs.aonlinelibrary.wiley.com/inchibrary.wiley.com/inchibrary.wiley.com/em-and-conditions) on Wiley Online Library on [18/12/2023], See the Terms and Conditions (http://oinineibrary.wiley.com/em-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons

serovars that are considered relevant for public health, are not in sight. These consolidated findings suggest that some revisions of programmes such as changes to the frequency of sampling on farms and the vaccination policy, especially for groups with long production cycles, could be considered in order to reduce the risk of Salmonella spreading. Broiler meat is another source of Salmonella and the contamination of broiler farms has been increasingly associated with persistent serovars, such as S. Infantis (Mughini-Gras et al., 2021). These data confirm that it is vital to identify efficient and costeffective measures to control contamination, primarily on farms, which may even include evaluating measures aimed at non-regulated Salmonella serovars (Newton et al., 2021) in relation to the epidemiological situation in each MS. Pigs are another species that deserves attention as a putative source of salmonellosis (Chanamé Pinedo et al., 2022). S. Typhimurium and its monophasic variant, which are among the top serovars causing human infections, also appear as some of the serovars most commonly associated with this animal species. In pigs, the Salmonella holding prevalence varies widely across MSs, as was ascertained with the EU baseline survey conducted in 2008 (EFSA, 2009b). Over the last few years, these differences may also have become larger due to the fact that some MSs (e.g. Norway, Finland, Sweden, Denmark, Germany and Estonia) have implemented control programmes for Salmonella in their pig populations (Bonardi et al., 2021) in an attempt to reduce the risk of this pathogen reaching humans (Ferrari et al., 2019). For bovine animals, similar to pigs, no harmonised control programmes are implemented at EU level, although some MSs have their own national control programmes in place. For this animal species, the prevalence situation is even more unclear than for pigs, because no baseline surveys have been carried out so far at EU level. Moreover, also looking at the 2021 food data and Salmonella PHC data on carcases at the slaughterhouse, it is evident how meat has continued to play a central role in the dissemination of the pathogen (one in seven broiler carcases and one in 13 turkey carcases were positive for Salmonella at the slaughterhouse). The safety of carcases and derived meat products is dependent on both farm and slaughterhouse performance (Cegar et al., 2022). Regarding poultry meat, although NCPs for poultry populations have been implemented for many years, the EU trends for different species have stabilised both for Salmonella spp. and for 'target serovars' without any significant decrease in the prevalence notified recently. Control at farm level is undeniably important for reducing the prevalence of Salmonella in the next phases of the process; however, further implementation aiming to reduce the contamination of final products could also address the slaughter process. The risk categorisation of slaughterhouses based on process hygiene criteria has been suggested as one of the essential components of the risk-based meat safety assurance system (EFSA BIOHAZ Panel, 2012). Therefore, efforts to control Salmonella at the primary production level should be combined with the categorisation of slaughterhouses in terms of their capacity to reduce carcase contamination (Cegar et al., 2022). Food chain information, accompanying animals to be admitted to the slaughterhouse, serves as the bridge between these two phases (Bonardi et al., 2021). Flocks from high-risk farms can be directed to low-risk slaughterhouses, thanks to their capacity to reduce carcase contamination. High-risk slaughterhouses could accept flocks from low-risk farms or could apply additional final carcase interventions (EFSA BIOHAZ Panel, 2012).

Regulation (EU) 2019/627 has defined the obligation, for competent authorities, to verify the correct implementation by the FBOp of PHC for *Salmonella* in carcases at the slaughterhouse. Data collected in this context confirmed that, especially for poultry (broiler and turkey) carcases, the proportion of positive samples, identified by sampling conducted by the CA, was significantly higher than that notified for samples taken by the FBOp. This discrepancy was also observed for national control programmes for poultry, where separate reporting on controls carried out by the CA and FBOp is mandatory. The prevalence of target *Salmonella* serovars in samples from controls conducted by competent authorities was consistently higher than for controls carried out by the FBOp, for both broilers and fattening turkeys. Investigations to define the reasons for discrepancies between the results of sampling conducted by the CA and FBOp should be encouraged as an essential prerequisite to ensure the trustworthiness of the data collected in both contexts.

Although animal-based foods are the most common sources of *Salmonella* transmission to humans (Ferrari et al., 2019), and while prevalence data collected in 2021 from food samples confirmed that the pathogen is seldom isolated from non-animal products, large multi-country outbreaks related to non-animal products (e.g. melons, sesame-based products and chocolate products) (ECDC and EFSA, 2021b,a, 2022a) recently occurred in the EU. Therefore, for such products, efforts to control *Salmonella* contamination must also be maintained at a high level.



18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilattico


dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

## 3. Listeria monocytogenes

Summary data substantiating this chapter, as well as additional information on related projects and internet sources are published for this report on the EFSA Knowledge Junction on Zenodo here. Summary statistics on *Listeria monocytogenes* food monitoring data and on listeriosis foodborne outbreaks reported in the framework of Directive 2003/99/EC, with downloadable files, are retrievable using the EFSA *Listeria monocytogenes* dashboard and the EFSA foodborne outbreaks dashboard, respectively available here and here. Summary statistics on human surveillance data with downloadable files are retrievable using the ECDC Surveillance Atlas of Infectious Diseases here.

# **Listeria** in the EU, 2021



## **Human cases in foodborne outbreaks**



23 Foodborne outbreaks


8 Strong-evidence outbreaks15 Weak-evidence outbreaks

104 Cases of illness

48 Hospitalisations

12 Deaths

## Foodborne outbreaks





### 3.1. Key facts

- In 2021, 27 MSs reported 2,183 confirmed invasive human cases of *Listeria monocytogenes* infection to ECDC. These cases resulted in 923 hospitalisations and 196 deaths in the EU. Listeriosis was the fifth most commonly reported zoonosis in humans in the EU, and is one of the most serious foodborne diseases under EU surveillance.
- The EU notification rate was 0.49 per 100,000 population, 14.0% higher compared with the rate in 2020 (0.43 per 100,000 population). Reporting in 2020 which showed the lowest number of human cases/lowest rate since the start of listeriosis surveillance in 2007 was impacted by the COVID-19 pandemic and the withdrawal of the United Kingdom from the EU.
- Compared with the rate before the COVID-19 pandemic (2017–2019 annual mean), the 2021 EU notification rate increased by 4.3% and decreased by 2.0% with and without data from the United Kingdom, respectively.
- The overall EU trend for listeriosis in the period 2017–2021 did not show any statistically significant increase or decrease.
- The overall EU case fatality rate was high (13.7%), similar to 2020 and slightly lower than 2019 (13.0% and 17.6%, respectively).
- *L. monocytogenes* infections were most commonly reported in the age group 'over 64 years' and particularly in the age group 'over 84 years'.
- 24 MSs reported a total of 244,357 samples from different 'ready-to-eat' food categories, from the distribution or manufacturing stages.
- The occurrence of *L. monocytogenes* gives an indication of the reasonably foreseeable contamination rate in these 'ready-to-eat' food categories. Results varied according to the 'ready-to-eat' food category, the sampling stage, the number of tested samples and the number of reporting countries. In the framework of objective sampling, all samplers and sampling units included, occurrences remained generally rare (< 0.1%) to low (> 1% to 10%) in these categories. The highest values (from 2% to 5%) were observed for fish and fishery products, meat products from bovines or pigs, fruits and vegetables and cheeses from sheep milk.
- At distribution, the proportions of positive results for single sample enumeration tests for L. monocytogenes carried out by the Competent Authorities (CAs) as part of verification of L. monocytogenes food safety criteria listed in Regulation (EC) No 2073/2005 remained very low to low (< 1.0%) in 9 out of 11 RTE food categories. The highest proportions at distribution were observed in 'ready-to-eat' 'meat products, fermented sausage' (3.1%) and 'fish' (1.5%).</li>
- At manufacturing, the proportions of single samples positive for *L. monocytogenes* based on the detection test were systematically higher compared with those at the distribution level, for all categories of 'ready-to-eat' food. The highest proportions at manufacturing were observed for fishery products (3.1%), products of meat origin other than fermented sausages (2.5%) and fish (1.8%).
- In primary production, the percentage of positive units was very low in cattle (1.1%), which is the most sampled animal species in the EU. The low number of data reported by MSs reflects the absence of minimum legal requirements for harmonised sampling and reporting at primary production.
- With the present report, EFSA has also published two new interactive communication tools on: the EFSA story map (available here) and the dashboard (available here).

## 3.2. Surveillance and monitoring of *Listeria monocytogenes* in the EU

### EFSA story map on Listeria monocytogenes

The EFSA story map on *Listeria monocytogenes* is a new interactive communication tool developed by EFSA in 2022, available online (here) and geared to the general public. This story map provides general information on the pathogen and its epidemiology, including information on where the pathogen can be found, how people and animals get infected, the occurrence of this pathogen in different sources, the disease it causes and how to prevent infection. In addition, this story map also illustrates the monitoring activities implemented in the EU and the role of EFSA with respect to these activities. Users can easily display and explore the content of the different sections in the story map, browsing the dynamic maps, images, text and multimedia features.



#### 3.2.1. Humans

Surveillance of listeriosis in humans in the EU focuses on invasive forms of *L. monocytogenes* infection, mostly manifesting as septicaemia, influenza-like symptoms, meningitis, or spontaneous abortion. Diagnosis of *Listeria* infections in humans is generally carried out by culture, from blood, cerebrospinal fluid and vaginal swabs, or by nucleic acid detection. Since 2018, MSs have been able to submit whole genome sequencing (WGS) data for *L. monocytogenes* to TESSy to be used for EU-wide surveillance and cross-sectoral comparison.

Surveillance of listeriosis is mandatory in all MSs, except in one country (Belgium). The EU case definition was used by 22 countries. Four countries (Denmark, Germany, Hungary and Italy) reported using different case definitions, and one (France) did not specify which case definition was used.

All countries had comprehensive surveillance systems. The surveillance systems for listeriosis cover the whole population in all MSs, except in Belgium, where the estimated coverage of the surveillance system is 80%. This estimated coverage was used in the calculation of notification rates for Belgium. In 2020–2021, Spain did not receive data from all regions that normally report, so the case numbers may therefore not be complete. Bulgaria reported aggregated data, while the other 26 countries reported case-based data.

#### 3.2.2. Food, animals and feed

Monitoring of *L. monocytogenes* is carried out along the food chain, at primary production, manufacturing and distribution.

The public health risk associated with RTE foods depends mainly on the effectiveness of control measures implemented by food business operators (FBOps), including:

- Good Agricultural Practices (GAP) at primary production.
- Good Manufacturing Practices (GMP) and Hazard Analysis and Critical Control Point (HACCP) programmes at manufacturing and distribution.
- Microbiological criteria for RTE foods as defined by Regulation (EC) No 2073/2005.

Official sampling is scheduled by national Competent Authorities (CAs) to verify whether FBOps correctly implement the legal framework of their own-check programmes. Official control samples are thus part of the verification of *L. monocytogenes* food safety criteria (FSC). Data provided to EFSA within this context enable a descriptive summary of contamination levels, especially of RTE foods, at the EU level.

The rationale for surveillance and monitoring of *L. monocytogenes* in animals, feed and food at the different stages along the food chain and the number of samples provided to EFSA are shown in Figure 8.


At the manufacturing and distribution stages, in 2021, 24 MSs reported 301,195 samples tested for *L. monocytogenes* involving different RTE food categories. Due to the impact of the COVID-19 pandemic on food surveillance activities, 2021 results were not comparable with 2020 results. Compared with 2019, the number of reporting MSs was stable (25 MSs in 2019), and the number of tested samples at these stages increased by 11.9% (222,659 samples tested in 2019). In 2020, 24 MSs reported 184,570 sampling units.

At primary production, data are far more comparable with previous years, with 13 MSs reporting 23,564 samples in 2021, compared with 22,567 samples in 2020 (13 MSs) and 22,135 samples in 2019 (13 MSs). Most of the reported data in animals and feed are generated by non-harmonised monitoring schemes across MSs, and for which mandatory reporting requirements do not exist. Data on the occurrence of *L. monocytogenes* in feed are only collected as part of clinical investigations in farm animals. Hence, monitoring data on *L. monocytogenes* in animal feed are rarely available.



18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Isituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Isituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Isituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM). Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Isituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM). Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Isituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM). Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Isituto Zooprofilatico Sperimenta (https

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common



**Figure 8:** Overview of *Listeria monocytogenes* testing along the food chain according to the sampling stage, the sampler and the objective of the sampling, EU

## 3.3. Data analyses

## 3.3.1. Data on RTE food in the context of Regulation (EC) No 2073/2005 on microbiological criteria

These data are based on single samples collected by CAs as part of verification of *L. monocytogenes* FSC listed in Regulation (EC) No 2073/2005, which are to be complied with by FBOps. These FSC are specified by RTE food category and by sampling stage, and are underpinned by the results of either detection or enumeration analytical methods (Table 33). Data were extracted from the database using the values 'official sampling' for the sampler, 'single units' for the sampling unit and 'objective sampling' for the sampling strategy (Table 36).

**Table 33:** Listeria monocytogenes FSC as described in Regulation (EC) No 2073/2005 for the different RTE categories across the food chain, 2021

	RTE foods intended for	Other RTE foods				
Sampling stage	infants and RTE foods for special medical purposes	Able to support growth of <i>Lm</i>	Unable to support growth of <i>Lm</i>			
Manufacturing <sup>(a)</sup>	NA	Based on detection method: Lm not detected in 25 g of sample $(n = 5, c = 0)^{(c)}$	NA			
Distribution <sup>(b)</sup>	Based on detection method: Lm not detected in 25 g of sample (n = 10, c = 0)	Based on enumeration method: limit of 100 CFU/g $(n = 5, c = 0)^{(d)}$	Based on enumeration method: limit of 100 CFU/g (n = 5, c = 0)			

 ${\it Lm: Listeria\ monocytogenes;\ NA:\ not\ applicable;\ RTE:\ ready-to-eat.}$ 

- (a): Before the food has left the immediate control of the food business operator that has produced it.
- (b): Products placed on the market during their shelf-life.



- (c): n = number of units comprising the sample (number of sample units per food batch that are required for testing); c = the maximum allowable number of sample units yielding unsatisfactory test results. In a two-class attributes sampling plan defined by n = 10, c = 0 and a microbiological limit of 'not detected in 25 g', in order for the food batch to be considered acceptable, *L. monocytogenes* must not be detected in qualitative (detection) analyses of 25 g food test portions obtained from each one of 10 sample units taken from the batch. If even one of the sample units from the batch is found to contain *L. monocytogenes* (detected in 25 g), then the entire batch is deemed unacceptable. This criterion applies to products before they have left the immediate control of the producing food business operator, when the operator is not able to demonstrate, to the satisfaction of the CA, that the product will not exceed the limit of 100 CFU/g throughout the shelf-life.
- (d): This criterion applies if the manufacturer is able to demonstrate, to the satisfaction of the CA, that the product will not exceed the limit of 100 CFU/g throughout the shelf-life. The operator may fix intermediate limits during the process that should be low enough to guarantee that the limit of 100 CFU/g is not exceeded at the end of the shelf-life.

Data reported by MSs were classified into the different categories of RTE food/sampling stages based on the assumptions described in the EU Summary zoonoses and foodborne outbreaks report of 2016 (EFSA and ECDC, 2017). Briefly, these assumptions are: all sampling units that were collected from 'cutting plants' and 'processing plants' were considered units collected at the manufacturing stage, while sampling units that were obtained from 'catering', 'hospital or medical care facility', 'retail', 'wholesale', 'restaurant or cafe or pub or bar or hotel or catering service', 'border inspection activities', 'packing centre' and 'automatic distribution system for raw milk' were considered units collected at distribution. When the stage was 'not available' or 'unspecified', data were also considered part of the distribution stage. Other assumptions concerning the RTE categories are presented in the footnotes of Table 36.

#### 3.3.2. Other monitoring data for Listeria monocytogenes in RTE food

The overall occurrence of *L. monocytogenes* in different RTE food categories, regardless of the stage, was assessed from detection results reported by MSs (Appendix B). All levels of sampling unit (single and batches), sampling stages, sampler and sampling contexts (surveillance, monitoring and surveillance – based on Regulation (EC) No 2073/2005) were considered. Only data obtained from the sampling strategies 'objective sampling' and 'census sampling' were used, excluding data reported from 'convenient sampling', 'suspect sampling', 'selective sampling' and 'other' contexts.

Since data were mostly reported by a limited number of MSs and are of a heterogeneous nature, results may not be representative of the EU level or directly comparable across years.

#### 3.3.3. Monitoring data for Listeria monocytogenes in animals and feed

For animals and feed, all sampling strategies were included, even data reported for 'suspect sampling' and 'selective sampling'.

#### EFSA dashboard on Listeria monocytogenes

The EFSA dashboard on *Listeria monocytogenes* (available online here) is a graphical user interface for searching and querying the large amount of data collected each year by EFSA from EU MSs and other reporting countries based on Zoonoses Directive 2003/99/EC. The *Listeria monocytogenes* dashboard shows summary statistics for the monitoring results of the pathogen with regard to major RTE food categories, *Listeria monocytogenes*-positive official samples in the context of FSC in accordance with Regulation (EC) No 2073/2005 and official samples exceeding the FSC limit of 100 CFU/g for specified food matrices. Other monitoring statistics for *Listeria monocytogenes* in RTE foods are also displayed. The *Listeria monocytogenes* data and related statistics can be displayed interactively using charts, graphs and maps in the online EFSA dashboard. In this tool, the main statistics can also be viewed (and downloaded) in tabular format. Detailed information on the use and features of the *Listeria monocytogenes* dashboard can be found in the user guide available on Zenodo here and can also be downloaded from the online tool. Links to the dashboard are available in the relevant sections of this chapter.

18314732, 2022, 12, Downloaded from https://efs.aonlinelibrary.wiley.com/doi/10.2903j.d/sa.2022.72666 by Istituto Cooperinents of the applicable Cerawice Common.

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilattico

Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

#### 3.4. Results

#### 3.4.1. Overview of key statistics, EU, 2017–2021

Table 34 summarises EU-level statistics on human listeriosis and on samples from RTE food tested for *L. monocytogenes* during the period 2017–2021. Food data of interest reported were classified into the major categories and aggregated by year to obtain an annual overview of the volume of data submitted. The sampling efforts of the MSs in 2021 for *L. monocytogenes* in some major RTE food categories can be found In Appendix A. More detailed descriptions of these statistics are provided in the below subsections and in the chapter on foodborne outbreaks.

**Table 34:** Summary of *Listeria monocytogenes* statistics related to invasive human infections and on sampled major RTE food categories in the EU, 2017–2021

	2021 <sup>(a)</sup>	2020	2019 <sup>(b)</sup>	2018 <sup>(b)</sup>	2017 <sup>(b)</sup>	Data source
Humans						
Total number of confirmed cases	2,183	1,887	2,621	2,544	2,474	ECDC
Total number of confirmed cases/100,000 population (notification rates)	0.49	0.43	0.46	0.47	0.47	ECDC
Number of reporting MSs	27	27	28	28	28	ECDC
Infection acquired in the EU	1,482	1,285	1,816	1,640	1,639	ECDC
Infection acquired outside the EU	4	5	14	8	4	ECDC
Unknown travel status or unknown country of infection	697	597	791	896	831	ECDC
Number of outbreak-related cases	104	120	349	159	39	EFSA
Total number of outbreaks	23	16	21	14	10	EFSA
Sampled major RTE food categories <sup>(c)</sup>						
Meat and meat products						
Number of sampling units	107,198	40,291	64,971	58,060	45,322	EFSA
Number of reporting MSs	23	22	22	22	24	EFSA
Fish and fishery products						
Number of sampling units	29,273	11,212	13,366	14,031	12,603	EFSA
Number of reporting MSs	24	23	22	22	24	EFSA
Milk and milk products						
Number of sampling units	66,633	49,132	61,866	59,313	56,428	EFSA
Number of reporting MSs	23	23	23	23	23	EFSA
Products intended for infants or special r	nedical pu	ırposes				
Number of sampling units	2,764	2,394	2,346	2,433	1,943	EFSA
Number of reporting MSs	19	19	19	18	21	EFSA
Other products						
Number of sampling units	94,817	81,541	80,110	28,143	27,946	EFSA
Number of reporting MSs	23	24	24	23	23	EFSA

ECDC: European Centre for Disease Prevention and Control; EFSA: European Food Safety Authority; MSs: Member States; RTE: ready-to-eat.

For a further interactive look at *Listeria monocytogenes* monitoring results, dashboards have been created (*different filters can be applied to query the data*) (*link*).

<sup>(</sup>a): Data on food and animal samples from the United Kingdom (Northern Ireland) are taken into account for 2021. In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/Northern Ireland, EU requirements on data sampling are also applicable to Northern Ireland.

<sup>(</sup>b): Data from the United Kingdom are taken into account for 2017–2019, because the United Kingdom was an EU MS, but it became a third country on 1 February 2020.

<sup>(</sup>c): Number of sampling units tested by detection or enumeration method; number of reporting MSs. More details on the number of samples per MS and for non-MSs can be found in Appendix A.

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilattico

dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

#### 3.4.2. Human listeriosis

In 2021, 27 MSs reported 2,183 confirmed cases of invasive listeriosis in humans (Tables 34 and 35). The EU notification rate was 0.49 cases per 100,000 population, 14.0% higher than 2020 (0.43 per 100,000 population), a year for which the COVID-19 pandemic affected the comparability of data. Comparing the notification rate in 2021 with the rate before the COVID-19 pandemic (2017–2019 annual mean), this rate increased by 4.3% and decreased by 2.0% with and without the data from the United Kingdom, respectively. In 2021, the highest notification rates were observed for Finland, Denmark, Sweden and Slovenia, with 1.3, 1.1, 1.0 and 0.9 cases per 100,000 population, respectively. The lowest notification rates were reported by Bulgaria, Croatia, Cyprus, Greece and Romania ( $\leq$  0.20 per 100,000).

Most (67.9%; 1,482) listeriosis cases with known origin of infection were reported to have been acquired in the EU in 2021 (Table 34). Only four travel-associated listeriosis cases were reported from outside the EU (the United Kingdom, Syria, India and the Dominican Republic) in 2021 vs. five outside the EU in 2020. No data on travel status or country of infection were reported for 697 cases (31.9%).

**Table 35:** Reported confirmed human cases of invasive listeriosis and notification rates per 100,000 population in EU MSs and non-MS countries, by country and year, 2017–2021

	population	in EU MS	s and n	on-MS	count	ries, b	y coun	try an	a year,	2017-	-2021	
	2021				20	20	20:	19	20:	18	2017	
Country	National coverage <sup>(a)</sup>	Data format <sup>(a)</sup>	Confine cases	and	Confine cases	and	Confi cases rat	and	Confine cases	and	Confirmed and ra	
			Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate
Austria	Υ	С	38	0.43	41	0.46	38	0.43	27	0.31	32	0.36
Belgium <sup>(b)</sup>	N	С	65	0.70	54	0.59	66	0.72	74	0.81	73	0.80
Bulgaria	Υ	Α	3	0.04	4	0.06	13	0.19	9	0.13	13	0.18
Croatia	Y	С	8	0.20	5	0.12	6	0.15	4	0.10	8	0.19
Cyprus	Y	С	1	0.11	2	0.23	1	0.11	1	0.12	0	0
Czechia	Υ	С	24	0.22	16	0.15	27	0.25	31	0.29	30	0.28
Denmark	Y	С	62	1.1	43	0.7	61	1.0	49	0.85	57	0.99
Estonia	Υ	С	5	0.38	3	0.23	21	1.6	27	2.0	4	0.30
Finland	Υ	С	70	1.3	94	1.7	50	0.91	80	1.5	89	1.6
France	Υ	С	435	0.64	334	0.50	373	0.56	338	0.50	370	0.55
Germany	Υ	С	560	0.67	546	0.66	571	0.69	678	0.82	721	0.87
Greece	Υ	С	21	0.20	20	0.19	10	0.09	19	0.18	20	0.19
Hungary	Y	С	35	0.36	32	0.33	39	0.40	24	0.25	36	0.37
Ireland	Υ	С	14	0.28	6	0.12	17	0.35	21	0.43	14	0.29
Italy	Y	С	241	0.41	155	0.26	202	0.34	178	0.29	164	0.27
Latvia	Υ	С	10	0.53	8	0.42	6	0.31	15	0.78	3	0.15
Lithuania	Y	С	7	0.25	7	0.25	6	0.21	20	0.71	9	0.32
Luxembourg	Υ	С	4	0.63	4	0.64	3	0.49	5	0.83	5	0.85
Malta	Y	С	0	0	5	0.97	5	1.0	1	0.21	0	0
Netherlands	Υ	С	86	0.49	90	0.52	103	0.60	69	0.40	108	0.63
Poland	Υ	С	120	0.32	57	0.15	121	0.32	128	0.34	116	0.31
Portugal	Υ	С	0	0	47	0.46	56	0.54	64	0.62	42	0.41
Romania	Υ	С	11	0.06	2	0.01	17	0.09	28	0.14	10	0.05
Slovakia	Y	С	13	0.24	7	0.13	18	0.33	17	0.31	12	0.22
Slovenia	Y	С	19	0.90	26	1.2	20	0.96	10	0.48	13	0.63
Spain <sup>(c),(d)</sup>	N	С	224	_	191	_	504	_	370	_	284	_
Sweden	Υ	С	107	1.0	88	0.85	113	1.1	89	0.88	81	0.81
EU Total 27	_	-	2,183	0.49	1,887	0.43	2,467	0.49	2,376	0.50	2,314	0.51
United Kingdom	-	_	_	-	_	-	154	0.23	168	0.25	160	0.24




		2021				2020		2019		18	2017	
Country	National Data coverage <sup>(a)</sup> format <sup>(a)</sup>		Confirmed cases and rates									
			Cases	Rate								
EU Total <sup>(e)</sup>	_	_	2,183	0.49	1,887	0.43	2,621	0.46	2,544	0.47	2,474	0.47
Iceland	Υ	С	5	1.4	4	1.1	4	1.1	2	0.57	6	1.8
Norway	Υ	С	20	0.37	37	0.69	27	0.51	24	0.45	16	0.30
Liechtenstein	Y	С	0	0	58	0.67	36	0.42	53	0.62	45	0.53
Switzerland <sup>(f)</sup>	Υ	С	33	0.38								

<sup>-:</sup> Data not reported.

- (a): Y: yes; N: no; A: aggregated data; C: case-based data.
- (b): Sentinel system; notification rates calculated with an estimated population coverage of 80% for Belgium.
- (c): The notification rate was not calculated since information on estimated coverage was not available.
- (d): Data not complete in 2020–2021, rate not calculated.
- (e): Cases reported by the United Kingdom for the period 2017–2019 were also considered for this estimation (EU-28). When the United Kingdom data were collected for the period 2017–2019, the United Kingdom was an EU MS, but it became a third country on 1 February 2020.
- (f): Switzerland provided data directly to EFSA. The human data for Switzerland include data from Liechtenstein for the years 2017–2020.

The distribution by month appears to be relatively stable. Over the last 5 years (2017–2021), there has been slightly greater reporting of cases in the second half of the year (Figure 9). Although a slight increase in cases was observed in 2021, the overall EU trend for listeriosis cases in the period 2017–2021 did not show any statistically significant increase or decrease. A significantly (p < 0.05) decreasing trend was reported by Romania over the period 2017–2021.



Source: Austria, Belgium, Czechia, Cyprus, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Ireland, Italy, Latvia, Lithuania, Malta, Netherlands, Poland, Romania, Slovakia, Slovenia and Sweden.

Figure 9: Trends in reported confirmed human cases of listeriosis in the EU by month, 2017–2021

18314732, 2022, 12, Downloaded from https://efs.aonlinelibray.wiley.com/doi/10/2033/jfsa.2022/7666 by Istituto Zoproficitieto's Speriment ad ell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023], See the Terms and Conditions (https://onlinelibray.wiley.com/ems-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons.



18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/erms

Information on hospitalisation was provided by 16 MSs for 43.8% of all confirmed cases in 2021, similar to the previous year. Among the cases with information on hospitalisation status, 96.5% were hospitalised totalling 923 hospitalised cases. Listeriosis had the highest proportion of hospitalised cases of all zoonoses under EU surveillance. Outcome was reported for 1,427 confirmed cases (65.4%). Fourteen MSs reported a total of 196 deaths from listeriosis in 2021. This represented a 17.4% increase compared with 2020 (167 deaths). There has been a steady increase in the annual number of deaths between 2010 and 2019 (annual average: 217) which dropped in 2020. However, the overall EU case fatality rate among cases with known outcome was 13.7%, similar to previous years (13.0%) and lower than in 2019 (17.6%).

France reported the highest number of fatal cases (75), followed by Spain (34), Poland (25) and Germany (20). *L. monocytogenes* infections were most commonly reported in the age group over 64. At the EU level, the proportion of listeriosis cases in this age group has steadily increased from 56.1% in 2008 to 64.5% in 2019, and 72.5% in 2020. In the age group over 84, there was an increase from 7.3% to 17.1% in 2019 and 2020, respectively. Within fatal cases of listeriosis, 58.1% of cases were in the age group 64–84 years, while 22.8% were in the age group over 84 years.

### 3.4.3. Listeria monocytogenes in RTE food

In the food sector 24 MSs reported 244,357 samples in different RTE categories at the distribution or manufacturing stages. This sampling effort increased by 80% compared with 2020 (136,346 sampling units) and was comparable with 2019 (218,438 sampling units).

#### Data collected in the context of Regulation (EC) No 2073/2005

In total, 17 MSs reported data in the context of the FSC according to the specifications mentioned above (Section 3.3.1) for 11 RTE food categories (Table 36).

Overall *L. monocytogenes* occurrences reported from official sampling were low, both at distribution and manufacturing (from 0% to 3.1%). Food categories for which data were reported both at manufacturing and distribution systematically showed lower occurrences at distribution. The highest occurrence was found at distribution for 'Product of meat origin, fermented sausages' (3.1%) and at manufacturing for 'Fishery products' (3.1%) and 'Products of meat origin, other than fermented sausages' (2.5%).

For all food categories, the proportion of samples exceeding the limit of the criteria at distribution (100 CFU/g) was very low (0% to 0.66%).

**Table 36:** Proportions (%) of positive single samples at the manufacturing and distribution stage from official sampling by CAs in the context of verification of implementation by FBOps of the *Listeria monocytogenes* FSC in accordance with Regulation (EC) No 2073/2005, EU, 2021

	Manufactur	ing stage <sup>(b)</sup>	Distribution <sup>(c)</sup>							
	Analytical method									
RTE food category <sup>(a)</sup>	<b>Detection</b> (EN ISO 11290-1)	Enumeration (EN ISO 11290-2)	<b>Detection</b> (EN ISO 11290-1)		<b>eration</b> 11290-2)					
	% positive samples (N samples tested, N reporting MS) <sup>(d)</sup>		% positive samples (N samples tested, N reporting MS)	% positive samples (N samples tested, N reporting MS) <sup>(d)</sup>	% of samples exceeding 100 CFU/g <sup>(e)</sup>					
Foods intended for infants and for medical purposes: data reported from BE, BG, CY, ES, HU, IT, RO, SI, SK <sup>22</sup>			0 (N = 1,358; 9 MSs)							

 $<sup>^{22}</sup>$  GR reported enumeration data (N = 6, 0 positive samples).



	Manufactur	ing stage <sup>(b)</sup>		Distribution <sup>(c)</sup>	
			nalytical method	l	
	<b>Detection</b> (EN ISO 11290-1)	Enumeration (EN ISO 11290-2)	<b>Detection</b> (EN ISO 11290-1)	Enume (EN ISO	
RTE food category <sup>(a)</sup>	% positive samples (N samples tested, N reporting MS) <sup>(d)</sup>		% positive samples (N samples tested, N reporting MS) <sup>(d)</sup>	% positive samples (N samples tested, N reporting MS) (d)	% of samples exceeding 100 CFU/g <sup>(e)</sup>
Fish: data reported from BE, BG, CY, CZ, DK, EE, ES, FR, GR, HR, HU, IT, LV, SK	1.8 (N = 501; 8 MSs)			1.5 (N = 1,516; 14 MSs)	0.66
Fishery products: data reported from BE, BG, CY, DK, EE, ES, FR, HR, IT, LV, RO, SI, SK	3.1 (N = 716; 8 MSs)			0.62 (N = 1,285; 11 MSs)	0.08
Cheeses, soft and semi-soft: data reported from BE, BG, CZ, EE, ES, GR, HR, HU, IT, LV, RO, SK	0.95 (N = 4,922; 10 MSs)			0.05 (N = 1,995; 10 MSs)	0
<b>Cheeses, hard:</b> data reported from BG, ES, IT, SK				0 (N = 244; 4 MSs)	0
Cheeses, unspecified: data reported from BE, ES HU, IT, RO, SI	0.77 (N = 1,686; 5 MSs)			0.19 (N = 530; 4 MSs)	0
Other dairy products (excluding cheeses) – entire category: data reported from BE, BG, CZ, EE, ES, GR, HR, HU, IT, LV, RO, SI, SK				0 (N = 1,231; 10 MSs)	0
Milk: data reported from BG, CZ, EE, ES, GR, HR, IT, RO, SK	0.26 (N = 388; 7 MSs)			0 (N = 285; 6 MSs)	0
Products of meat origin, fermented sausages: data reported from BE, DK, ES, HR, HU, SI, SK				3.1 (N = 447; 7 MSs)	0.45
Products of meat origin, other than fermented sausages: data reported from BE, BG, CZ, DK, EE, ES, GR, HR, HU, IT, LV, PL, RO, SI, SK	2.5 (N = 14,136; 12 MSs)			0.19 (N = 4,271; 14 MSs)	0.02
Other products: data reported from BE, BG, CY, CZ, EE, ES, GR, HU, HR, IT, LV, RO, SI, SK	0.72 (N = 1,658; 8 MSs)			0.65 (N = 3,849; 14 MSs)	0.03

1831/372, 2022, 12, Downloaded from https://efsa.onlinelibmry.wiley.com/oi/10.2903j.efsa.2022.7666 by. Istituto Zooprofilatioo Speriment dell'Umbria e del Marche (IZSUM). Wiley Online Library on [1812/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/erms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons. License

MSs: Member States; N: number of single samples tested.

Grey boxes are not applicable in relation to the analytical method for the specific food category and sampling stage in the context of Regulation (EC) No 2073/2005.

- (a): In the absence of relevant physicochemical data (pH, aw), EFSA assumes that foods listed under 'fish and fishery products', 'soft and semi-soft cheeses', 'unspecified cheeses', 'milk', 'products of meat origin other than fermented sausages', 'other dairy products' and 'other products' belong to the category of foods that are able to support the growth of *L. monocytogenes*. EFSA assumes that 'fermented sausages' and 'hard cheeses' belong to the category of foods that are unable to support the growth of *L. monocytogenes*, because foods classified under these two categories of RTE products undergo ripening/fermentation and are expected to have low pH and moderate aw values. For 'other dairy products', EFSA presents the results in a conservative way, by considering all foods included in this category as capable of supporting the growth of *L. monocytogenes*.
- (b): Includes sampling units that were collected from 'cutting plants' and 'processing plants'.
- (c): Includes sampling units that were obtained from 'catering', 'hospital or medical care facility', 'retail', 'wholesale', 'not available', 'unspecified', 'restaurant or cafe or pub or bar or hotel or catering service', 'automatic distribution system for raw milk', 'border inspection' and 'packing centre'.
- (d): Proportion (%) of positive samples (detection of *L. monocytogenes* in 25 g of sample for qualitative analyses, or number of *L. monocytogenes* > or < 100 CFU/g for enumeration analyses); in parentheses the number of tested samples and the number of reporting MSs.
- (e): Proportion of samples exceeding 100 CFU/g for enumeration analyses.

For a further interactive look at *Listeria monocytogenes* monitoring results, dashboards have been created (*different filters can be applied to query the data*) (*link*).

#### Other monitoring data for Listeria monocytogenes in RTE food

Details on the occurrence of *L. monocytogenes* (detection results) in the main RTE food matrices in 2021, together with 2020 and 2019 results, are presented in Appendix B. The below text summarises these results for the major food categories.

#### Fish and fishery products, RTE

In 2021, 22 MSs and four non-MSs (Albania, Iceland, North Macedonia and Serbia) reported such data for fish or fishery products.

In the EU, as for previous years, the occurrence of L. monocytogenes remained one of the highest of all RTE food categories from 3.5% to 5.4%, with an overall mean of 4.7% (N = 9,967). The number of tested samples from fish batches was particularly high for Poland (4,893 tested samples reported in 2021). Excluding Poland, four MSs (France, Italy, the Netherlands and Spain) contributed 67% of the remaining reported data for these categories.

#### Meat and meat products, RTE

In 2021, 20 MSs and three non-MSs (Albania, North Macedonia and Serbia) reported detection data for meat and meat products.

In the EU, the overall occurrence of L. monocytogenes, all matrices included, was 2.3% (N = 40,710 tested units for detection). In all, 68.4% of tested units were assigned to the four main animal species (pigs, bovine animals, broilers and turkeys). The remaining 31.6% of tested samples were reported from other animals, unspecified or mixed meats. RTE meat and meat products from pigs were by far the most frequently tested (60.8%). RTE meat and meat products from bovine animals, broilers and turkeys represented 5.4%, 1.9% and 0.31% of all tested samples, respectively.

#### Pork meat products

Eighteen MSs and three non-MSs (Albania, North Macedonia and Serbia) reported data for pork meat products. In the EU, the overall occurrence of L. monocytogenes remained comparable with previous years: 2.7% in 2021 (N = 24,751), 3.0% in 2020 (N = 6,585) and 4.2% in 2019 (N = 14,035). In 2021, four MSs provided 91% of data on pork meat products (Czechia, Poland, Romania and Spain).

Poultry meat products (broilers and turkeys)

Fourteen MSs and one non-MS (Albania) reported data for RTE poultry meat products (broiler and turkey meat products). The overall occurrence of L. monocytogenes in sampled RTE poultry meat products (N = 895) in EU was 1.3%. This is comparable with 2020 (0.65%, N = 464) and 2019 results (1.9%, N = 931). All positive samples in 2021 (N = 12) were from RTE broiler meat as all 126 samples from RTE meats from turkeys were negative. Six MSs provided 90% of data on poultry meat products (Bulgaria, Czechia, Hungary, Poland, Romania and Spain).

18314732, 2022, 12. Downloaded from https://eis.ao.nlinelibrary.wiley.com/doi/10.2903j.cfs.a.2022.7266 by Isitiuto Zoprofilatios Sperimenta dell'Umbria e delle Marche (ZSUM). Wiley Online Library on 11812023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/rems-and-conditions) on Wiley Online Library for rules of use O.A articles are governed by the applicable Creative Commons

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (thps://onlinelibrary.wiley.com/terms

-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons

#### Bovine meat products

Fourteen MSs and two non-MSs (Albania and Serbia) reported data for RTE bovine meat products. In the EU, the overall occurrence of L monocytogenes reported in RTE bovine meat products was 3.9% (N = 2,217). Four MSs provided 89% of data on bovine meat products (Czechia, Luxembourg, the Netherlands and Poland). For comparison, occurrences were 7.4% in 2020 (N = 856) and 4.2% in 2019 (N = 1,248).

#### Milk and milk products, RTE

In 2021, 20 MSs and four non-MSs (Albania, Montenegro, North Macedonia and Serbia) reported data for RTE milk and milk products, all categories included.

In the EU, the overall occurrence of L. monocytogenes in RTE milk products, all matrices included, was 0.51% (N = 26,154 tested units for detection). This occurrence was 0.69% for cheeses (N = 14,985) and 0.30% for milk (N = 1,642). Seven MSs (Czechia, Hungary, Italy, the Netherlands, Poland, Romania and Spain) provided 81% of data. Additionally, 36.5% of data were assigned to milk or specified cheeses from bovine, goat or sheep origin (Appendix B). The remaining data were reported from other dairy products and cheeses made from mixed milk or unspecified animals.

Among assigned tested samples, 79.1% of cheese samples and 80.5% of milk samples were of bovine origin. Compared with previous years, occurrences observed in all categories of cheeses were rare to very low (< 1%), regardless of the animal sector, except for cheeses from unpasteurised sheep milk where higher occurrences were observed (ranging from 1.7% to 4.6%).

#### Fruits, vegetables and juices, RTE

Fifteen MSs reported data for RTE fruits, vegetables and juices. The overall occurrence was 3.0% (N = 1,407). A total of 80.9% of data were reported by four MSs (Germany, Hungary, Italy and Spain). The results are comparable with previous years. The higher occurrence reported for batches in this category in 2021 compared with previous years may not be representative as a result of a low number of tested samples (N = 24).

## 3.4.4. Listeria spp. in animals

In 2021, 12 MSs and two non-MSs (North Macedonia and Switzerland) reported data on several animal categories (food-producing, wild-, zoo- and pet animals, including birds) from different species (Table 37).

In the EU, reported data were mainly results from 'animals' (98.7%, N = 23,257) compared with other sampling unit levels ('herd/flock' and 'holding'). Considering all sampling unit levels, cattle were the major animal species concerned by *Listeria* testing (70.8% of total units tested, N = 16,687), followed by sheep (15.2%, N = 3,586) and pigs (9.1%, N = 2,137). The sample size, as well as the sampling strategy and the proportion of positive samples, varied considerably among the reporting countries and animal species. Most EU data were reported by 2 MSs: the Netherlands (62.4%) and Ireland (26.0%).

Among the positive units, only limited positive findings were reported as *L. ivanovii* (3.2%) and *L. innocua* (0.80%). As in previous years, many positive findings for *Listeria* (37.8%) were reported as other or unspecified species.

In 2021, a few data (all negative) were reported by Romania in silage (N = 101 units tested) and Greece in feed (N = 65 units tested).

**Table 37:** Summary of *Listeria* species statistics related to major animal species, MSs, 2021

Animal species	N reporting MSs	N tested units	% positive units	N positive units for <i>L.</i> monocytogenes	N positive units for <i>L. ivanovii</i>	N positive units for <i>L. innocua</i>	N positive units for other <i>Listeria</i> species
Cattle	12	16,687	1.1	157	0	0	31
Sheep	10	1,154	5.8	14	2	0	51
Pigs	7	2,137	< 0.01	1	0	0	0
Others	12	3,586	3.3	45	10	3	59
Total EU	12	23,564	1.6	217	12	3	141



For a further interactive look at *Listeria monocytogenes* monitoring results, dashboards have been created (*different filters can be applied to query the data*) (*link*).

#### 3.5. Discussion

In 2021, the number of confirmed cases of human listeriosis was 2,183, corresponding to an EU notification rate of 0.49 per 100,000 population and resulting in an increase of 14% in the notification rate compared with the rate in 2020. Fewer cases were reported in 2020, which could partially be explained by the impact of the COVID-19 pandemic on national healthcare systems and the withdrawal of the United Kingdom from the EU. Listeriosis remains one of the most serious foodborne diseases under EU surveillance due to the high rate of hospitalisations, and high morbidity and mortality, particularly among elderly people.

The general trend for listeriosis in 2017–2021 showed no statistically significant increase or decrease, except for Romania with a statistically significant decrease. Listeriosis had a fairly stable notification rate even during the COVID-19 pandemic, when compared with other foodborne diseases. Clinical manifestations in susceptible categories such as fever, meningitis and septicaemia require hospitalisation or medical treatment and subsequent notification to the national health system (Radoshevich and Cossart, 2018; Quereda et al., 2021). Furthermore, outbreaks, mainly related to the consumption of ready-to-eat foods, meat and meat products, smoked fish and frozen vegetables are not necessarily linked to social and community events (ECDC and EFSA, 2018a,b, 2019). Most infections and outbreaks are domestic.

Listeriosis continues to be one of the foodborne infections with the highest number of fatal cases in the EU, particularly among elderly people. With the ageing of the European population (20.8% of the European population is over 65) (Eurostat, 2022) and the increase in chronic age-related diseases (EFSA BIOHAZ Panel, 2020c), more groups are entering the high-risk categories for severe forms of *Listeria* infections. It is important to continue raising awareness of listeriosis and its risks associated with certain consumption habits and types of food (e.g. RTE fish products and frozen vegetables) (ECDC and EFSA, 2018b; EFSA BIOHAZ Panel, 2018; ECDC and EFSA, 2019).

The molecular characterisation of clinical isolates of *Listeria* is now based on WGS, combined with core genome multi-locus sequence typing (cgMLST) for strain typing. Almost all MSs have moved from pulsed-field gel electrophoresis (PFGE) (Bergis et al., 2021) to WGS (ECDC, 2019), ensuring greater accuracy in typing *L. monocytogenes*. Prior to the use of WGS, a system of rapid and accurate identification of clusters was lacking in sensitivity and, at the European level, many outbreaks were not detected early (Van Walle et al., 2018).

In the food sector, the sampling effort at manufacturing and distribution remained focused on RTE products of animal origin. The occurrence of L. monocytogenes as assessed from detection results reported by MSs (covering objective sampling of single samples and batches) varied according to the food category and sample stage. Occurrences remained generally rare to low in RTE foods. The highest values were observed for fish and fishery products (3.5–5.4%), meat products from bovine or pig origin (2.7–3.9%), fruits and vegetables (2.5%) and hard cheeses from raw or low heat-treated sheep milk (4.6%).

As in previous years, the highest proportions of positive official control samples as part of verification of L. monocytogenes FSC were observed at manufacturing compared with distribution. One exception concerned fermented sausages, for which occurrence at distribution was 3.1%. The results from official sampling showed that the proportion (%) of samples at distribution exceeding the limit of  $100 \, \text{CFU/g}$  was very low (< 1%).

In primary production, cattle were the most frequently sampled animal species and presented a low proportion of positive units (1.1%).

Interpreting national data and trends for the occurrence of *L. monocytogenes* in food or animals over time must be carried out with caution. Currently, surveillance data come from systems that are not fully harmonised and that vary in nature and effectiveness depending on the MS. Moreover, data may be reported by a limited number of MSs depending on each food/animal category.

Combining human, animal and food epidemiological data with molecular and genotyping data provides an efficient methodology to better understand the ecology of this pathogen at different stages of the food chain, and will improve the investigation of listeriosis outbreaks affecting one or more MSs (ECDC et al., 2021). *L. monocytogenes* surveillance in the EU currently uses tools based on genotyping to characterise food and animal contamination and listeriosis outbreaks. ECDC and EFSA have jointly created interoperable databases in order to quickly identify outbreaks. In 2022, the EFSA



18314732, 2022, 12, Downloaded from https://dfs.aonlinelibrary.wiley.com/doi/10.29035j.dfs.2.022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2035]. See the Terms and Conditions (https://onlinelibrary.wiley.com/erms

One Health WGS System that interoperates with the ECDC Molecular Typing system was opened to MSs for the exchange of cgMLST profiles and minimum metadata (EFSA, 2022d). The aim is to collect typing information to detect clusters of foodborne disease and to generate hypotheses on the possible food vehicles involved, supporting risk managers. At the international level, the FAO and WHO (2022) also issued an open call for experts to obtain more representative data on *L. monocytogenes* in foods and to develop a new, full farm-to-table risk assessment.



18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico

Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

## 4. Shiga toxin-producing Escherichia coli (STEC)

Summary data substantiating this chapter, as well as additional information on related projects and internet sources are published for this report on the EFSA Knowledge Junction on Zenodo here. Summary statistics on STEC infections in humans associated with foodborne outbreaks reported in the framework of Directive 2003/99/EC, with downloadable files, are retrievable using the EFSA foodborne outbreaks dashboard available here. Summary statistics on human surveillance data with downloadable files are retrievable using the ECDC Surveillance Atlas of Infectious Diseases here.


## Shiga toxin-producing Escherichia coli (STEC) in the EU, 2021



## Human cases in foodborne outbreaks



## Foodborne outbreaks





### 4.1. Key facts

- In 2021, the number of confirmed cases of human STEC infection was 6,084. This made STEC infection the fourth most commonly reported foodborne gastrointestinal infection in humans in the EU.
- The EU notification rate in 2021 was 2.1 per 100,000 population, representing an increase of 36.9% compared with the rate in 2020 (1.5 per 100,000 population).
- Compared with the rate before the COVID-19 pandemic (2017–2019 annual mean), the 2021 EU notification rate increased by 9.9% and 14.2% with and without the data from the United Kingdom, respectively.
- In 2020, STEC cases reported to ECDC showed the lowest rate since 2007, when the disease surveillance started. The reporting was most likely impacted by the COVID-19 pandemic and withdrawal of the United Kingdom from the EU.
- The overall trend for STEC infections did not show any statistically significant increase or decrease in 2017–2021.
- In 2021, 22 MSs reported the presence of STEC in 3.6% of 17,516 food samples taken according to an 'objective sampling' strategy, compared with 2.4% and 2.8% in 2020 and 2019, respectively.
- 'Sprouted seeds' were tested by 10 MSs in the context of Regulation (EC) No 2073/2005 and one batch was positive for STEC out of 617 sampling units.
- Overall, STEC was most commonly found in 'fresh meat derived from different animal species'
  (7% STEC-positive), followed by bakery products (6.3%) and 'non-ready-to-eat' (RTE) 'milk
  and milk products' (2%), whereas 'fruits and vegetables' was the least contaminated category
  (0.5%).
- Sixteen MSs tested 7,444 RTE food samples for STEC with 112 (1.5%) positives, including 20 (1.6%) from 'meat and meat product samples', 39 (1.7%) from 'milk and milk product samples', 1 (0.3%) from 'spices and herbs' and 10 STEC-positive samples from 'fruits, vegetables and juices' (0.5%).
- Of the food isolates, 27.4% were provided with information on the serogroup and many of these belonged to the top 20 STEC serogroups reported in human infections to ECDC in 2021.
- Most of the virulotypes of STEC isolates from food and animals were also identified in severe STEC infections in humans. Only 32.5% (N = 284) of the STEC isolated from food in 2021 were reported with information on virulence gene typing (stx1 or stx2 and eae) and 9% (N = 79) were provided with the stx gene subtypes.
- Testing of animal samples increased compared with previous years, but still was not widely carried out in the EU. In fact, 3,746 animal samples were reported by seven MSs in 2021 with 6.1% of positives. Most of the sample units (88.5%) were taken from a single animal species, cattle.

## 4.2. Surveillance and monitoring of STEC infections in the EU

## 4.2.1. **Humans**

In 2021, all 27 MSs reported information on STEC infections in humans.

STEC surveillance is mandatory in 24 MSs, voluntary in two MSs (Belgium and France) and one country (Italy) adopted another, non-specified system. The surveillance systems for STEC infections cover the whole population in all EU MSs except for three MSs (France, Italy and Spain). The notification rates were not calculated in these three countries for the following reasons: (a) in France, STEC surveillance in humans is based on paediatric haemolytic-uraemic syndrome (HUS) cases; (b) in Italy, STEC surveillance is sentinel and primarily based on the HUS cases reported through the national registry of HUS; (c) no estimation for population coverage of STEC cases was provided by Spain. In 2020 and 2021, Spain did not receive data from all regions; therefore, the number of reported cases might not be complete. Twenty-six MSs reported case-based data, whereas Bulgaria reported aggregated data.

The EU case definition was used by 26 countries. Four countries reported using a different case definition, and in two instances, the case definition used was not indicated.



#### 4.2.2. Food and animals

Data collected in the context of Regulation (EC) No 2073/2005

The food safety criterion (Regulation (EC) No 2073/2005) specifies that sprouts placed on the market during their shelf-life shall not contain detectable STEC O157, O26, O111, O103, O145 and O104:H4 in 25 g of product. The ISO TS 13136:2012 method with the adaptation developed by the EURL for *E. coli* for detecting O104:H4 or alternative methods validated according to the requirements of the ISO standard 16140 are indicated as reference methodologies for testing.

Nevertheless, the sampling objectives and the sampling frequency vary or are interpreted differently among MSs, resulting in non-harmonised data.

Other STEC monitoring data from food and animals

All the other food and animal testing data originate from the reporting obligations of MSs under Directive 2003/99/EC (i.e. the Zoonoses Directive). Due to the absence of explicitly indicated sampling strategies in this directive, the data generated by MSs are based on investigations with non-harmonised sampling programmes. Therefore, STEC monitoring data in accordance with Directive 2003/99/EC are not comparable between MSs and preclude assessing temporal and spatial trends at the EU level.

In certain food categories, different sampling designs and inaccuracies due to limited numbers of samples also preclude an accurate estimation of prevalence.

Nevertheless, descriptive summaries of sample statistics at the EU level can be compiled and used to indicate the circulation of certain STEC types in food and animals, provided that the above-mentioned relevant limitations of the data set are taken into consideration.

To improve the quality of EU data on STEC monitoring of food and animals, EFSA issued technical specifications for harmonised monitoring and reporting of STEC in animals and foodstuffs in 2009 (EFSA, 2009c). With an additional Scientific Opinion, EFSA encouraged MSs to extend the monitoring and reporting of data on STEC serogroups (EFSA BIOHAZ Panel, 2013c). More recently, it has been recommended that the presence of the main virulence genes be reported, considering the most recent developments in STEC testing and risk assessment (FAO and WHO, 2018; NACMCF, 2019; EFSA BIOHAZ Panel, 2020b). Finally, the latest published EFSA Scientific Opinion on the pathogenicity assessment of STEC presents important considerations related to the virulence of the different STEC types and underlines the significance of determining the virulence gene combinations (virulotypes) of the isolated STEC strains, with an emphasis on *stx* gene subtyping, which would facilitate a more precise assessment of the risk connected with different STEC isolates (EFSA BIOHAZ Panel, 2020b).

### 4.3. Data analyses

### 4.3.1. Occurrence in food and animals

The monitoring data on sprouts as part of Regulation (EC) No 2073/2005 were aggregated and summarised for trend watching according to the following 'filters': Sampling context: 'surveillance, based on Regulation No 2073/2005'; Sampling unit type: 'single'; Sampling stage: as appropriate; Sampling strategy: 'objective sampling' and Sampler: 'official sampling'.

For the description of the occurrence of STEC-positive samples in the different food categories, the subset of the monitoring data with 'objective sampling' specified as sampling strategy was used (N = 17,516), meaning that the reporting MSs collected the samples according to a planned strategy based on the selection of random samples statistically representative of the population to be analysed.

## 4.3.2. Serogroups and virulence features in food and animals

The full data set was used for the descriptive analyses of STEC findings in food (N = 23,659) and animals (N = 3,746), primarily those on virulence gene types and on their frequency distribution.

To analyse the STEC serogroups, the data were grouped according to the test method used, because 3.8% of food samples and 41.5% of animal samples were still assayed using methods detecting only *E. coli* O157. This distinction between methods was necessary to minimise the impact of *E. coli* O157-specific methods on the analysis of the serogroup distribution given that the latter do not identify other STEC possibly present in the samples.

In 2021, the methods targeting any STEC included the ISO TS 13136:2012 method (ISO, 2012), stx gene PCR-based methods and the DIN10118:2004 (DIN, 2004a) standard based on the



18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.cfsa.2022.7666 by Istituto Zooprofilattico

Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

immunochemical detection of Stx by colony blot. The methods designed to detect only *E. coli* O157 were the ISO 16654:2001 method (ISO, 2001), the NMKL 164:2005 method (NMKL, 2005) and the DIN 10167:2004–03 (DIN, 2004b) method. The method indicated as the OIE method for *E. coli* O157 is an adaptation of the ISO 16654:2001 method used for animal samples (WOAH, 2018b).

#### 4.4. Results

#### 4.4.1. Overview of key statistics, EU, 2017–2021

Table 38 summarises EU-level statistics on human STEC infections and on samples from food and animals tested for STEC during 2017–2021. Food and animal data were classified into major categories and aggregated by year to obtain an annual overview of the volume of data submitted. More detailed descriptions of these statistics are provided in the below subsections and in the chapter on foodborne outbreaks.

**Table 38:** Summary of STEC statistics related to humans, major food categories and the main animal species, EU, 2017–2021

	2021	2020	2019 <sup>(a)</sup>	2018 <sup>(a)</sup>	2017 <sup>(a)</sup>	Data source
Humans						
Total number of confirmed cases	6,084	4,489	7,801	8,167	6,071	ECDC
Total number of confirmed cases/100,000 population (notification rates)	2.1	1.5	1.9	2.1	1.7	ECDC
Number of reporting MSs	27	27	28	28	28	ECDC
Infection acquired in the EU	4,355	3,370	4,836	5,783	4,747	ECDC
Infection acquired outside the EU	117	148	751	693	525	ECDC
Unknown travel status or unknown country of infection	1,612	971	2,214	1,691	799	ECDC
Number of foodborne outbreak-related cases	275	208	273	390	260	EFSA
Total number of foodborne outbreaks	31	34	42	50	48	EFSA
Food						
All						
Number of sampling units	23,659	22,119	25,030	20,498	19,351	EFSA
Number of reporting MSs	22	22	22	20	22	EFSA
Meat and meat products						
Number of sampling units	12,160	10,866	14,110	9,250	10,706	EFSA
Number of reporting MSs	19	17	20	17	18	EFSA
Milk and milk products						
Number of sampling units	4,094	4,665	5,479	5,339	3,485	EFSA
Number of reporting MSs	11	10	13	14	10	EFSA
Fruits and vegetables						
Number of sampling units	3,976	3,398	2,696	3,371	2,323	EFSA
Number of reporting MSs	14	15	13	13	15	EFSA
Animals <sup>(b)</sup>						
All						
Number of sampling units	3,746	2,112	2,588	1,631	2,217	EFSA
Number of reporting MSs	7	6	9	5	7	EFSA
Cattle (bovine animals)						
Number of sampling units <sup>(b)</sup>	3,316	868	1,615	1,112	1,681	EFSA
Number of reporting MSs	5	3	7	5	6	EFSA
Small ruminants: sheep, goats, deer						
Number of sampling units <sup>(b)</sup>	151	227	320	188	210	EFSA
Number of reporting MSs	2	2	6	4	2	EFSA

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Islituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

ECDC: European Centre for Disease Prevention and Control; EFSA: European Food Safety Authority; MSs: Member States; STEC: Shiga toxin-producing *Escherichia coli*.

(a): Data from the United Kingdom are taken into account for 2017–2019, because the United Kingdom was an EU MS, but it became a third country on 1 February 2020.

(b): Individual animals.

#### 4.4.2. STEC infections in humans

In 2021, 6,084 confirmed cases of STEC infections were reported in the EU (Table 39). Twenty-four MSs reported at least one confirmed STEC case and three MSs reported zero cases. In 2021, the EU notification rate was 2.1 per 100,000 population. In 2021, there was an increase of 9.9% and 14.2% compared with the average annual notification rate calculated for years 2017–2019 with and without the data from the United Kingdom, respectively. If 2020 data are considered, the increase in the notification rate was higher: 36.9%. However, 2020 data collection was likely affected by the COVID-19 pandemic.

The highest country-specific notification rates among the EU MSs were observed in Ireland, Denmark and Malta (17.54, 15.87 and 13.18 cases per 100,000 population, respectively). Five countries (Romania, Portugal, Greece, Slovakia and Poland) reported  $\leq 0.1$  cases per 100,000 population.

Most STEC cases reported were infected in the EU (71.6% domestic cases or travel in the EU, 1.9% travel outside EU and 26.5% of unknown travel history or unknown country of infection) (Table 38). Overall, for the year 2021, 98.6% of the 4,355 reported STEC cases in humans who acquired the infection in the EU (Table 38) were domestic (acquired within the home country) and 1.4% were acquired through travel in the EU. The proportion of human STEC cases infected within the EU remained stable during 2017–2021, excluding the decrease observed in 2020 during the pandemic.

In 2021, the number of cases infected through travel outside the EU continued to be considerably lower than that reported in 2019, as already observed for the 2020 data. In fact, the proportion decreased from 9.7% in 2019 to 3.1% in 2020 (EFSA and ECDC, 2021) and reached 2.6% in 2021, probably reflecting the disruption in travel caused by the COVID-19 pandemic still in the first half of 2021.

Germany, Sweden, the Netherlands, Austria and Finland reported the highest number of travel-associated cases (107, 97, 33, 18 and 16, respectively), altogether representing 92.2% of all the imported cases (EU and non-EU).

Turkey was most frequently reported as the probable country of infection, followed by Egypt and Kosovo among the non-EU countries (36.8%, 11.1% and 5.1% of the cases with a known probable country of infection, respectively).

**Table 39:** Reported human cases of STEC and notification rates per 100,000 population in EU MSs and non-MS countries, by country and year, 2017–2021

		2021			2020	2020 2019		2018		2017			
Country	National coverage <sup>(a)</sup>	National Data		Confirmed cases and rates									
	_		Cases	Rate									
Austria	Υ	С	383	4.3	288	3.2	284	3.2	305	3.5	250	2.8	
Belgium	Υ	С	124	1.1	84	0.7	131	1.1	112	0.98	123	1.1	
Bulgaria	Υ	Α	0	0	0	0	0	0	0	0	0	0	
Croatia	Υ	С	12	0.30	8	0.20	22	0.54	10	0.24	7	0.17	
Cyprus	Υ	С	0	0	0	0	0	0	0	0	0	0	
Czechia	Υ	С	36	0.34	32	0.30	33	0.31	26	0.25	37	0.35	
Denmark	Υ	С	927	15.9	445	7.6	623	10.7	493	8.5	263	4.6	
Estonia	Υ	С	7	0.53	10	0.75	6	0.45	7	0.53	3	0.23	
Finland	Υ	С	288	5.20	175	3.2	311	5.6	210	3.8	123	2.2	
France <sup>(b)</sup>	N	С	298	-	262	_	335	_	259	_	260	_	
Germany	Υ	С	1,635	2.0	1,409	1.7	1,907	2.3	2,226	2.7	2,065	2.5	




18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (thps://onlinelibrary.wiley.com/terms

		2021					20	19	20	18	2017	
Country	National coverage <sup>(a)</sup>	Data ) format <sup>(a)</sup>	Confirmed cases and rates									
			Cases	Rate								
Greece	Y	С	10	0.09	3	0.03	5	0.05	1	0.01	3	0.03
Hungary	Y	С	24	0.25	8	0.08	23	0.24	14	0.14	12	0.12
Ireland	Υ	С	878	17.5	734	14.8	798	16.3	966	20.0	795	16.6
Italy <sup>(b)</sup>	N	С	65	-	45	_	62	_	73	_	92	_
Latvia	Υ	С	13	0.69	2	0.10	48	2.5	3	0.16	1	0.05
Lithuania	Y	С	0	0	0	0	0	0	0	0	0	0
Luxembourg	Υ	С	10	1.6	0	0	4	0.65	3	0.50	1	0.17
Malta	Y	С	68	13.2	43	8.4	53	10.7	41	8.6	9	2.0
Netherlands	Υ	С	484	2.8	323	1.9	459	2.7	488	2.8	392	2.3
Poland	Υ	С	7	0.02	3	0.01	14	0.04	6	0.02	4	0.01
Portugal	Υ	С	2	0.02	5	0.05	1	0.01	2	0.02	1	0.01
Romania	Υ	С	6	0.03	14	0.07	36	0.19	20	0.10	11	0.06
Slovakia	Υ	С	5	0.09	1	0.02	3	0.06	12	0.22	3	0.06
Slovenia	Υ	С	48	2.3	30	1.4	31	1.5	32	1.5	33	1.6
Spain <sup>(c),(d)</sup>	N	С	101	-	74	_	269	0.57	126	0.27	86	_
Sweden	Υ	С	653	6.3	491	4.7	756	7.4	892	8.8	504	5.0
EU Total 27	_	_	6,084	2.0	4,489	1.5	6,214	1.8	6,327	1.9	5,078	1.7
United Kingdom	_	_	-	_	-	-	1,587	2.4	1,840	2.8	993	1.5
EU Total <sup>(e)</sup>	-	_	6,084	2.1	4,489	1.5	7,801	1.9	8,167	2.1	6,071	1.7
Iceland	Υ	С	7	1.9	4	1.1	27	7.6	3	0.86	3	0.89
Norway	Y	С	437	8.1	331	6.2	511	9.6	494	9.3	381	7.2
Liechtenstein	Y	С	6	15.4	715	8.3	966	11.3	887	10.4	746	8.8
Switzerland <sup>(f)</sup>	Υ	С	915	10.6								

<sup>-:</sup> Data not reported.

- (a): Y: yes; N: no; A: aggregated data; C: case-based data.
- (b): Sentinel surveillance; mainly haemolytic-uraemic syndrome (HUS) cases are notified.
- (c): Notification rate was not calculated because information on estimated coverage was not available.
- (d): Data incomplete for 2021 and 2020, rate not estimated.
- (e): Cases reported by the United Kingdom for the 2017–2019 period were also considered for this estimation (EU-28). When the United Kingdom data were collected for the period 2017–2019, the United Kingdom was an EU MS, but it became a third country on 1 February 2020.
- (f): Switzerland provided the data directly to EFSA. The human data for Switzerland include data from Liechtenstein for years 2017–2020.

The seasonal trend in confirmed STEC cases observed in the EU between 2012 and 2020 was maintained in 2021, with more cases reported during the summer months (Figure 10). The observed STEC infection seasonality is in line with that reported in the literature (Sapountzis et al., 2020). The overall trend for STEC in 2017–2021 did not show any statistically significant increase or decrease, even considering the drop observed in 2020, likely due to the COVID-19 pandemic. At the MS level, a statistically significant increasing trend (p < 0.05) was observed over years 2017–2021 in Denmark and Malta.



Source: Austria, Cyprus, Denmark, Estonia, Finland, France, Germany, Greece, Ireland, Italy, Lithuania, Luxembourg, Malta, Netherlands, Poland, Romania, Slovenia and Sweden.

**Figure 10:** Trends in reported confirmed human cases of STEC infection in the EU by month, 2017–2021

Seventeen MSs provided information on hospitalisation for 35.1% of all confirmed STEC cases in the EU in 2021. Out of the 2,133 cases with known hospitalisation status, 42.2% were hospitalised. The highest proportions of hospitalised cases were reported by Greece, Poland and Slovakia (100% each). However, these MSs only reported a few cases of infection (10, 7 and 5, respectively). Other MSs also reporting high proportions of hospitalised cases were Italy (83.1%), Latvia (76.9%), Estonia (71.4%) and Romania (66.7%). Noteworthily, the high proportion of hospitalised cases observed in Italy is due to the sentinel surveillance system adopted, which is mainly based on HUS notification. A total of 362 HUS cases were reported in 2021 in almost all age groups, with the highest proportion of patients in the youngest age groups from 0–4 years (233 cases; 64.4%) to 5–14 years (71 cases; 19.6%). The most common serogroups among HUS cases were O26 (34% of all cases with serogroup reported), O157 (19.6%), O80 (11%) and O145 (7.6%).

In 2021, 18 fatalities in patients with a STEC infection were reported in the EU, compared with 13 deaths in 2020. Seven MSs reported one to five fatal cases each and 13 MSs reported no fatal cases. This resulted in an EU case fatality of 0.4% among the 4,366 confirmed cases with known outcome (71.8% of all reported confirmed cases). Deaths were mostly reported in the 0–4 years age group (27.8%), followed by the 85+ age group (16.7%).

## 4.4.3. STEC in food

For the year 2021, 22 MSs provided results from analyses of 23,659 food units (batches or single samples).

The most recent source attribution analysis available for STEC underlined that 'bovine meat and products thereof', 'milk and dairy products' and 'vegetables, fruits and products thereof' were the vehicles most frequently implicated in STEC infections in the EU during the 2012–2017 period (inclusive) (EFSA BIOHAZ Panel, 2020b), confirming the results of previous reports (FAO and WHO, 2018). These categories are indeed the most tested in the EU and in 2021 represented 74.5%

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/continelibrary.wiley.c



//efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico

Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

of the food sample units tested, considering the subset of the monitoring data with 'objective sampling' specified as sampling strategy (N = 17,516) (Table 40).

**Table 40:** Occurrence of Shiga toxin-producing *E. coli* (STEC) in the main food categories, EU, 2021 and 2017–2020

		2021		2017–2020 <sup>(a)</sup>			
Food	N reporting MSs	N sampling units	Positive N (%)	N reporting MSs	N sampling units	Positive N (%)	
RTE food							
All	16	7,444	112 (1.5)	19	24,429	226 (0.93)	
Meat and meat products	8	1,276	20 (1.6)	9	5,116	72 (1.4)	
Meat and meat products from bovine animals	5	643	11 (1.7)	7	2,517	45 (1.8)	
Meat and meat products from pigs	3	155	1 (0.65)	6	494	5 (1.0)	
Other meat and meat products	6	478	8 (1.7)	6	2,105	22 (1.0)	
Milk and milk products	8	2,271	39 (1.7)	11	7,507	98 (1.3)	
Milk	2	199	4 (2.0)	5	727	33 (4.5)	
Raw milk <sup>(b)</sup>	1	107	2 (1.9)	4	714	33 (4.6)	
Cheese	8	1,930	32 (1.7)	11	6,363	63 (1.0)	
Dairy products excluding cheeses (butter, cream, ice cream, whey, yoghurt and fermented dairy products)	3	142	3 (2.1)	5	417	2 (0.48)	
Bakery products	2	587	37 (6.3)	3	106	2 (1.9)	
Fruits, vegetables and juices	11	1,922	10 (0.52)	10	6,350	9 (0.14)	
Spices and herbs	3	296	1 (0.34)	7	1,953	8 (0.41)	
Ready-to-eat salads	2	301	0 (0)	4	531	1 (0.19)	
Seeds, sprouted	10	617	1 (0.16)	11	1,730	0	
Non-RTE food						-	
All	17	10,072	518 (5.1)	22	40,921	1,329 (3.2)	
Meat and meat products	16	7,848	503 (6.4)	21	32,172	1,217 (3.8)	
Milk and milk products	6	714	14 (2.0)	9	2,916	103 (3.5)	
Fruits, vegetables and juices	9	1,188	0 (0)	10	2,716	1 (0.04)	
Fresh meat <sup>(c)</sup>							
All	16	6,700	469 (7.0)	21	27,219	1,022 (3.8)	
Fresh meat from bovine animals	16	5,095	288 (5.7)	21	21,017	514 (2.4)	
Fresh meat from pigs	7	604	100 (16.6)	9	1,226	63 (5.1)	
Fresh meat from goats	2	2	0 (0)	3	26	1 (3.8)	
Fresh meat from sheep	4	505	49 (9.7)	6	2,943	309 (10.5)	
Other fresh meat	7	494	32 (6.5)	10	2,007	135 (6.7)	

STEC: Shiga toxin-producing Escherichia coli; N: Number; MSs: Member States; RTE: ready-to-eat.

### Data collected in the context of Regulation (EC) No 2073/2005

As regards 2021 data for STEC on sprouted seeds from the at the distribution level in the context of Regulation (EC) No 2073/2005, 617 samples were tested by 10 MSs with one positive batch (Table 40). This included 388 single official control samples, taken by CAs to verify the compliance of FBOp with the STEC food safety criterion, with none positive. As noted in previous years, testing sprouted seeds is still not widely applied at the EU level, although a microbiological criterion for this food commodity has been laid down in Regulation (EC) No 2073/2005 since 2013.

<sup>(</sup>a): Data from the United Kingdom are taken into account for 2017–2019, because the United Kingdom was an EU MS, but it became a third country on 1 February 2020.

<sup>(</sup>b): Raw RTE milk sampling units are a subset of RTE milk.

<sup>(</sup>c): Fresh meat sampling units are a subset of the two main categories (RTE food and non-RTE food).



#### Other STEC monitoring data from food

Overall, 3.6% of the 17,516 food sample units tested by 21 MSs, and collected using an objective sampling strategy, were positive for STEC. For years 2020, 2019, 2018 and 2017, the reported percentages of STEC-positive food samples were lower, namely 2.4%, 2.8%, 2.8% and 1.5%, respectively (EFSA and ECDC, 2021). In Table 40, these monitoring results are summarised, and a distinction is made between RTE food, non-RTE food and 'fresh meat'.

#### RTE food

As regards RTE food, testing of 7,444 samples was reported. The food category with the highest number of sample units tested was 'milk and milk products' with 2,271 samples assayed, notably cheeses (25.9%), followed by 'fruits, vegetables and juices' (25.8%), 'meat and meat products' (17.1%), 'seeds, sprouted' (8.3%) and bakery products (6.3%). In total, 112 RTE food samples were found to be positive for STEC: 1.6% in 'meat and meat products' (notably of bovine origin), 1.7% in 'milk and milk products' particularly milk, 0.5% in the 'fruits, vegetables and juices' category and 0.3% in 'spices and herbs' (Table 40).

#### RTE and non-RTE food

To provide overall statistics, the following descriptive analyses are based on merged data from RTE food and non-RTE food.

Meat and meat products

Bovine meat

In 2021, 5,095 units of fresh and unprocessed bovine meat were tested. At the distribution stage, 2,815 units were sampled by 12 MSs with 4.1% of positives. Samples (N = 2,280) taken at the manufacturing level were the most contaminated (7.5%, 9 MSs), particularly at the slaughterhouse with 9.1% of the samples testing positive for STEC.

Sheep and goat meat

Small ruminants are an important reservoir of STEC as reported in the literature (Persad and LeJeune, 2014). In 2021, four MSs reported the results of an investigation on 505 sample units of fresh sheep meat with 9.7% of these being STEC-positive, whereas two MSs reported on fresh goat meat with no STEC-positive sampling units out of the two tested (Table 40). This proportion of contaminated samples is consistent with that from the analysis of 2017–2020 data (10.5%). However, this food commodity is generally tested in surveys by very few MSs.

The sampling stage that yielded the highest proportion of positive sheep meat samples was the distribution level, with 14.3% of samples producing a STEC isolate. Conversely, the samples taken at the slaughterhouse showed a much lower STEC contamination rate (4.6%).

Meat from other ruminants

Only three MSs provided information on the presence of STEC in fresh deer meat samples. In total, 101 samples were taken and 17 were found to be contaminated with STEC (16.8%). The sampling stage with highest percentage of samples contaminated by STEC was the manufacturing level (29.4%). Deer hunting may have contributed to this result because the use of firearms and the slaughtering procedures may favour gut perforation and the contamination of meat, respectively.

Meat from other animal species

Seven MSs tested fresh pig meat in 2021 and reported data on 604 samples, with 100 of these being positive for STEC (16.6%) (Table 40).

Fresh meat from animal species other than bovine, ovine, goat, pig and deer species was tested in 2021 by seven MSs that reported on the analyses carried out on 494 sample units (Table 40). These included samples taken from poultry, ducks, wild and farmed game, geese, horses, rabbits, wild boars and unspecified meat. Thirty-two samples were reported as STEC-positive (6.5%) (Table 40).

Meat products and meat preparations

Meat products and meat preparations other than fresh meat were sampled in 2021 by 11 MSs that tested 2,485 samples, resulting in 58 isolated STEC strains (2.3%).

18314732, 2022, 12, Downloaded from https://fs.aonlinelibrary.wiley.com/inchibrary.wiley.com/inchibrary.wiley.com/em-and-conditions) on Wiley Online Library on [18/12/2023], See the Terms and Conditions (http://oinineibrary.wiley.com/em-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons



18314732, 2022, 12, Downloaded from https://disa.onlinelibrary.wiley.com/doi/10.2903/j.disa.2022.7666 by Isituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.2903/j.disa.2022.7666 by Isituto Zooprofilation Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.2903/j.disa.2022.7666 by Isituto Zooprofilation Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.2903/j.disa.2022.7666 by Isituto Zooprofilation Sperimenta dell'Umbria e delle Marche (IZSUM). Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.2903/j.disa.2022.7666 by Isituto Zooprofilation Sperimenta dell'Umbria e dell'Um

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

#### Milk and milk products

Overall, STEC was found in 53 (1.8%) out of 2,985 samples of RTE and non-RTE milk and milk products including cheese reported by eight MSs (Table 40).

In 2021, seven MSs reported on the testing of 503 sample units of raw cows' milk with seven positive units (1.4%). One MS reported monitoring results on 13 sample units of raw goats' milk, and two MS reported only 8 samples of raw sheep milk. Both categories recorded one positive sample each.

The presence of STEC in RTE dairy products other than milk and cheeses was reported by three MSs, which tested 142 sample units of butter, cream, ice cream, yoghurt and fermented dairy products. Three positive samples were detected (Table 40).

For the cheese samples, 1,930 samples were tested for the presence of STEC, with 32 (1.7%) positive units from eight MSs in 2021.

### Vegetables and fruits

STEC were found in 10 (0.3%) out of 3,110 samples of fruits and vegetables (Table 40). In total, 1,214 sample units of RTE spice and herbs, salads and sprouted seeds were tested and 2 STEC-positive units were reported (0.2%) (Table 40).

#### Other foodstuffs

This category contains miscellaneous food commodities not comprised in the previously mentioned categories, and included cereals and meals, bakery products, fish and fishery products, infant formula and others. For the whole category, 865 samples were analysed by five MSs with 39 (4.5%) positive samples reported from the cereals and meals (2) and bakery products (37) food categories.

#### 4.4.4. STEC in animals

For the year 2021, results from 3,746 sampling units (single heads or herds or flocks) from animals were reported by seven MSs, (Table 41), and no data from non-MSs have been reported. This number has increased compared with the number of animals tested during 2017–2020.

The most tested animal category in 2021 was cattle, with 3,316 sample units tested (88.5% of all animal samples tested) by five MSs and 6.1% of positives (Table 41). The number of samples tested for this animal category showed an approximately four-fold increase with respect to the previous year and a two-fold increase considering the 2017–2019 period (EFSA and ECDC, 2021). The proportion of positive samples varied significantly with sampling stage. The 322 sample units taken at the farm level yielded 17.4% of positives, and the testing of the 2,994 samples taken at the slaughterhouse returned 139 STEC positive samples (4.6%).

The other animal species tested included pigs, Cantabrian chamois, deer and sheep among others, with only few samples assayed per category by three MSs.

**Table 41:** Summary of STEC statistics related to major animal species, reporting EU MSs, 2021

Animals	N reporting MSs	N tested sampling units	Positive sampling units		
			N	%	
Cattle 5		3,316	195	5.9	
Sheep and goats	1	48	7	14.6	
Other ruminants <sup>(a)</sup>	2	121	1	0.83	
Pigs	1	51	6	11.8	
Other animals/not specified	specified 2 210		20	9.6	
Total	7	3,746	229	6.1	

MSs: Member States; STEC: Shiga toxin-producing Escherichia coli.

(a): Other ruminants include Cantabrian chamois, deer and water buffalos.



#### 4.4.5. Focus on STEC strain features: virulence genes and serogroups

#### **Humans**

Data on STEC serogroups were reported in 2021 by 22 MSs for 56.5% of the confirmed human cases. The most frequently reported serogroups were O157 (15.2% of the human cases reported with information on the serogroup) and O26 (14.8%). These two serogroups together represented 30% of the total number of confirmed human cases with known serogroups in 2021. The proportion of non-typable STEC isolates rose to 25.9% in 2021 (18.3% in 2020).

Data on virulotypes (based on Shiga toxin genes stx1, stx2 and the intimin-coding gene eae) were reported for 35% (N = 2,130) of confirmed STEC infections (N = 6,084) in 2021. The most frequently reported virulence gene combination in strains isolated from severe cases (HUS, bloody diarrhoea and hospitalised cases) was stx2+/eae+, accounting for 34.1% of the strains with a known virulotype (Table 42), half of which were from HUS cases (84 strains). The proportion of the second most common virulotype stx1+/stx2+/eae+ accounted for 26.2% of these cases (Table 42). The most common stx gene subtypes in strains from severe human infections were stx2a (38.3% of isolates with reported stx gene subtyping data), stx1a (27.7%), stx2c (8.8%) and stx2d (9.0%) (Table 43).

**Table 42:** Virulotypes (*stx* type and presence of *eae*) in food, animal and human STEC isolates causing severe infection (haemolytic-uraemic syndrome (HUS), hospitalisation and bloody diarrhoea) in 2021 and comparison with those associated with severe disease in humans during 2012–2017, EU

Virulence	N of animal	N of food	N of human	Relative frequency of the virulotype in <sup>(b)</sup>				
genes profile	isolates in 2021 <sup>(a)</sup>	isolates in 2021 <sup>(a)</sup>	isolates in 2021 (%)	HUS	Hospitalisation	Bloody diarrhoea		
stx2; eae+	53	23	164 (34.1)	17.7	42.0	40.2		
stx1; stx2; eae+	17	10	126 (26.2)	5.9	35.7	64.8		
stx1; eae+	43	34	58 (12.1)	1.2	27.4	27.3		
stx2; eae-	_	139	74 (15.4)	2.7	24.3	14.8		
stx1; eae-	_	61	34 (7.1)	0.3	20.3	14.1		
stx1; stx2, eae-	-	17	25 (5.2)	1.4	15.3	19.4		
Total	77	284	481 (100)	_	_	_		

STEC: Shiga toxin-producing *Escherichia coli*; HUS: haemolytic-uraemic syndrome. The *stx* genes were characterised at the type level (*stx1* and *stx2*).

**Table 43:** Virulotypes (*stx* subtype) of food and human STEC isolates causing severe infection (haemolytic-uraemic syndrome, hospitalisation and bloody diarrhoea) in 2021 and comparison with those associated with severe disease in humans during 2012–2017, EU

stx genes subtypes combinations	N of food	N of human	Relative frequency of the stx gene subtype combinations in (b)							
	isolates in 2021 <sup>(a)</sup>	isolates in 2021 (%)	HUS		Hospitalisation		Bloody diarrhoea			
			eae+	eae-	eae+	eae-	eae+	eae-		
stx2a	20	184 (38.3)	27.4	10.4	56.4	32	58.4	26.3		
stx1a	15	133 (27.7)	1.2	0	27.6	20.7	27.3	8		
stx2d	7	43 (9.0)	NR	10.3	NR	33.3	NR	16		
stx2c	7	42 (8.8)	4.3	5	19.8	NR	23.9	NR		
stx2b	1	29 (6.0)	NR	0.5	NR	21.3	NR	10.5		

18314732, 2022, 12, Downloaded from https://sta.onlinelibrary.wiley.com/doi/10.2903/j.dsa.2022.72666 by Isitiuto Zooprofilatios Sperimental dell'Umbria e delle Marche (ZSUM), Wiley Online Library on [18122023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/em-and-conditions) on Wiley Online Library for rules of use; O.A articles are governed by the applicable Creative Common

<sup>(</sup>a): Due to the low number of isolates virulotyped for food and animals, only the number of isolates is shown.

<sup>(</sup>b): Relative frequencies (%) of the different combinations of *stx* gene types with or without the *eae* gene in STEC isolated from severe disease (TESSy data, 2012–2017) (EFSA BIOHAZ Panel, 2020b).



18314732, 2022, 12, Downloaded from https://drs.aonlinelibrary.wiley.com/doi/10.2003j.efsa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/erms

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

sty gangs	N of food	N of human isolates in 2021 (%)	Relative frequency of the <i>stx</i> gene subtype combinations in <sup>(b)</sup>							
stx genes subtypes combinations	isolates in 2021 <sup>(a)</sup>		HUS		Hospitalisation		Bloody diarrhoea			
			eae+	eae-	eae+	eae-	eae+	eae-		
stx1c	2	18 (3.8)	NR	0.6	NR	18.9	NR	19.5		
stx2f	ND	18 (3.8)	3.8	NR	21	NR	8.7	NR		
stx2c;stx2a	ND	5 (1.0)	29	NR	57.1	NR	65.5	NR		
stx2g	3	3 (0.6)	NR	_	_	NR	NR	NR		
stx2c;stx2d	ND	3 (0.6)	_	_	_	_	_	_		
stx2e	8	1 (0.2)	_	NR	NR	NR	NR	31.8		
stx2b;stx2c	ND	1 (0.2)	_	_	_	_	_	_		
stx2d;stx1a	5		_	_	_	_	_	_		
stx1d	2		_	_	_	_	_	_		
stx2a;stx1a	7		20.8	4.5	59.3	NR	56.6	NR		
stx2d;stx2c;stx2a	1		_	_	_	_	_	_		
stx2c;stx1a	1		_	_	_	_	_			
Total	79	480	_	_	_	_	_	_		

Note: Two STEC isolates with subtypes stx2a and stx2c/stx2a, respectively, were reported in animals in 2021.

STEC: Shiga toxin-producing *Escherichia coli*; HUS: haemolytic-uraemic syndrome. The *stx* genes were characterised at the subtype level.

NR: Data present in the TESSy data set used with fewer than 20 isolates.

ND: Not detected.

- -: not present in the TESSy database during the 2012-2017 period.
- (a): Due to the low number of isolates virulotyped for food, only the number of isolates is given.
- (b): Relative frequencies (%) of the different combinations of *stx* gene subtypes in STEC isolated from severe disease (TESSy data. 2012–2017). Human data from: EFSA Journal 2020;18(1):5967.

#### **Food**

Most of the 20 most common STEC serogroups isolated from human infections were also found in the STEC isolated from food in 2021, except for serogroups O55, O63, O80, O125 and O177. The information on the serogroup was available for 239 (27.4%) out of 872 strains isolated from the overall 23,659 sample units tested in 2021.

Information on *stx1* and/or *stx2* and *eae* genes was provided for 284 (32.6%) of the STEC strains (Table 42). Only 79 STEC isolates were subtyped for the *stx* gene (Table 43). This was 27.8% of the isolates with *stx* types and *eae* genes reported, and was 9.1% of the total number of STEC isolated from food in 2021. Tables 42 and 43 show the combinations of the virulence genes determined in the food, animal and human STEC isolates in 2021 and the numbers and frequencies of these combinations in the STEC isolated from severe human disease in the EU in 2012–2017, as analysed in the latest pathogenicity assessment of STEC (EFSA BIOHAZ Panel, 2020b). Given the scarce amount of data on the virulence genes characterised in food and animal isolates in 2021, the number of isolates are displayed instead of the relative frequency for each virulotype.

#### **Animals**

This section includes the analysis of the data related to the 3,746 sampling units from animals tested, of which 6.1% (N = 229) were positive for the presence of STEC.

For the analysis of the distribution of STEC serogroups and virulotypes, all STEC strains isolated from the animal sampling units were considered and from some units multiple isolates were obtained. In total there were 245 STEC strains and of 201 strains (82%) were provided with information on the serogroups. However, 36 isolates were obtained using the analytical methods detecting the O157 serogroup only. The isolates with serogroup information belonged to 22 different O-groups, with 8 of these represented in the top 20 serogroups isolated from human disease in 2021.

For the analysis of the virulence genes of STEC strains, 113 STEC animal isolates (46.1%) were provided with the virulotype based on the identification of the stx1, stx2 and eae genes (Table 42).



Two MSs also carried out *stx* gene subtyping and reported this information for only two STEC strains (*stx2a*/*stx2c*; *stx2a*).

All data provided by the reporting countries were used to generate atlases of the STEC serogroups identified in the different food and animal categories compared with those determined in human isolates for years 2017–2021 (Appendix C) and for 2021 for food and animal isolates only (Appendices D and E).

#### 4.5. Discussion

In 2020, there was a decline in the notification rates of STEC infections, probably due to the COVID-19 pandemic. In 2021, the number of reported cases was comparable to those reported during the pre-pandemic period (2017–2019).

In 2021, more than half of the confirmed human cases reported by EU MSs had information on the serogroup with the most frequently reported being O157, followed by O26, although the two proportions diverged by only 0.4%.

It has been proposed that the highest predictive power in terms of pathogenicity potential of STEC strains resides in the characterisation of the Shiga toxin-coding genes (*stx*) and, to a lesser extent, the intimin-coding *eae* gene (FAO and WHO, 2018; NACMCF, 2019; EFSA BIOHAZ Panel, 2020b). Moreover, *stx* gene subtyping helps to identify the strains that have a higher frequency of association with severe disease (EFSA BIOHAZ Panel, 2020b). Regarding subtyping capacity, more than half of the MSs' national public health laboratories reported the ability to perform whole-genome sequencing for STEC isolates (EFSA BIOHAZ Panel, 2020b). This is a promising perspective, possibly fostering increased reporting of typing and subtyping data for STEC isolates in the coming years.

In 2021, more than one third of the STEC cases were reported with information on the stx (stx1 or stx2) and the eae genes and only 22.5% of the isolates with a reported stx1, stx2 and eae virulotype were provided with information on stx gene subtypes. Based on the analysis of the stx subtypes reported in TESSy from 2012 to 2017 (EFSA BIOHAZ Panel, 2020b), all STEC virulence gene combinations and most of the stx gene subtypes identified in 2021 can be associated with severe illness, albeit at different frequencies.

More than 40% of the cases with information on hospitalisation status were hospitalised. The highest proportions of hospitalised cases were reported by Greece, Poland and Slovakia, countries that accounted for 100% of the hospitalised cases; however, these countries reported low numbers of cases, below or equal to 10. A total of 362 HUS cases were reported in 2021 in almost all age groups, with the highest proportion of patients in the youngest age groups, from 0 to 4 years. Most cases of deaths were reported in this age group followed by the 85+ age class.

In 2021, 22 EU MSs reported monitoring results of STEC in 23,659 food samples. Not all reporting MSs have tested all food categories equally. After aggregating the food samples into macro-categories in 2021, the number of MSs testing and reporting data on the presence of STEC in food ranged from 19 MSs reporting the testing for STEC in meat and vegetables (including seeds) to 11 MSs testing milk and dairy products. Sprouted seeds were tested by 10 MSs, considering the samples taken in the context of Regulation (EC) No 2073/2005. As noted in previous years, although the microbiological criterion for the presence of STEC in seeds and sprouts has a normative value, the sampling of this food categories in the EU appears to be infrequent.

The analytical procedures for testing food in the EU have been substantially harmonised. In 2021, all the reporting countries used the ISO TS 13136:2012 or equivalent method to test 22,760 samples (96.2%) out of the 23,659 total samples tested. The remaining data reported by some MSs (four) for specific surveys were obtained with the ISO 16654:2001 or equivalent method. This method detects serogroup O157 only and does not give information on any other STEC serogroups possibly present in the sample.

The general extent of observed STEC contamination in food, assessed using food samples taken according to an 'objective sampling' strategy, was 3.6%. As observed in previous years, the frequencies of STEC contamination varied among the different major food categories, RTE and non-RTE. The most contaminated food categories included commodities of animal origin, whereas vegetables and fruits were the least contaminated or did not yield positive results at all. The finding of 6.3% of bakery products samples contaminated with STEC is interesting. However, most of the contaminated samples were reported by one single MS, which tested the majority of the samples. Such a high level of contamination might be related to the reported contamination of the flour, which has been identified as the cause of some outbreaks of STEC infections in the US and Canada

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (thps://onlinelibrary.wiley.com/terms

(Gill et al., 2019). Additionally, behaviours such as tasting and manipulating raw dough should also be considered in relation to the risk of STEC infections from contaminated flour (Crowe et al., 2017).

STEC-positive units were detected in all types of RTE food. Importantly, only a few MSs reported data for certain food categories or with a limited sampling effort for certain foods (e.g. one MS reported 107 raw milk sample results). The testing of RTE food commodities for STEC is important because these foods are consumed without any treatment to reduce or eliminate the possible presence of the pathogen, posing a direct risk to the consumer.

In 2021, 27.4% of the food isolates were provided with information on the serogroup, compared with 28.2% and 34.4% observed in 2020 and 2019, respectively, and 41.8% in 2018 (EFSA and ECDC, 2021). Serogroup determination clearly shows a decreasing trend, which probably reflects the awareness that this character is of limited importance in assessing the virulence of STEC strains. Most of the top 20 STEC serogroups isolated from human infections in 2021 were also found in the STEC isolated from food in the same year.

As regards the animal monitoring results for 2021, overall, 6.1% of the 3,746 samples taken were STEC-positive. However, in 2021, the samples tested from cattle represented 88.5% of the total animal sample units assayed by five MSs and 5.9% of these were contaminated with STEC. In any case, the number of animal sampling units tested, although considerably increased with respect to 2020, continued to be very low, possibly biasing the estimates.

The analysis of the presence and subtypes of virulence genes is important for pathogenicity assessment (EFSA BIOHAZ Panel, 2020b). Unfortunately, this level of characterisation is still far from being routinely carried out for food and animal isolates, and only 32.6% of the STEC isolated from food in 2021 were reported together with the information on the stx gene types (stx1 or stx2) and the intimin-coding gene eae. The number of strains with information on stx gene subtyping was even lower (N = 79), representing 27.8% of the isolates with stx types and eae genes reported, and 9.1% of the total number of STEC isolated from food in 2021. Given that this characterisation strategy represents the basis for a molecular risk assessment of STEC circulating in the vehicles of infection, MSs should be encouraged to continue to adopt this approach.

The virulotyping of STEC isolated from food in 2021, although conducted on a small subset of isolates, confirmed that all the virulotypes identified, based on the *stx* types and the *eae* gene, matched those associated with the STEC strains isolated from severe disease (EFSA BIOHAZ Panel, 2020b) (Table 42). As far as the *stx* gene subtyping is concerned, 9 out of the 13 combinations identified in food isolates were also represented among those associated with severe disease (Table 43).

On the other hand, 46.1% of animal isolates (N = 113) were reported with data on the characterisation of the virulence genes, but only two animal isolates had undergone stx gene subtyping by two MSs. Also in this case, all the stx1/stx2/eae profiles and the two subtype combinations identified – both eae positive – were represented in STEC isolated from human severe disease during the 2012–2017 period (EFSA BIOHAZ Panel, 2020b) (Tables 42 and 43).

Given the predictive power of the *stx* gene subtyping, a wider adoption of this approach is recommended. Raising awareness on the need to enhance the characterisation of STEC strains will lead to an improved risk assessment of STEC, ultimately supporting actions to mitigate the impact of STEC on public health.



e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

## 5. Tuberculosis due to Mycobacterium bovis or Mycobacterium caprae

Summary data substantiating this chapter, as well as additional information on related projects and internet sources are published for this report on the EFSA Knowledge Junction on Zenodo here. Summary statistics on human surveillance data with downloadable files are retrievable using the ECDC Surveillance Atlas of Infectious Diseases here.

## Tuberculosis in the EU, 2021

due to Mycobacterium bovis or Mycobacterium caprae

Human cases	Notification rate (per 100,000 population)	0.03	irena —	creasing ecreasing able	
111 Cases of illness					
55 Cases in individuals of EU origin		NA	Hospitalisations *		
47 Cases in individuals originating from outside the EU		10	Deaths *		
9 Unknown infections			* Data refers to 2020		

## **Human cases in foodborne outbreaks**



## 5.1. Key facts

- In 2021, the number of confirmed cases of human tuberculosis due to *Mycobacterium bovis* or *Mycobacterium caprae* was 111, corresponding to an EU notification rate of 0.03 cases per 100,000 population. This is an increase of 12.4% compared with the rate in 2020 (0.026 per 100,000 population).
- Compared with the EU notification rate before the COVID-19 pandemic (2017–2019 annual mean), there was a decrease of 28.3% and 25.3% with and without the data from the United Kingdom, respectively.
- In both 2021 and 2020, the *M. bovis* or *M. caprae* reporting recorded the lowest number of human cases and rates since 2017, because it was impacted by the COVID-19 pandemic and withdrawal of the United Kingdom from the EU.
- In 2021, the *M. bovis* and *M. caprae* case notification rate was similar (0.03 cases per 100,000) among EU MSs with disease-free status and EU MSs with non-disease-free status in the bovine population.
- Almost half (49.5%) of *M. bovis* and *M. caprae* cases in humans were of EU origin (native cases and/or cases originating from other MSs).
- No foodborne outbreaks due to *M. bovis* or *M. caprae* have ever been reported to EFSA since the start of the data collection on foodborne outbreaks in 2004; 2021 was no exception.
- In 2021, the overall prevalence of bovine tuberculosis due to *M. bovis* or *M. caprae* increased slightly (0.6%) compared with the previous year, and the number of infected cattle herds in the EU increased from 7,372 to 9,690 herds. This increase was mainly due to the United Kingdom (Northern Ireland) data, which were not included in the EU 2020 statistics. Infection of cattle with *M. tuberculosis* was not reported to EFSA.

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/erms

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

- Fourteen MSs and the United Kingdom (Northern Ireland) detected the presence of bovine tuberculosis in 2021. Similar to previous years, the distribution of infected herds was heterogeneous and spatially clustered, with a national herd-level prevalence ranging from < 0.01% (Poland) to 11.3% (the United Kingdom (Northern Ireland)).
- Seventeen MSs had disease-free status during 2021. Ten MSs and the United Kingdom (Northern Ireland) were non-disease-free, of which three MSs had disease-free zones.
- Overall, 139 cattle herds proved infected with *M. bovis* or *M. caprae* in the disease-free zones (0.014%), confirming that infection occurs rarely in these zones.
- In the non-disease-free zones of 10 MSs and the United Kingdom (Northern Ireland) 9,551 cattle herds (1.3% of total herds) showed infection with *M. bovis* or *M. caprae* in 2021. The United Kingdom (Northern Ireland) (11.3%), Ireland (4.6%) and Spain (1.3%) were the only countries that reported prevalence higher than 1%. No infected herds were reported by Malta or Cyprus.
- In the last 10 years (2012–2021), the annual number of infected cattle herds and the prevalence of bovine tuberculosis in non-disease-free zones has decreased by 47.5% and 4.2%, respectively. This decrease is mainly attributable to the withdrawal of the United Kingdom from the EU in 2020. In fact, the annual prevalence of herds infected with *M. bovis* or *M. caprae* in all the zones of the United Kingdom was higher than 10% between 2012 and 2019. For the same reason, the inclusion of the United Kingdom (Northern Ireland) among the EU zones explains the increase in both the number of infected herds and the prevalence recorded in 2021.

# 5.2. Surveillance and monitoring of tuberculosis due to *Mycobacterium bovis* or *Mycobacterium caprae* in the EU

#### **5.2.1.** Humans

The notification of tuberculosis in humans is mandatory in all 27 EU MSs and covers the whole population. All countries report case-based data. The proportion of tuberculosis cases caused by *M. bovis* and *M. caprae* was calculated using the preliminary estimate of the total number of confirmed tuberculosis cases in 2021 among reporting MSs' species-specific data. No human data on *M. bovis* and *M. caprae* cases are available for France because this MS does not report species-specific data within the *Mycobacterium tuberculosis* complex (MTBC) for human tuberculosis cases. In addition, Latvia did not report any MTBC data during 2018–2020 and Iceland in 2021.

Given that tuberculosis is a chronic disease with a long incubation period, it is not possible to assess travel-associated cases in the same way as for other, acute-onset zoonoses detailed in this report. Instead, a distinction is made between individuals with the disease originating from an EU MS (cases of EU origin) and those originating from outside the EU (case originating outside of the EU). In the analysis, origin is mainly based on the reported birthplace, except for cases from Austria, Belgium, Greece, Hungary and Poland, whose origin is based on their reported citizenship. The treatment outcome for tuberculosis due to *M. bovis* and *M. caprae* is assessed 1 year (12 months) after case notification, because the shortest duration for treatment completion is 6 months, according to the international treatment guidelines for tuberculosis.

#### **5.2.2.** Animals

Bovine tuberculosis surveillance data

On 21 April 2021, Commission Implementing Regulation (CIR) (EU) 2020/2002<sup>23</sup> under the Animal Health Law (AHL) entered into force, with Union notification and reporting covering infection with *Mycobacterium tuberculosis* complex (*M. bovis, M. caprae* and *M. tuberculosis*) (MTBC), and bovine tuberculosis is also covered by Directive 2003/99/EC. Article 2 of the AHL states that its scope applies to transmissible diseases, including zoonoses, without prejudice to the rules laid down in Directive 2003/99/EC. Therefore, the annual zoonosis data reporting requirements for MSs in accordance with

<sup>&</sup>lt;sup>23</sup> Commission Implementing Regulation (EU) 2020/2002 of 7 December 2020 laying down rules for the application of Regulation (EU) 2016/429 of the European Parliament and of the Council with regard to Union notification and Union reporting of listed diseases, to formats and procedures for submission and reporting of Union surveillance programmes and of eradication programmes and for application for recognition of disease-free status, and to the computerised information system, OJ L 412, 8.12.2020, pp. 1–28.



18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (thps://onlinelibrary.wiley.com/terms

-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons

Directive 2003/99/EC, implemented by EFSA using specific tools, manuals and guidance, are not affected by the entry into force of CIR (EU) 2020/2002. The EU MSs need to report to the EU ADIS<sup>24</sup> on outbreaks of infection with MTBC in bovine (cattle, buffalo and bison), caprine and ovine animal populations, in other even-toed ungulates (Artiodactyla) and in terrestrial mammals. Annual summary reports are published online. CIR (EU) 2020/2002 details compulsory notification and annual reporting requirements from MSs to ADIS. MSs or zones thereof with disease-free status must also notify ADIS of MTBC outbreaks in terrestrial animals.

In accordance with the Zoonoses Directive 2003/99/EC, MSs must report annual surveillance data for bovine tuberculosis. These data originate from the compulsory national eradication and surveillance programmes that the MSs implement in accordance with EU legislation (Animal Health Law). The reports submitted by the MSs are harmonised and can be used for the assessment of the epidemiological situation and analysis of trends in MSs and MS zones.

In accordance with the EU legislation in force, disease-free status can be assigned by the European Commission to an MS or a zone thereof if during the previous 3 years at least 99.8% of the bovine herds have maintained their disease-free status from infection with MTBC, and the herd incidence rate during the last year has not exceeded 0.1%. Because the risk of infection with MTBC in disease-free zones is different from that of non-disease-free zones, these zones have been considered separately in this report.

Differently to the procedure for the above-mentioned statistics for humans, for which infections with *M. bovis* and *M. caprae* are treated separately, all cases of tuberculosis in cattle caused by the infectious MTBC members (*M. bovis*, *M. caprae*) are considered cases of bovine tuberculosis and were taken into account to summarise the EU situation on bovine tuberculosis. Whenever possible, reporting MSs distinguish descriptively between *Mycobacterium tuberculosis* complex, *M. bovis* and *M. caprae*. *M. tuberculosis* in cattle was not reported to EFSA during 2016–2020, whereas rare cases were reported earlier.

#### Mycobacterium surveillance data from food and from animals other than bovine animals

*Mycobacterium* spp. monitoring data from food and from animals other than bovine animals are submitted to EFSA in accordance with Directive 2003/99/EC. Data collected allow for descriptive summaries to be compiled at the EU level, but do not allow trend watching or trend analyses (Table 1).

In accordance with CIR (EU) 2020/2002, notification to ADIS and surveillance rules apply to other Artiodactyla (such as camelids, caprine animals and cervids) as species susceptible to infection with MTBC.

### 5.3. Results

#### 5.3.1. Overview of key statistics, EU, 2017–2021

Table 44 summarises the EU-level statistics on human tuberculosis due to *M. bovis* or *M. caprae* and on bovine tuberculosis during 2017–2021. More detailed descriptions of these statistics are provided in the below subsections.

**Table 44:** Summary statistics related to tuberculosis due to *Mycobacterium bovis* and *Mycobacterium caprae* in humans and bovine animals (stratified by disease-free<sup>(a)</sup> and non-disease-free status zones), EU, 2017–2021

	2021 <sup>(b)</sup>	2020	2019 <sup>(c)</sup>	2018 <sup>(c)</sup>	2017 <sup>(c)</sup>	Data source
Humans						
Number of confirmed <i>M. bovis</i> cases	103	96	141	168	204	ECDC
Number of confirmed M. caprae cases	8	3	11	13	9	ECDC
Total number of confirmed cases	111	99	152	181	213	ECDC
Total number of confirmed cases/100,000 population (notification rates)	0.03	0.03	0.03	0.04	0.05	ECDC

<sup>&</sup>lt;sup>24</sup> EU Animal Diseases Information System (ADIS). More information is available at https://food.ec.europa.eu/animals/animal-diseases/animal-disease-information-system-adis\_en



18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Isituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (thps://onlinelibrary.wiley.com/terms

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

	2021 <sup>(b)</sup>	2020	2019 <sup>(c)</sup>	2018 <sup>(c)</sup>	2017 <sup>(c)</sup>	Data source
Number of EU MSs that reported data on <i>M. bovis</i> or <i>M. caprae</i> cases	26	25	26	26	27	ECDC
M. bovis or M. caprae cases in individuals of EU origin	55	53	107	105	143	ECDC
M. bovis or M. caprae cases in individuals originating from outside the EU	47	35	40	68	62	ECDC
M. bovis or M. caprae cases in individuals of unknown origin	9	11	5	8	8	ECDC
Total number of foodborne outbreaks	0	0	0	0	0	EFSA
Number of outbreak-related cases	0	0	0	0	0	EFSA
Animals						
Bovine animals						
Number of infected herds in disease-free zones	139	139	143	172	134	EFSA
Number of reporting disease-free MSs	17	17	17	17	18	EFSA
Number of infected herds in non-disease-free zones	9,551	7,233	16,277	18,801	18,857	EFSA
Number of reporting non-disease-free MSs	11	9 <sup>(d)</sup>	11	11	10	EFSA

ECDC: European Centre for Disease Prevention and Control; EFSA: European Food Safety Authority; MSs: Member States.

- (a): The Member States or zones thereof with disease-free status from infection with the *Mycobacterium tuberculosis* complex (*M. bovis, M. caprae* and *M. tuberculosis*).
- (b): Data on animals from the United Kingdom (Northern Ireland) are taken into account for 2021. In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/Northern Ireland, the EU requirements on data sampling are also applicable to Northern Ireland.
- (c): Data from the United Kingdom are taken into account for 2017–2019, because the United Kingdom was an EU MS, but it became a third country on 1 February 2020.
- (d): No data were reported from Bulgaria.

### 5.3.2. Tuberculosis due to *Mycobacterium bovis* and *Mycobacterium caprae* in humans

In 2021, 111 confirmed human cases of tuberculosis due to *M. bovis* or *M. caprae* were reported by eight MSs (Austria, Belgium, Germany, Ireland, Italy, the Netherlands, Spain and Sweden) (Table 45). Tuberculosis cases due to *M. bovis* (103 cases) were reported in all these countries, and only three MSs (Austria, Germany and Spain) also notified cases caused by *M. caprae* (8 cases). Overall, tuberculosis cases due to *M. bovis* or *M. caprae* accounted for a small proportion (0.3%) of total tuberculosis cases reported in the EU by 26 MSs with species-specific data on MTBC available in 2021 (see Section 5.2.1). Eighteen MSs did not report any cases.

The EU notification rate in 2021 was 0.03 cases per 100,000 population, which is similar to the rate observed in 2020 (Table 45). The EU notification rate decreased by 28.3% and 25.3% compared with the mean notification rate observed over the pre-pandemic years 2017–2019 with and without the data from the United Kingdom, respectively.

In 2021, the highest notification rate was reported by Belgium (0.09 per 100,000), followed by Spain (0.07 per 100,000).

Among the 17 MSs with disease-free status in 2021, 16 MSs reported on MTBC species. *M. bovis* and *M. caprae* human cases were reported in five MSs. The notification rate in these 16 MSs reporting on MTBC species was 0.026 cases per 100,000 population. The same notification rate for *M. bovis* and *M. caprae* human cases (0.026 cases per 100,000 population) was also observed in the 10 MSs with non-disease-free status in 2021.

Approximately half of the cases reported in 2021 (55/111; 49.5%) were of EU origin (native cases and/or cases originating from other MSs). The other cases originated from outside the EU (N = 47; 42.3%) or had unknown origin (N = 9; 8.1%) (Table 44). Notification rates of M. bovis and M. caprae human cases of EU origin were similar in disease-free MSs (N = 27; 48.2%) and non-disease-free MSs (N = 29; 51.8%).

Information on treatment outcome after 12 months was reported for 91.9% (91/99) of the human *M. bovis* and *M. caprae* cases notified in 2020. The treatment was reported to be successful for



58 cases (63.7%). Three cases (3.3%) were still being treated at 12 months. Deaths were reported in 10 cases (11.0%). Twenty cases (22.0%) were reported as lost to follow-up.

Regarding drug resistance, only one case was reported to be resistant to isoniazid treatment out of 77 *M. bovis* or *M. caprae* cases with test results reported in 2021 for rifampicin and isoniazid. This finding confirms that drug resistance is relatively rare among *M. bovis* or *M. caprae*, considering that neither rifampicin-resistant nor multidrug-resistant cases were reported in 2021.

**Table 45:** Reported cases of human tuberculosis due to *Mycobacterium bovis* and *Mycobacterium caprae* and notification rates per 100,000 population in EU MS and non-MS countries by country and year, 2017–2021

		2	2021			20	20	20	19	20	18	20	17
Country	Status	National coverage	Data format <sup>(a)</sup>	cases	rmed s and tes	cases	rmed s and tes	Confi cases rat	s and	cases	rmed s and tes	cases	rmed s and tes
				Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate
Austria	DFS	Υ	С	4	0.04	0	0	3	0.03	2	0.02	2	0.02
Belgium	DFS	Υ	С	10	0.09	6	0.05	0	0	5	0.04	6	0.05
Bulgaria		Y	С	0	0	0	0	0	0	0	0	0	0
Croatia		Y	С	0	0	0	0	0	0	0	0	0	0
Cyprus		Y	С	0	0	0	0	0	0	0	0	0	0
Czechia	DFS	Υ	С	0	0	0	0	0	0	1	0.01	0	0
Denmark	DFS	Υ	С	0	0	0	0	0	0	0	0	1	0.02
Estonia	DFS	Υ	С	0	0	0	0	0	0	0	0	0	0
Finland	DFS	Υ	С	0	0	1	0.02	0	0	0	0	0	0
France <sup>(b)</sup>	DFS	_	_	_	_	_	_	_	_	_	_	_	_
Germany	DFS	Υ	С	42	0.05	37	0.04	51	0.06	64	0.08	48	0.06
Greece		Y	С	0	0	2	0.02	1	0.01	0	0	1	0.01
Hungary	DFS	Υ	С	0	0	0	0	0	0	0	0	0	0
Ireland		Υ	С	2	0.04	4	0.08	7	0.14	7	0.14	4	0.08
Italy		Y	С	12	0.02	6	0.01	11	0.02	17	0.03	21	0.03
Latvia	DFS	Υ	С	0	0	_	_	_	_	_	_	0	0
Lithuania	DFS	Υ	С	0	0	0	0	0	0	0	0	0	0
Luxembourg	DFS	Υ	С	0	0	0	0	0	0	0	0	0	0
Malta		Y	С	0	0	0	0	0	0	0	0	0	0
Netherlands	DFS	Υ	С	5	0.03	6	0.03	5	0.03	11	0.06	11	0.06
Poland	DFS	Υ	С	0	0	0	0	0	0	0	0	0	0
Portugal		Υ	С	0	0	0	0	0	0	0	0	0	0
Romania		Y	С	0	0	1	0.01	1	0.01	0	0	2	0.01
Slovakia	DFS	Υ	С	0	0	0	0	0	0	0	0	0	0
Slovenia	DFS	Υ	С	0	0	0	0	0	0	0	0	0	0
Spain		Y	С	32	0.07	30	0.06	35	0.07	46	0.10	73	0.16
Sweden	DFS	Υ	С	4	0.04	6	0.06	3	0.03	4	0.04	3	0.03
EU Total 27		_	_	111	0.03	99	0.03	117	0.03	157	0.04	172	0.05
United Kingdom <sup>(c)</sup>	_		-	_	-	_	-	35	0.05	24	0.04	41	0.06
EU Total		_	_	111	0.03	99	0.03	152	0.03	181	0.04	213	0.05
Iceland <sup>(d)</sup>		Y	С	_	_	0	0	0	0	0	0	0	0
Norway	DFS	Y	С	0	0	0	0	1	0.02	0	0	3	0.06
Liechtenstein	DFS	Y	С	0	0	3	0.03	4	0.05	3	0.04	3	0.04
Switzerland <sup>(e)</sup>	DFS	Υ	С	3	0.03								

18314732, 2022, 12, Downloaded from https://eftsaonlinelibrary.wiley.com/doi/10.2903j.eftsa.2022.7666 by Islituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18122023]. See the Terms and Conditions, wiley convierms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Ceravive Commons.

com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilattico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms


- EU: European Union.
- -: Data not reported.
- DFS: Disease-free status, i.e. free of infection with M. bovis and M. caprae in bovine animal population.
- (a): Y: yes; N: no; A: aggregated data; C: case-based data.
- (b): Not reporting species of the *M. tuberculosis* complex.
- (c): Cases reported by the United Kingdom for the period 2017–2019 were also considered for this estimation (EU-28). When the United Kingdom data were collected for the period 2017–2019, the United Kingdom was an EU MS, but it became a third country on 1 February 2020.
- (d): In Iceland, which has no special agreement regarding animal health (status) with the EU, the last outbreak of bovine tuberculosis was in 1959.
- (e): Switzerland provided data directly to EFSA. The human data for Switzerland include data from Liechtenstein for the years 2017–2020.

DFS All Zones of the MS have disease-free status.

Not all Zones of the MS have disease-free status.

No Zones of the MS have disease-free status.

Figure 11 shows, for the year 2021, the number of confirmed tuberculosis cases due to *M. bovis* and to *M. caprae* in individuals of EU origin overlaid with the national aggregated herd prevalence of bovine tuberculosis.



**Figure 11:** Map of the number of confirmed tuberculosis cases due to *Mycobacterium bovis* and *Mycobacterium caprae* in individuals of EU origin, and national herd prevalence of tuberculosis in the bovine population in EU MS and non-MS countries, 2021



Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

#### 5.3.3. Mycobacterium in food

No Mycobacterium spp. monitoring data from food were submitted for the year 2021.

#### 5.3.4. Tuberculosis in bovine animals


#### **Bovine tuberculosis surveillance data**

Seventeen MSs had DFS from MTBC during 2021 (Figure 12). Of the remaining 10 MSs and the United Kingdom (Northern Ireland), three MSs had disease-free zones or provinces:

- Italy: 11 regions and 12 provinces;
- Portugal: 1 region (Algarve) and all of the Azores islands, except São Miguel;
- Spain: 3 autonomous communities (Canary Islands, Galicia and Asturias).

Seven MSs had no DFS from MTBC zones. The United Kingdom (Northern Ireland) has no disease-free region either.

Norway, Switzerland and Liechtenstein had disease-free status, in accordance with the EU legislation. In Iceland, which has no special agreement with the EU on animal health status, the last outbreak of bovine tuberculosis was reported in 1959.



**Figure 12:** Status of countries for infection with the *Mycobacterium tuberculosis* complex (*M. bovis, M. caprae* and *M. tuberculosis*) in the bovine animal population, EU MSs and non-MSs, 2021

During 2021, in the EU, the overall prevalence of cattle herds infected with MTBC was very low (9,690 out of 1,726,451 herds; 0.6%), slightly higher than 2020 (0.4%). Compared with the previous year, the number of bovine herds infected with MTBC in the EU increased from 7,372 to 9,690 herds in



18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilattico

Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

(https://onlinelibrary.wiley.c

2021. This increase was mainly due to 2021 data from the United Kingdom (Northern Ireland), which has a high prevalence, whereas its 2020 data were not included in the overall EU statistics. Thirteen MSs (11 disease-free and 2 non-disease-free) reported no cases of bovine tuberculosis in cattle (Table 46). The remaining MSs reported bovine tuberculosis with a wide range of prevalence at the national level. MTBC infection in cattle herds primarily affected the EU zones with non-disease-free status: the overall occurrence of bovine tuberculosis in non-disease-free zones (1.3%) was 92.8 times higher than in disease-free zones (0.014%).

Table 46: Status of countries as regards bovine tuberculosis and related prevalence, EU, 2021

Member state (MS)	Status	N of infected herds in disease- free zones	Prevalence (%) of infected herds in DFS <sup>(a)</sup> zones	N of infected herds in non- DFS <sup>(a)</sup> zones	Prevalence (%) of infected herds in non-DFS <sup>(a)</sup> zones
Austria	DFS	6 <sup>(b)</sup>	0.01	_	_
Belgium	DFS	5 <sup>(c)</sup>	0.02	_	_
Bulgaria		_	_	6	0.01
Croatia		_	_	2	0.01
Cyprus		_	_	0	0
Czechia	DFS	0	0	_	_
Denmark	DFS	0	0	_	_
Estonia	DFS	0	0	_	_
Finland	DFS	0	0	_	_
France	DFS	99 <sup>(c)</sup>	0.06	_	_
Germany	DFS	8	0.01	-	_
Greece		_	_	140	0.76
Hungary	DFS	4 <sup>(c)</sup>	0.02	_	_
Ireland		_	_	5,071	4.6
Italy		9 <sup>(c)</sup>	0.02	179	0.35
Latvia	DFS	0	0	_	_
Lithuania	DFS	0	0	_	_
Luxembourg	DFS	0	0	_	_
Malta		_	_	0	0
Netherlands	DFS	0	0	_	_
Poland	DFS	8 <sup>(c)</sup>	< 0.01	_	_
Portugal		0	0	99	0.35
Romania		_	_	31	0.01
Slovakia	DFS	0	0	_	_
Slovenia	DFS	0	0	_	_
Spain		0	0	1,424	1.33
Sweden	DFS	0	0	_	_
United Kingdom (Northern Ireland)		-	-	2,599	11.3
EU Total (27 + XI)		139	0.014	9,551	1.3

MS: Member State; DFS: Disease-free status.

(c): Only Mycobacterium bovis identified.

DFS All Zones of the MS have disease-free status.

Not all Zones of the MS have disease-free status.

No Zones of the MS have disease-free status.

<sup>(</sup>a): Data from the United Kingdom (Northern Ireland) are taken into account for 2021. In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/Northern Ireland, the EU requirements on data sampling also apply to Northern Ireland, which is non-disease-free.

<sup>(</sup>b): Only Mycobacterium caprae identified.




18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Isituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (thps://onlinelibrary.wiley.com/terms

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

#### MSs and MSs' zones with disease-free status from Mycobacterium tuberculosis complex

The majority of the whole EU cattle herd population (56.1%) is located in the disease-free zones of the 20 MSs having such zones. However, the number of cattle herds has been steadily decreasing (-29.4% in 2021 compared with 2012), impacting on the calculation of the prevalence of herds infected with MTBC. From 2012 to 2021, there was a marked decrease in the annual number of infected cattle herds (-38.8%), whereas the prevalence has remained stable. Infected herds were rare events (Figure 13).

Seven MSs with disease-free zones reported a total of 139 MTBC-infected bovine herds (Table 46), confirming that the detection of bovine tuberculosis in disease-free zones is sporadic. When comparing data from 2021 and 2020, the number of MTBC-infected cattle herds remained the same, with an almost unchanged prevalence  $(0.013\% \ vs.\ 0.014\%)$ , whereas the total number of cattle herds has decreased by 6.5%.



- (\*): In contrast to years 2012–2019, the year 2020 does not include United Kingdom data. Since 1 February 2020, the United Kingdom has withdrawn from the EU and has become a third country.
- (\*\*): In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/Northern Ireland, the European Union requirements on data sampling also apply to Northern Ireland.

**Figure 13:** Prevalence of cattle herds infected with *Mycobacterium tuberculosis* complex in disease-free (DFS) zones, EU, 2012–2021

### MSs and MSs' zones with non-disease-free status from Mycobacterium tuberculosis complex

The zones with non-disease-free status, belonging to 10 MSs and the United Kingdom (Northern Ireland), account for 43.9% of the whole EU cattle herd population, which has been steadily decreasing over time (–45.2% in 2021 compared with 2012; Figure 14). The significant decrease in 2020 is due to the withdrawal of the United Kingdom from the EU. Moreover, Bulgaria did not report data in that year. The increase in the number of 2021 cattle herds can be attributed to the data provided by Bulgaria after its 2020 hiatus and by the United Kingdom (Northern Ireland).


Eight MSs and the United Kingdom (Northern Ireland) reported a total of 9,551 cattle herds infected with MTBC (Table 46). Over the last 10 years, the annual number of infected cattle herds in



non-disease-free zones has decreased (-47.5% in 2021 compared with 2012). During the same period, the overall prevalence of bovine tuberculosis decreased from 1.32% to 1.26% (-4.2%). This decrease can be attributed to the withdrawal of the United Kingdom from the EU in 2020.

In 2021, prevalence varied widely among MSs with non-disease-free status: Ireland (4.6%) and Spain (1.3%) were the only MSs that reported prevalence higher than 1%; no infected herds were reported by Malta or Cyprus (Table 46). The United Kingdom (Northern Ireland), which has no disease-free zones, reported prevalence of 11.3%. Compared with 2020, the prevalence of infected herds increased in 2021, mainly due to the inclusion of United Kingdom (Northern Ireland) data, which were not included in the 2020 EU-level statistics.

Most MSs reported MTBC infections without specifying the species involved. Infection with *M. bovis* was specifically detected in Bulgaria, France, Germany, Hungary, Ireland, Italy, Poland, Romania and the United Kingdom (Northern Ireland), whereas infection with *M. caprae* was specifically reported by Austria, Germany and Romania. *M. tuberculosis* in cattle was not reported.



- (\*): In contrast to years 2012–2019, the year 2020 does not include the United Kingdom data. Since 1 February 2020, the United Kingdom has withdrawn from the EU and has become a third country. No data were reported from Bulgaria.
- (\*\*): In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/Northern Ireland, the European Union requirements on data sampling also apply to Northern Ireland.


**Figure 14:** Prevalence of cattle herds infected with *Mycobacterium tuberculosis* complex in non-disease-free (non-DFS) zones, EU, 2012–2021

Figure 15 shows the prevalence trends in the five MSs (Greece, Ireland, Italy, Spain, Portugal) and the United Kingdom (Northern Ireland) presenting the highest prevalence between 2005 and 2021. In most of the MSs (Greece, Ireland, Spain) and the United Kingdom (Northern Ireland), no significant improvements were observed over time given that these data refer to zones where activities related to eradication programmes are more difficult due to unfavourable local environmental conditions (free-range animals, common grazing land, transhumance, etc.) (Ciaravino et al., 2021; Byrne et al., 2022). However, in Italy, Spain and Portugal, several zones obtained disease-free status during this period.

18314732, 2022, 12, Downloaded from https://efs.aonlinelibray.wiley.com/doi/10/2033/jfs/a2022/7666 by Istituto Zoproficitieto's Speriment ad ell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023], See the Terms and Conditions (https://onlinelibray.wiley.com/ems-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons.

com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilattico

Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [1812/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA article are governed by the applicable Creative Common



**Figure 15:** Prevalence of bovine tuberculosis-infected herds in non-disease-free zones of five MSs and the United Kingdom (Northern Ireland), 2005–2021

Non-Member States and pre-accession countries

Bovine tuberculosis was not detected in 2021 in Iceland, Liechtenstein, Norway or Switzerland. Among the pre-accession countries, Montenegro, as in 2020, reported no infected herds, whereas the Republic of North Macedonia reported a prevalence of 0.13% MTBC-infected herds (20/14,898), a value slightly lower than that reported in 2020 (0.15%).

### Mycobacterium surveillance data from animals other than bovine animals, and complementary reporting from cattle

In 2021, three MSs (Finland, Slovenia, Sweden) and one non-MS (Norway) reported surveillance data on infection in farmed deer, with no cases of MTBC infection detected. For the first time in 2021, Greece reported cases of infection with MTBC in goats via ADIS.

Complementary to their 2021 reports from cattle, MSs also reported cases of MTBC infection in other mammal species. In particular, *M. bovis* was detected in farmed alpacas, small ruminants (sheep and goats), pigs, cats, wild deer, wild boars and badgers, whereas *M. caprae* was reported in wild red deer in Austria and in wild boars in Hungary.

#### 5.4. Discussion

In 2021, the reporting of human cases of tuberculosis due to *M. bovis* and *M caprae* at the EU level was the lowest in the last 5 years, with the exception of 2020 and taking into account the withdrawal of the United Kingdom from the EU. This finding should be interpreted with caution. It may be an important indication of the successful application of the eradication programmes for tuberculosis in bovine animals in EU MSs. On the other hand, it may be a consequence of the disruption in healthcare services due to COVID-19. Like for other zoonoses, the reporting of fewer *M. bovis* and *M. caprae* cases in 2020, i.e. the first pandemic year, may reflect the impact of the COVID-19 pandemic on healthcare systems, even for a long-standing condition such as tuberculosis. A WHO document released in early in the COVID-19 pandemic in Europe called attention to the need to maintain the continuity of essential services for people affected with tuberculosis during the COVID-19 pandemic (WHO, 2020). Tuberculosis-related services and programmes were perturbed by the COVID-19

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Isituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (thps://onlinelibrary.wiley.com/terms

-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons

pandemic and the impact of the COVID-19 pandemic on tuberculosis services — leading to hindered access to care, delayed diagnosis and delayed reporting — was particularly severe in 2020 (Migliori et al., 2021; Nikolayevskyy et al., 2021). The increased reporting of cases in 2021 at the EU level and in many MSs can be considered a sign of the healthcare systems' resilience during the second pandemic year. The significant reduction in the proportion of *M. bovis* and *M. caprae* cases lost to follow-up for the evaluation of treatment outcome after 12 months is another important indication of a progressive recovery of tuberculosis programmes to their pre-pandemic capacity. In 2020, this proportion accounted for almost half (49.3%) of the cases with reported treatment outcome, whereas in 2021 it was only 22.0%, although this proportion is still much higher compared with pre-pandemic years (lost-to-follow-up cases were 1.2% in 2019, 2.2% in 2018 and 2.4% in 2017). However, compared to *M. tuberculosis*, *M. bovis* and *M. caprae* represent a minor proportion of all MTBC cases in the human population and the limited number of human *M. bovis* and *M. caprae* cases involves some degree of uncertainty, which hampers coming to definitive conclusions.

The characteristics of *M. bovis* and *M. caprae* cases reported in 2021 and indicators of treatment outcome among cases occurred in 2020 did not substantially differ from those reported in prepandemic years. Successful treatment and deaths were reported in 63.7% and 11% of cases, respectively. This finding suggests that there was no direct effect of COVID-19 on patients with *M. bovis* and *M caprae*, albeit the high proportion of lost-to-follow-up cases challenges this conclusion.

The notification rate of *M. bovis* and *M. caprae* cases in 2021 was the lowest among all zoonoses analysed in this report, apart from rabies, meaning that exposure of humans to *M. bovis* and *M. caprae* and the development of clinical disease is apparently a rare event, although the consequences are severe for patients. It is possible that the annual number of cases of zoonotic tuberculosis in the EU and the overall burden may be much higher, because not all MSs implement routine surveillance able to discriminate *M. bovis* and *M. caprae* cases from *M. tuberculosis* cases. Moreover, lower access to healthcare services by the non-EU native population, which account for 42.3% of the total number of tuberculosis cases, may also contribute to the underdiagnosis or underreporting of *M. bovis* and *M. caprae* cases.

The high proportion of deaths among patients evaluated for treatment outcome that are reported every year at the 12-month follow-up argues for the need to continue eradication programmes in the bovine population to reduce the risk of exposure to *M. bovis* and *M. caprae* in the human population. A call for action launched in 2017 acknowledged the importance of a cross-sectorial and multidisciplinary approach linking animal, human and environmental health, particularly to protect the most vulnerable and marginalised communities (Olea-Popelka et al., 2017).

*M. bovis* and *M. caprae* notification rates did not differ significantly between EU countries with disease-free status in the bovine population and those with non-disease-free status. This finding appears to suggest that the health status of the bovine population does not directly influence the risk of human infection by *M. bovis* and *M. caprae*, which remains a rare event and therefore highly subject to peculiar individual habits and behaviours. On the other hand, notification rates are strongly influenced by the sensitivity of surveillance at the country level and differences in case reporting among MSs are considerable. For example, Germany which is a disease-free country regarding its bovine population, reported more than one third of the total *M. bovis* and *M. caprae* human cases notified in the EU (42 cases, 37.8%), revealing highly sensitive surveillance. Data reported to ECDC show that the proportion of MSs not reporting any *M. bovis* and *M. caprae* cases in the last 5 years were higher among MSs with non-disease-free status in the bovine population (5/10) than among MSs with disease-free status (7/16; 2021). This comparison suggests lower surveillance sensitivity in some MSs in the non-disease-free status group.

*M. bovis* and *M. caprae* cases originating from outside the EU were reported in 2021 (42.3%) and 2020 (35.4%) in higher proportions compared with MTBC cases originating from outside the EU and EEA in 2020 (33.0%) (ECDC and WHO, 2022). Although these statistics are not fully comparable among the two groups because the 2021 data do not include cases originating from EEA countries, this difference raises the question of exposure to *M. bovis* and *M. caprae* being higher in people originating from outside the EU. For example, employees in slaughterhouses and cattle farms are frequently non-EU natives, which may lead to differential occupational exposure to aerosols from infected animals and their carcases in this population.

The regulatory framework for bovine tuberculosis changed substantially in the EU during 2021. Tuberculosis caused by MTBC infection is currently considered a disease to be controlled in all MSs with the goal of eradicating the disease in bovine animals throughout the EU, while being kept under surveillance in other mammals. All MSs must have a surveillance or a control-and-eradication

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Islituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.

programme approved by the European Commission. This should lead, in the coming years, to progressive improvement in the already satisfactory epidemiological situation. In 2021, the overall EU prevalence of cattle herds infected with MTBC was 0.6%, slightly higher than in 2020. Bovine tuberculosis was reported by 14 MSs and by the United Kingdom (Northern Ireland). Its distribution was highly heterogeneous and spatially clustered in the EU, with herd prevalence ranging from 0% to 11.3%, confirming the results of a previous EFSA report (EFSA AHAW Panel, 2014).

Seventeen MSs had disease-free status and three non-disease-free MSs also had such zones. The number of disease-free zones increased during 2021. Thirteen of these MSs reported no cases of bovine tuberculosis in cattle. In the disease-free zones, the detection of bovine tuberculosis remained a rare event, as in the previous years. From 2012 to 2021, the overall annual number of infected cattle herds, the prevalence and the total number of cattle herds decreased.

Eight non-disease-free MSs and the United Kingdom (Northern Ireland) detected bovine tuberculosis during 2021, with an overall infected herd prevalence of 1.3%. When comparing 2021 with 2020 data, the overall annual number of infected cattle herds, the prevalence and the total number of cattle herds increased in these non-disease-free zones. This increase is mainly due to the withdrawal of the United Kingdom from the EU in 2020 and to the subsequent (2021) inclusion of data from the United Kingdom (Northern Ireland), a region with high bovine tuberculosis prevalence. In the last 10 years (2012–2021), the overall annual number of infected cattle herds reported in the non-disease-free zones decreased by 47.5%, whereas prevalence decreased only by 4.2%. These different trends can partly be attributed to (1) the withdrawal of the United Kingdom from the EU; (2) the decreased number of herds; (3) the gradual obtention of disease-free status in zones within non-disease-free MSs and (4) unfavourable environmental conditions hindering the eradication process in several non-disease-free zones (Byrne et al., 2022).

There is a major obstacle to bovine tuberculosis eradication in cattle in areas where infection is endemic in wildlife animals. Successfully tackling bovine tuberculosis also involves addressing the wildlife reservoir of the disease. Ireland introduced a vaccination policy in the Eurasian badger (*Meles meles*), a species known as a maintenance host of *M. bovis*, in 2018 and is also, among other control measures, reducing the badger population (Gormley and Corner, 2013). On the other hand, in the United Kingdom (Northern Ireland), no active badger intervention or vaccination took place in 2021. Stagnating or increasing trends in the prevalence of bovine tuberculosis-infected cattle herds demonstrate that eradication of this disease is a challenge, owing to the complex interactions between the pathogen, hosts and the local environments (EFSA AHAW Panel, 2014). MS-specific evaluations of status, trends and of the relevance of bovine tuberculosis as a source of disease for humans can be found in the 2021 Annual National Zoonoses Country Reports referenced in Section 5.5.

In 2021, *M. bovis* was reported to be isolated – apart from bovine animals – from a wide range of mammal species, both domestic and wild, reflecting that this pathogen has a broad host range. *M. caprae*, acknowledged to cause bovine tuberculosis, was reported in cattle, and also in wild boar and red deer.




18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilattico

dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

#### 6. Brucella

Summary data substantiating this chapter, as well as additional information on related projects and internet sources are published for this report on the EFSA Knowledge Junction on Zenodo here. Summary statistics on brucellosis foodborne outbreaks reported in the framework of Directive 2003/99/EC, with downloadable files, are retrievable using the EFSA foodborne outbreaks dashboard available here. Summary statistics on human surveillance data with downloadable files are retrievable using the ECDC Surveillance Atlas of Infectious Diseases here.

# Brucella in the EU, 2021




### **Human cases in foodborne outbreaks**



### Foodborne outbreaks

Food vehicles causing strong-evidence outbreaks



and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common



- In 2021, the number of confirmed cases of human brucellosis was 162 in the EU. The EU notification rate of 0.03 per 100,000 population was the same as that reported in 2020, which was the lowest since surveillance began in the EU in 2007.
- Compared with the rate before the COVID-19 pandemic (2017–2019 annual mean), there was a decrease of 52.2% and 51.1% with and without the data from the United Kingdom, respectively.
- From 2017 to 2021 there has been a significantly declining trend of confirmed human cases of brucellosis in the EU.
- Three MSs (Germany, Greece and Italy) have had significantly decreasing 5-year trends from 2017 to 2021.
- Brucella melitensis was reported as the aetiological agent in 51 (87.9%) out of 58 human cases of brucellosis with information reported on the Brucella species.
- Croatia became disease-free from infection with *B. abortus*, *B. melitensis* and *B. suis*<sup>25</sup> in 2021 in bovine animal populations. In total, 21 MSs and the United Kingdom (Northern Ireland) were disease-free while 6 MSs (Bulgaria, Greece, Hungary, Italy, Portugal and Spain) were non-disease-free. Overall, in the disease-free zones of the EU, there were eight infected herds in 2021, demonstrating a rare occurrence (prevalence < 0.001%). In the non-disease-free zones, bovine brucellosis remained very low, with 546 herds reported to be infected (0.43%). The number of infected herds stabilised at between 648 and 485 during 2017–2021.
- Spain became disease-free from infection with *B. abortus*, *B. melitensis* and *B. suis* in ovine and caprine populations in 2021. In total, 20 MSs and the United Kingdom (Northern Ireland) were disease-free while seven MSs (Bulgaria, Croatia, overseas French regions, Greece, Italy, Malta and Portugal) were non-disease-free. Overall, in the disease-free zones of the EU, there were 15 infected herds in 2021, demonstrating an extremely low prevalence (< 0.01%). In the non-disease-free zones, brucellosis in sheep and goats remained very low, with 331 herds reported to be infected (0.18%). The number of infected herds in these zones decreased from 815 in 2017 to 331 in 2021.
- Brucellosis is still an animal health concern with public health relevance in southern European countries that are not brucellosis disease-free.

#### 6.2. Surveillance and monitoring of *Brucella* in the EU

#### 6.2.1. Humans

In 2021, 26 MSs reported information on brucellosis in humans. Surveillance is mandatory in 25 MSs. Belgium has another (unspecified) system. The surveillance systems cover the whole population in all MSs reporting brucellosis data. Denmark has no surveillance system in place for brucellosis, and the disease is not notifiable or reported at the EU level. All countries reported case-based data except Belgium and Bulgaria, which reported aggregated data. The EU case definition was used by 23 of these 26 countries, all of which had a comprehensive surveillance system. France reported using a different case definition, and Germany and Italy did not specify which case definition was used.

In 2021, the regulatory framework for brucellosis in animals changed substantially in the EU. Infection with *B. abortus*, *B. melitensis* and *B. suis* in bovine animals, and sheep and goats became a disease to be controlled in all MSs with the goal of eradicating the disease in these animals throughout the EU, while being kept under surveillance in other even-toed ungulates (Artiodactyla). Previous, repealed legislation established requirements relating to Officially brucellosis-free status in cattle (OBF) and/or officially *B. melitensis*-free status in sheep and goats (ObmF), in certain Member States and regions of Member States.



#### 6.2.2. Food and animals

# Monitoring data for *Brucella* from bovine animals, and from sheep and goats, originating from the National Control and Eradication Programmes and/or from countries or zones with disease-free status

On 21 April 2021, Commission Implementing Regulation (CIR) (EU) 2020/2002<sup>26</sup> under the Animal Health Law (AHL) entered into force, with Union notification and reporting covering infection with *Brucella abortus*, *B. melitensis* and *B. suis*. Reporting on brucellosis (any *Brucella* species) is also covered by the Directive 2003/99/EC, Annex I list A 'brucellosis and agents thereof'. Article 2 of the AHL states that its scope applies to transmissible diseases, including zoonoses, without prejudice to the rules laid down in Directive 2003/99/EC. Therefore, the zoonoses annual data reporting requirements for MSs in accordance with Directive 2003/99/EC, implemented by the European Food Safety Authority (EFSA) using specific tools, manuals and guidance, are not affected by the entry into force of the CIR (EU) 2020/2002. The EU MSs need to report outbreaks of infection with *Brucella abortus*, *B. melitensis* and *B. suis* in bovine, caprine and ovine animal populations, and other terrestrial animals to the EU ADIS, <sup>27</sup> and annual summary reports are published online. CIR (EU) 2020/2002 details compulsory notification and annual reporting requirements from MSs to ADIS.

In accordance with the EU legislation in force, disease-free status can be assigned by the European Commission to an MS or a zone thereof if during the previous 3 years at least 99.8% of bovine herds have maintained their disease-free status from infection with brucellosis, and the herd incidence rate during the last year has not exceeded 0.1%. Because the risk of infection with brucellosis in disease-free zones is different from that of non-disease-free zones, these zones have been considered separately in this report.

Regulation (EU) 2016/429 of the European Parliament and of the Council concerns approval of the disease-free and non-vaccination status of MSs or zones or compartments. CIR (EU) 2018/1882 includes Bos ssp., Bubalus ssp., Bison ssp., Ovis ssp., Capra ssp. as species for notification, surveillance, prevention, certification and compulsory eradication. Commission Delegated Regulation 2020/689 (Annex IV) describes the conditions required to grant, maintain or suspend the status of free from infection with B. abortus, B. melitensis and B. suis for an establishment keeping bovine animals, and sheep and goats.

The reports submitted by the MSs are based on Regulation (EU) 2016/429 and subsequent legislation and are essential for assessment of the epidemiological situation in MSs and MS zones, whether declared disease-free (DF) in cattle and/or in sheep and goats.

Annual surveillance programmes are carried out in DFS zones to confirm the absence of infection with *Brucella abortus*, *B. melitensis* and *B. suis* in cattle, and in sheep and goats. In non-DF zones, control and eradication programmes for brucellosis in cattle, and sheep and goats are in place. These data are comparable across MSs because the monitoring schemes are harmonised, and the data collected and reported to EFSA originate from the census-as-sampling framework or a randomised design. These data can be used to carry out trend analyses both at the EU and MS levels, to perform trend watching and produce descriptive summaries and to assess the impact of control and eradication programmes (Table 1).

## Monitoring data for *Brucella* from food, and animals other than bovine animals, and sheep and goats

Monitoring data for *Brucella* from food, and animals other than bovine animals, and sheep and goats are submitted to EFSA in accordance with Directive 2003/99/EC. They preclude trend analyses and trend watching at the EU level (Table 1). In accordance with CIR (EU) 2020/2002, notification to ADIS and surveillance rules apply to other Artiodactyla species (such as camelids or cervids) as susceptible species for brucellosis. Data collected allow for descriptive summaries to be compiled at the EU level.

#### 6.3. Results

#### 6.3.1. Overview of key statistics, EU, 2017–2021

Table 47 displays statistics at the EU level on human and animal brucellosis, along with data on detection of *Brucella* in food, between 2017 and 2021. Results are described in detail in this chapter.

EFSA Journal 2022;20(12):7666

<sup>&</sup>lt;sup>26</sup> Commission Implementing Regulation (EU) 2020/2002 of 7 December 2020 laying down the rules for the application of Regulation (EU) 2016/429 of the European Parliament and of the Council with regard to Union notification and Union reporting of listed diseases, to formats and procedures for submission and reporting of Union surveillance programmes and of eradication programmes, and for application for recognition of disease-free status, and to the computerised information system, OJ L 412, 8.12.2020, pp. 1–28.

<sup>&</sup>lt;sup>27</sup> EU Animal Diseases Information System (ADIS). More information is available at: https://food.ec.europa.eu/animals/animal-diseases/animal-disease-information-system-adis\_en

**Table 47:** Summary of *Brucella* statistics related to humans, major food categories and main animal species (stratified by disease-free status<sup>(a)</sup> and non-disease-free status zones), EU, 2017–2021

	2021 <sup>(b)</sup>	2020	2019 <sup>(c)</sup>	2018 <sup>(c)</sup>	2017 <sup>(c)</sup>	Data source
Humans						
Total number of confirmed cases	162	132	309	332	378	ECDC
Total number of confirmed cases/100,000 population (notification rates)	0.03	0.03	0.06	0.08	0.09	ECDC
Number of reporting MSs	26	26	27	26	26	ECDC
Infection acquired in the EU	76	68	126	133	148	ECDC
Infection acquired outside the EU	21	14	50	51	46	ECDC
Unknown travel status or unknown country of infection	65	50	133	148	184	ECDC
Number of outbreak-related cases	2	2	2	0	2	EFSA
Total number of outbreaks	1	1	1	0	1	EFSA
Food						
Milk and milk products						
Number of sampling units	320	275	586	1,005	1,338	EFSA
Number of reporting MSs	3	3	2	3	3	EFSA
Animals						
Cattle (Bovine animals)						
Number of infected herds in disease-free zones	8	6	4	3	0	EFSA
Number of reporting disease-free MSs	22	19	20	20	20	EFSA
Number of infected herds in non-disease-free zones	546	603	485	563	648	EFSA
Number of reporting non-disease-free MSs	6	7	8	8	8	EFSA
Sheep and goats						
Number of infected herds in disease-free zones	15	3	1	0	7	EFSA
Number of reporting disease-free MSs	22	19	20	20	20	EFSA
Number of infected herds in non-disease-free zones	331	349	451	620	815	EFSA
Number of reporting non-disease-free MSs	6	7 <sup>(d)</sup>	8	8	8	EFSA

ECDC: European Centre for Disease Prevention and Control; EFSA: European Food Safety Authority; MSs: Member States.

#### 6.3.2. Human brucellosis

In 2021, 162 confirmed cases were reported in the EU, which was a slight increase compared to 2020. The notification rate was 0.03 cases per 100,000 population, which is equal to the notification rate in 2020. In 2021, 26 MSs provided data and information on brucellosis in humans (Table 47). Compared with the rate before the COVID-19 pandemic (2017–2019 mean annual), there was a decrease of 52.2% and 51.1%, with and without the data from the United Kingdom, respectively.

In 2021, 10 MSs (Bulgaria, Estonia, Finland, Hungary, Ireland, Latvia, Lithuania, Malta, Romania and Slovenia) reported zero cases (Table 48).

1831/4372, 2022, 1,2 Downloaded from https://ef.son.inlinibitary.wiley.com/oid/10/2003/jefssa2022/7666 by Islituic Copyor/Inlinib Sperimental dell'Umbria e delle Marche (ISUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://oillneibbrary.wiley.com/ems-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

<sup>(</sup>a): Disease-free status from infection with *B. abortus*, *B. melitensis* and *B. suis* in bovine animal populations, and/or ovine and caprine animal populations.

<sup>(</sup>b): Data on food and animal samples from the United Kingdom (Northern Ireland) are taken into account for 2021. In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/Northern Ireland, EU requirements on data sampling are also applicable to Northern Ireland.

<sup>(</sup>c): Data from the United Kingdom are taken into account for 2017–2019, because the United Kingdom was an EU MS over this period but it became a third country on 1 February 2020.

<sup>(</sup>d): No data were reported from Bulgaria.



18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilattico

Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

**Table 48:** Reported confirmed human cases of brucellosis and notification rates per 100,000 population in EU MSs and non-MSs countries, by country and year, 2017–2021


			202	1			2020		20	19	20	18	2017	
Country	Sta	tus	National coverage <sup>(c)</sup>	Data		rmed s and tes	cases	rmed s and tes	case	irmed s and tes	Confi cases rat	and	cases	rmed s and tes
	Bv <sup>(a)</sup>	Ov, Cp <sup>(b)</sup>	coverage **	Tormat (7	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate
Austria	DFS	DFS	Υ	С	6	0.07	8	0.09	6	0.07	7	0.08	6	0.07
Belgium	DFS	DFS	Υ	Α	7	0.06	4	0.03	3	0.03	9	0.08	8	0.07
Bulgaria			Y	Α	0	0	1	0.01	0	0	1	0.01	2	0.03
Croatia	DFS		Υ	С	2	0.05	1	0.02	3	0.07	3	0.07	1	0.02
Cyprus	DFS	DFS	Υ	С	1	0.11	0	0.00	0	0	0	0	0	0
Czechia	DFS	DFS	Y	С	1	0.01	0	0.00	4	0.04	4	0.04	1	0.01
Denmark <sup>(d)</sup>	DFS	DFS	_	-	_	_	_	_	-	_	-	_	-	_
Estonia	DFS	DFS	Y	С	0	0	0	0.00	1	0.08	1	0.08	0	0
Finland	DFS	DFS	Y	С	0	0	0	0.00	0	0	0	0	1	0.02
France	DFS		Y	С	21	0.03	19	0.03	34	0.05	0	0	21	0.03
Germany	DFS	DFS	Y	С	13	0.02	19	0.02	36	0.04	37	0.04	41	-
Greece			Υ	С	24	0.22	30	0.28	65	0.61	97	0.90	94	0.87
Hungary		DFS	Y	С	0	0	0	0.00	0	0	0	0	0	0
Ireland	DFS	DFS	Υ	С	0	0	0	0.00	0	0	0	0	2	0.04
Italy			Y	С	32	0.05	18	0.03	49	0.08	94	0.16	99	0.16
Latvia	DFS	DFS	Υ	С	0	0	1	0.05	0	0	0	0	0	0
Lithuania	DFS	DFS	Y	С	0	0	0	0.00	0	0	0	0	0	0
Luxembourg	DFS	DFS	Υ	С	1	0.16	0	0.00	0	0	0	0	0	0
Malta	DFS		Y	С	0	0	0	0.00	0	0	0	0	0	0
Netherlands	DFS	DFS	Υ	С	2	0.01	2	0.01	7	0.04	5	0.03	2	0.01
Poland	DFS	DFS	Y	С	1	< 0.01	0	0.00	2	0.01	0	0	2	0.01
Portugal			Y	С	10	0.10	9	0.09	33	0.32	19	0.18	16	0.16
Romania	DFS	DFS	Y	С	0	0.00	0	0.00	1	0.01	1	0.01	3	0.02
Slovakia	DFS	DFS	Y	С	6	0.11	2	0.04	1	0.02	0	0	1	0.02
Slovenia	DFS	DFS	Y	С	0	0	1	0.05	6	0.29	3	0.15	1	0.05
Spain <sup>(e)</sup>		DFS	Y	С	25	_	10	_	20	0.04	40	0.09	63	0.14
Sweden	DFS	DFS	Y	С	10	0.10	7	0.07	14	0.14	11	0.11	14	0.14
EU Total 27		-	-	-	162	0.03	132	0.03	285	0.06	332	0.08	378	0.09
United Kingdom <sup>(f)</sup>	_	-	_	-	_	-	_	-	24	0.04	_	-	-	-
EU Total		_	_	-	162	0.03	132	0.03	309	0.06	332	0.08	378	0.09
Iceland			Y	С	0	0	0	0	0	0	0	0	0	0
Norway	DFS	DFS	Y	С	3	0.06	2	0.04	4	0.08	3	0.06	3	0.06
Liechtenstein	DFS	DFS	Y	С	0	0	3	0.03	7	0.08	5	0.06	9	0.11
Switzerland <sup>(g)</sup>	DFS	DFS	Υ	С	6	0.07								

<sup>-:</sup> Data not reported.

DFS: Disease-free status from infection with *B. abortus*, *B. melitensis* and *B. suis* in bovine animal populations, and/or in ovine and caprine animal populations.

- (a): Bovine animal population.
- (b): Ovine and caprine animal populations.
- (c): Y: yes; N: no; A: aggregated data; C: case-based data.
- (d): No surveillance system.
- (e): Data not complete for 2020–2021, rate not estimated.
- (f): Cases reported by the United Kingdom for the period 2017–2019 were also considered for this estimation (EU-28). When the United Kingdom data were collected for the period 2017–2019, the United Kingdom was an EU MS, but it became a third country on 1 February 2020.
- (g): Switzerland provided data directly to EFSA. The human data for Switzerland include data from Liechtenstein for the years 2017–2020.

Clear seasonality was observed in the number of confirmed brucellosis cases in the EU, with more cases reported from April to August. There was a significantly (p < 0.01) declining EU trend from 2017 to 2021 (Figure 16). Three MSs (Germany, Greece and Italy) had significantly decreasing 5-year trends from 2017 to 2021.



Source: Austria, Cyprus, Czechia, Estonia, Finland, France, Germany, Greece, Hungary, Ireland, Italy, Lithuania, Latvia, Malta, Netherlands, Poland, Portugal, Romania, Slovakia, Slovenia and Sweden.

Figure 16: Trends in reported confirmed human cases of brucellosis in the EU, by month, 2017–2021

Ten MSs provided information on hospitalisation. Out of 162 human cases, 60 (37.0%) were reported with information on hospitalisation. Among these, 36 (60.0%) were hospitalised. This is a decrease of 6.7% compared to the data for 2020 (64.3%). Eleven MSs provided information on the outcome. Out of the 162 cases of human brucellosis, 59 (36.4%) were reported with information on disease outcomes. No fatalities out of the 59 cases were notified.

Out of 162 human cases, 58 (35.8%) were reported with information on the *Brucella* species by 12 MSs. *B. melitensis* was reported as the aetiological agent in 51 cases (87.9%); *B. suis* was reported in three cases (5.2%); *B. abortus* in two cases (3.4%); and other (not specified) *Brucella* species in two cases (3.4%).

The proportion of *B. melitensis* infections was the same compared to the data in 2020 (87.8%). Only 11 (30.6%) out of 36 hospitalised cases were reported with information on the *Brucella* species. *B. melitensis* was reported as the aetiological agent in 10 hospitalised cases and *B. abortus* in 1 hospitalised case.


The number of confirmed, domestically acquired brucellosis cases in humans (patients not having been outside the country of notification during the disease incubation period) is overlaid with the national prevalence data on *Brucella*-positive cattle herds, and sheep and goat herds in the EU in 2021 in Figure 17. Greece, Italy and Spain have the highest number of confirmed brucellosis cases in

1831472, 2022, 12, Downloaded from https://efs.aonlinelibray.wiley.com/doi/10/2033/jefs.a2022/7666 by Istituto Zooporficiation Sperimenta dell'Umbria e delle Marche (ZSUM), Wiley Online Library on [18/12/2023], See the Terms and Conditions (http://onlinelibray.wiley.com/rems-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Cerawise Commons of the Common of the Common



dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

humans, while Greece, Italy and Portugal have the highest prevalence of *Brucella*-positive ruminant herds. Italy, which has reported a high number of human brucellosis cases over the years, did not report the origin of infection for 2021.



**Figure 17:** Number of confirmed, domestically acquired brucellosis cases in humans and national prevalence of *Brucella*-positive cattle herds, and sheep and goat herds, in EU MSs and non-MS countries, 2021

#### 6.3.3. Brucella in food

Very few monitoring data for *Brucella* were submitted in 2021, as was the case in previous years. In total, 307 samples of 'milk', 'cheese' and 'other dairy products' were collected from processing plants by three MSs (Italy, Portugal and Spain). Two Italian samples from 'raw sheep's milk' taken at the processing plant level identified as positive for an unspecified *Brucella* species.

#### 6.3.4. Brucella in animals

Monitoring data for *Brucella* from bovine animals, and from sheep and goats, originating from the National Control and Eradication Programmes and/or from countries or zones with disease-free status

#### Cattle

The status of countries for brucellosis in cattle, as of 31 December 2021, showed a favourable situation, with 21 MSs and the United Kingdom (Northern Ireland) being disease-free in 2021 (Figure 18). Croatia obtained disease-free status (DFS) in 2021. Out of the six non-disease-free status



onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico

dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms


and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative C

MSs, Italy, Portugal and Spain have increasing numbers of disease-free zones. Since 2019, Spain has reported no positive herds, indicating an extremely low level of prevalence, and absence of infection. Bulgaria, Greece and Hungary have no disease-free zones.

Three countries had disease-free zones or provinces in cattle:

- Italy: 13 regions and 9 provinces.
- Portugal: 1 region (Algarve) and 14 districts (from the Centre region and from the Azores islands).
- Spain: 16 autonomous communities and 1 province (Badajoz).

Lichtenstein, Norway, Switzerland and the United Kingdom (Northern Ireland) are disease-free in accordance with the EU legislation. Iceland, which has no special agreement on animal health (status) with the EU, has never reported any brucellosis cases caused by *B. abortus*, *B. melitensis* or *B. suis* (Figure 18).<sup>28</sup>



**Figure 18:** Status of countries as regards brucellosis in bovine animal population (*Bison* ssp., *Bos* ssp., *Bubalus* ssp.), MSs and non-MSs, 2021

In 2021, the overall proportion of cattle herds infected with *B. abortus, B. melitensis* or *B. suis* in the EU remained very low (0.04%; 554 out of 1,719,963 herds), with 22 MSs and the United Kingdom (Northern Ireland) reporting no cases of brucellosis in cattle. Greece, Italy and Portugal reported most infected herds.

In DFS MSs or in the DFS zones of non-DFS MSs, the overall prevalence was extremely low (< 0.001) (Table 49). Only eight infected cattle herds were reported in these DFS zones of the EU (six in 2020), with Italy reporting seven positive herds and France one.

\_

<sup>&</sup>lt;sup>28</sup> French, Spanish overseas territories are defined as DFS zones; Portuguese overseas territories are defined as non-DFS zones.



Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions

In 2021, the prevalence of brucellosis-infected cattle herds remained very low in the non-DFS zones of the six non-disease-free MSs, with 546 positive herds (0.43%; Table 49), compared with 603 (0.38%) positive herds in 2020. The number of positive herds out of all herds reported by Italy in its non-DFS zones was 453 (504 in 2020), while this number was 24 in Portugal (27 in 2020). Greece reported 69 positive herds (72 in 2020).

**Table 49:** Status of countries as regards brucellosis in bovine animals (*Bos* ssp., *Bubalus* ssp. and *Bison* ssp.) and related prevalence, EU, 2021

Member State (MS)	Disease-free status from brucellosis in bovine animals <sup>(a)</sup>	N of infected herds in DFS zones	Prevalence (%) of infected herds in DFS <sup>(a)</sup> zones	N of infected herds in non- DFS <sup>(a)</sup> zones	Prevalence (%) of infected herds in non-DFS <sup>(a)</sup> zones
Austria	DFS	0	0	_	_
Belgium	DFS	0	0	_	_
Bulgaria		_			0
Croatia	DFS	0	0	_	_
Cyprus	DFS	0	0	_	_
Czechia	DFS	0	0	_	_
Denmark	DFS	0	0	_	_
Estonia	DFS	0	0	_	_
Finland	DFS	0	0	_	_
France	DFS	1	< 0.01	_	_
Germany	DFS	0	0	_	_
Greece		_	_	69	0.63
Hungary		_	_	0	0
Ireland	DFS	0	0	_	_
Italy		7	0.01	453	1.3
Latvia	DFS	0	0	_	_
Lithuania	DFS	0	0	_	_
Luxembourg	DFS	0	0	_	_
Malta	DFS	0	0	_	_
Netherlands	DFS	0	0	_	_
Poland	DFS	0	0	_	_
Portugal		0	0	24	0.11
Romania	DFS	0	0	_	_
Slovakia	DFS	0	0	_	_
Slovenia	DFS	0	0	_	_
Spain		0	0	0	0
Sweden	DFS	0	0	_	_
United Kingdom (Northern Ireland) <sup>(b)</sup>	DFS	0	0	_	_
EU Total (27 + XI)		8	< 0.001	546	0.43

<sup>-:</sup> not applicable (no such zones).

DFS: Disease-free status.

127

DFS All Zones of the MS have disease-free status.

Not all Zones of the MS have disease-free status.

No Zones of the MS have disease-free status.

<sup>(</sup>a): Disease-free status from infection with B. abortus, B. melitensis and B. suis, in bovine animal populations.

<sup>(</sup>b): Data from the United Kingdom (Northern Ireland) are taken into account for 2021. In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/Northern Ireland, EU requirements on data sampling are also applicable to Northern Ireland, which is disease-free.



18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wley.com/doi/10.2903j.gfsa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/erms

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons

Comparing data for 2012 with data for 2021, the overall annual number of reported infected cattle herds in non-DFS zones decreased by 53.7%, from 1,181 to 546, while the prevalence of infected cattle herds increased from 0.10% to 0.43% (Figure 19). This is due to the reduction in the number of non-DFS zones and the resulting drop in the total number of cattle herds of interest from 1,162,978 to 127,000 during the same period. No speciation of *Brucella* isolates was reported for cattle.





Figure 19: Prevalence of Brucella-positive cattle herds, in non-DF zones, EU, 2012–2021

Figure 20 displays trends between 2005 and 2021 in the reported prevalence of brucellosis test-positive cattle herds in Greece, Italy and Portugal. The prevalence in Greece showed considerable annual variation from a minimum of 2% in 2008 to a maximum of 12% in 2012. The prevalence in Italy has remained under 2% since 2012, with a minimum value of 1.3% in 2019. The prevalence in Portugal has decreased consistently from about 1% in 2005 to 0.11% in 2021.

onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico

Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common



**Figure 20:** Prevalence of *Brucella*-positive cattle herds, in two MSs with non-DFS zones (Italy and Portugal) and in one non-DFS MS (Greece), 2005–2021

In 2021, bovine brucellosis was not detected in the following non-MS countries: Iceland, Montenegro, Norway and Switzerland. In the Balkans area, the disease is still present in cattle, with 25 infected (0.17%) herds out of 14,898 herds reported by North Macedonia, along with positive results reported by Bosnia and Herzegovina, and Serbia from national monitoring data. Bosnia and Herzegovina, and Serbia reported 108 positive animals out of 111,785 (0.10%) and 13 positive animals out of 488,587 (0.003%), respectively.

#### Sheep and goats

The status of countries as regards ovine and caprine brucellosis, reflecting the situation on 31 December 2021, is presented in Figure 21 and in Table 50. In 2021, 20 MSs and the United Kingdom (Northern Ireland) had disease-free status, with Spain obtaining the disease-free status in 2021. Out of the seven other MSs, three had DFS zones (France, Italy and Portugal) and four MSs had no zones with a disease-free status (Bulgaria, Croatia, Greece and Malta). In France, no cases of brucellosis have been reported in small ruminants since 2003, and the whole of metropolitan France obtained disease-free status in 2021.

Three countries had disease-free zones or provinces for sheep and goats:


- France: all 13 metropolitan regions.
- Italy: 15 regions and 5 provinces.
- Portugal: 1 region (the autonomous region of the Azores).

In 2021, the overall proportion of sheep and goat herds infected with *B. abortus, B. melitensis* or *B. suis* in the EU remained very low (0.03%; 346 out of 1,069,048 herds). In 2021, 23 MSs and the United Kingdom (Northern Ireland) reported no cases of infection with *B. abortus, B. melitensis* or *B. suis* in sheep and goat herds (Table 50). Infected or positive herds were reported by Greece, Italy, Portugal and Spain.



onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico

dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.c



**Figure 21:** Status of countries as regards ovine and caprine brucellosis, MSs<sup>29</sup> and non-MS countries, 2021

In particular, in the DFS zones the prevalence was extremely low, with 21 MSs reporting no cases of brucellosis, and Italy and Spain together reporting 15 infected herds (3 in 2020 reported by Italy), resulting in an overall prevalence in the DFS zones lower than 0.01% (0.0004% in 2020).

In 2021, the seven MSs with non-DFS zones reported 331 infected herds (0.18%), compared to 349 (0.22%) in 2020 (Table 50). The number of infected herds reported by these MSs was 26 in Greece (33 in 2020), 107 in Italy (120 in 2020) and 198 in Portugal (196 in 2020). No infected herds were reported by Bulgaria, Croatia, France and Malta. For the last 2 years, Croatia has reported zero infected herds, indicating that in the coming years, eradication of sheep and goat brucellosis is an achievable goal.

<sup>&</sup>lt;sup>29</sup> French, Portuguese and Spanish overseas territories are defined as non-DFS zones.



**Table 50:** Status of countries for brucellosis in sheep (*Ovis* ssp.) and goat (*Capra* ssp.) populations and related prevalence, EU, 2021

Member State	Disease-free status from brucellosis in sheep and goats <sup>(a)</sup>	N of infected herds in DFS zones	Prevalence (%) of infected herds in DFS <sup>(a)</sup> zones	N of infected herds in non DFS <sup>(a)</sup> zones	Prevalence (%) of infected herds in non-DFS <sup>(a)</sup> zones
Austria	DFS	0	0	_	_
Belgium	DFS	0	0	_	_
Bulgaria		_	_	0	0
Croatia		_	_	0	0
Cyprus	DFS	0	0	_	_
Czechia	DFS	0	0	_	_
Denmark	DFS	0	0	_	_
Estonia	DFS	0	0	_	_
Finland	DFS	0	0	_	_
France <sup>(b)</sup>		0	0	_	_
Germany	DFS	0	0	_	_
Greece		_	_	26	0.10
Hungary	DFS	0	0	_	_
Ireland	DFS	0	0	_	_
Italy		6	< 0.01	107	0.40
Latvia	DFS	0	0	_	_
Lithuania	DFS	0	0	_	_
Luxembourg	DFS	0	0	_	_
Malta		_	_	0	0
Netherlands	DFS	0	0	_	_
Poland	DFS	0	0	_	_
Portugal		0	0	198	0.38
Romania	DFS	0	0	_	_
Slovakia	DFS	0	0	_	_
Slovenia	DFS	0	0	_	_
Spain	DFS	9	< 0.01	_	_
Sweden	DFS	0	0	_	_
United Kingdom (Northern Ireland) <sup>(c)</sup>	DFS	0	0	-	-
EU Total (27 + XI)		15	< 0.01	331	0.18

<sup>-:</sup> not applicable (no such zones).

DFS: Disease-free status.

All Zones of the MS have disease-free status.

Not all Zones of the MS have disease-free status.

No Zones of the MS have disease-free status.

Spain reported that nine sheep and goat herds in its DFS zones were infected with *Brucella melitensis*. No other speciation of isolates was reported for sheep and goats.

18314732, 2022, 12, Downloaded from https://efs.aonlinelibrary.wiley.com/doi/10.2903/jefs.a.2022.7666 by Istituto Zopprofilative Sperimenta dell'Umbria e delle Marche (ISSUM), Wiley Online library.wiley.com/rems-and-conditions) on Wiley Online library for rules of use. O. A articles are governed by the applicable Creative Common

<sup>(</sup>a): Disease-free status from infection with B. abortus, B. melitensis and B. suis in ovine and caprine animal populations

<sup>(</sup>b): The whole of metropolitan France has disease-free status.

<sup>(</sup>c): Data from the United Kingdom (Northern Ireland) are taken into account for 2021. In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/Northern Ireland, EU requirements on data sampling are also applicable to Northern Ireland, which is disease-free.



18314732, 2022, 12, Downloaded from https://eds.onlinelibrary.wiley.com/doi/10.2903j.efsa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/emms

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons

From 2012 to 2021, the overall annual number of reported infected sheep and goat herds in the non-disease-free zones decreased by 80.4%, from 1,693 to 331, while the prevalence of positive herds decreased by 60%, from 0.45% to 0.18% (Figure 22).




Figure 22: Prevalence of Brucella-positive sheep and goat herds, in non-DFS zones, EU, 2012–2021


Figure 23 displays trends from 2004 to 2020 in the reported prevalence of brucellosis test-positive sheep and goat herds in non-DFS zones of the three MSs reporting the majority of infected herds in the EU. The prevalence in Greece showed high annual variation from a minimum of 0.4% in 2015 to a maximum of 8.6% in 2012. This may be related to the low proportion of tested herds, which leads to a lack of precision in estimates. Italy and Portugal reported low (> 1-10%) to very low (0.1–1%) prevalence during this period, decreasing for both MSs.

Brucellosis was not detected in sheep and goat herds in 2021 in the following non-MS countries: Iceland, Montenegro, Norway and Switzerland. North Macedonia reported 44 infected herds (0.79%) out of 5,574 and submitted national monitoring data on ovine and caprine brucellosis. From national monitoring data, North Macedonia and Serbia reported 1,406 positive animals out of 502,864 (0.28%) and 8 positive animals out of 1,406,330 (0.001%), respectively.

onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico

Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common



**Figure 23:** Prevalence of *Brucella*-positive sheep and goat herds, in two MSs with non-DFS zones (Italy and Portugal) and in one non-DFS MS (Greece), 2005–2021

#### Monitoring data for Brucella from animals other than bovine animals, and sheep and goats

Similarly to 2021 reports on *Brucella* from cattle, and from sheep and goats, *Brucella* species were reported from a wide range of animal species. *B. canis* was isolated in dogs from six MSs (Finland, France, Italy, the Netherlands, Sweden and Romania) and from one non-MS (Norway). Five MSs reported 44 positive dogs out of 414 tested animals (10.6%). *B. microti* was isolated from 'farmed' frogs in France; *B. suis* biovar 2 or unspecified *Brucella* species were reported in pigs, wild boars and wild hares in seven MSs (Finland, France, Germany, Italy, Romania, Slovakia and Spain); *B. melitensis* was isolated from Alpine ibex in France; unspecified *Brucella* species were found in wild bears, wild deer and wild boars; and *B. ceti* was found in dolphins in Italy and marine mammals in Sweden.

#### 6.4. Discussion

The 5-year EU trend of confirmed cases of brucellosis in humans declined significantly from 2017 to 2021. The EU notification rate was 0.03 per 100,000 population, which was the same as that reported in 2020, the lowest since surveillance began in the EU in 2007. According to Kefaloudi et al. (2022), human brucellosis remains a persistent public health problem with an annual notification rate averaging 0.9/100,000 population (standard deviation [SD]: 0.35).

Because international travel was significantly reduced during the first part of the COVID-19 pandemic, it has been postulated that the risk of human infection of travellers to endemic areas traditionally suited to tourism was likely smaller. This could have had an impact on the trend for human brucellosis. More evidence-based data should be acquired to provide insights into the true efficacy of the eradication plans in animals on the incidence of brucellosis in humans.

Like in 2020, human cases in non-DFS countries, such as Greece and Portugal, can be inferred to be mainly of domestic origin, while human cases in DFS countries, such as Austria, Germany, France and Sweden are of non-domestic origin. Italy, which also reported several human cases, did not provide information on the origin of infection. Interestingly, both Bulgaria (non-disease-free) and Hungary (non-disease-free for cattle) did not report any cases of human brucellosis. Where such information was available, it was clear that human cases were commonly associated with hospitalisation.

18314732, 2022, 12, Downloaded from https://fs.aonlinelibrary.wiley.com/doi/10.9293/j-fs.a.2022.7566 by Istituto Zoporofilatios Sperimenta dell'Umbria e delle Marche (ZSUM), Wiley Online Library on [1812023], See the Terms and Conditions (https://onlinelibrary.wiley.com/rerms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Cerawise Common.

In 2021, human cases of brucellosis were mainly caused by infection with *B. melitensis*. Furthermore, *B. melitensis* was the major species involved in cases of hospitalisation, where speciation information was provided. This information is very important when optimising risk management to further reduce disease in humans, considering that *B. melitensis* is mainly, if not completely, associated with brucellosis in sheep and goats. Three human cases were attributed to *B. suis*. This finding should be monitored in the next few years to detect a possible trend for a potentially emerging condition.

Bovine brucellosis, and ovine and caprine brucellosis have been eradicated by most EU MSs. In MSs and zones with disease-free status, very few infected herds were reported for 2021: 8 infected cattle herds, and 15 infected sheep and goat herds.

Eradication of brucellosis in cattle has been achieved in Croatia and Spain. Metropolitan France has eradicated brucellosis in sheep and goats. Some MSs were not disease-free free from bovine brucellosis and/or from brucellosis in sheep and goats, and both infections were mostly reported in Greece, Italy and Portugal. In Greece, the proportion of tested herds remained limited, which may affect the precision of surveillance. The overall number of infected herds should be interpreted differently between countries, as the proportion of tested herds was highly variable between MSs.

Nevertheless, from 2012 to 2021, the overall annual number of reported positive ruminants in the non-DFS zones decreased, with a decrease in prevalence. In Italy, problems were concentrated in the southern part of the country and Sicily. Moreover, the infection was present in water buffalo from the provinces of Salerno and Caserta, where in the latter province a drastic increase in the prevalence and incidence of infection in buffalo has been observed in the last years. About 80% of Campania's buffalo population is found in Caserta. The infection is concentrated in the geographic areas with the highest density of animals and farms per km² (Ottaiano et al., 2022). In non-DFS zones, non-food-borne transmission of brucellosis to humans may still occur through direct contact with infected animals. People working with farm animals, including farmers, livestock breeders, butchers, abattoir workers and veterinarians, are known to be at increased risk of brucellosis in endemic countries.

In 2021, the regulatory framework for brucellosis in animals changed substantially in the EU. Infection with *B. abortus*, *B. melitensis* and *B. suis* in bovine animals, and sheep and goats is currently considered a disease to be controlled in all MSs with the goal of eradicating the disease in these animals throughout the EU, while being kept under surveillance in other even-toed ungulates (Artiodactyla). All MSs must have a surveillance, or a control and eradication programme in place, approved by the EC. This should lead, in the coming years, to further progressive improvement in the already satisfactory epidemiological situation.

Attention should also be paid to canine brucellosis cases due to *B. canis*, which is also considered zoonotic and which is covered by the Directive 2003/99/EC, Annex I list A 'brucellosis and agents thereof'. In the last 2 years, an increased number of cases was described in Italy (De Massis et al., 2021), France, the Netherlands (van Dijk et al., 2021) and the United Kingdom (England; source APHA). Moreover, despite the lack of bacterial speciation, it is important to highlight that many different zoonotic species of *Brucella* are isolated from many hosts, including marine mammals and amphibians. Rouzic et al. (2021) reported the first human case of brucellosis caused by an isolate whose genome is identical to that of a frog isolate from Texas, demonstrating the zoonotic potential of atypical amphibian-type *Brucella*.

In conclusion, although cases of brucellosis in humans and ruminants are declining in the EU, brucellosis is still an animal health concern with public health relevance, especially in southern European countries that are not brucellosis disease-free.



18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilattico

dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

#### 7. Trichinella

Summary data substantiating this chapter, as well as additional information on related projects and internet sources are published for this report on the EFSA Knowledge Junction on Zenodo here. Summary statistics on trichinellosis foodborne outbreaks reported in the framework of Directive 2003/99/EC, with downloadable files, are retrievable using the EFSA foodborne outbreaks dashboard available here. Summary statistics on human surveillance data with downloadable files are retrievable using the ECDC Surveillance Atlas of Infectious Diseases here.

# **Trichinella** in the EU, 2021




### **Human cases in foodborne outbreaks**



### Foodborne outbreaks

Food vehicles causing strong-evidence outbreaks





#### 7.1. Key facts

- In 2021, the number of confirmed cases of human trichinellosis was 77, corresponding to an EU notification rate of 0.02 per 100,000 population. This was a decrease of 32.5% compared with 2020 (0.03 per 100,000 population).
- Compared with the rate before the COVID-19 pandemic (2017–2019 annual mean), the 2021 EU notification rate decreased by 17.5% and 28.6% with and without the data from the United Kingdom, respectively.
- The overall trend for trichinellosis during the period 2017–2021 in the EU did not show any statistically significant increase or decrease.
- In 2021, no infections with *Trichinella* were reported in tested fattening pigs (54 million) or breeding pigs (0.9 million) kept under controlled housing conditions, confirming that farming conditions are a key factor in preventing infection with this zoonosis.
- In pigs not kept under controlled housing conditions, 0.0001% (120 out of 161 million) were positive for *Trichinella*. Romania accounted for most of the positive pigs (81), followed by Poland (19), Spain (13), Croatia (five) and Finland and France (one each).
- No *Trichinella* infections were detected in domestic solipeds in the EU in 2021, as during the 2017–2020 period.
- In 2021, the proportion of hunted wild boar that tested positive was 0.07%.
- The proportion of *Trichinella*-positive foxes (indicator animals) was 1.6% in 2021, while for 2020, a rate of 0.9% was reported.

#### 7.2. Surveillance and monitoring of *Trichinella* in the EU

#### **7.2.1.** Humans

In 2021, 26 EU MSs reported information on trichinellosis in humans. The surveillance of *Trichinella* infections is mandatory in all reporting MSs, except in Belgium and France, where surveillance systems are voluntary. There is no surveillance system in place for trichinellosis in Denmark and the disease is not notifiable or reported at the EU level. The surveillance systems cover the whole population in all MSs reporting trichinellosis data. All countries reported case-based data except Bulgaria, which reported aggregated data. The EU case definition was used by 22 MSs, other case definitions were used by three MSs (France, Germany and Italy) and Belgium did not specify the case definition used.

#### **7.2.2.** Animals

# Trichinella monitoring data for domestic pigs (both fattening and breeding animals), farmed wild boar and solipeds

In accordance with Commission Implementing Regulation (EU) 2015/1375<sup>30</sup>, all *Trichinella*-susceptible animals intended for human consumption in the EU, i.e. domestic pigs (both fattening and breeding animals), farmed wild boar and solipeds, should be tested for the presence of *Trichinella* larvae in the muscles unless carcases have undergone a freezing treatment (freezing inactivates the parasite). ISO 18743/2015 or an equivalent method should be used (Commission Implementing Regulation (EU) 2015/1375). Therefore, data on *Trichinella* infections in these animals are comparable across MSs because the monitoring schemes are harmonised and the data collected and reported to EFSA originate from census sampling (EFSA BIOHAZ Panel, 2013a,b) (Table 51).

Domestic pigs, farmed and hunted wild boar and other wild animals (e.g. bears) that are not processed to be placed on the EU market (e.g. those intended for own consumption) are exempted from Commission Implementing Regulation (EU) 2015/1375 and their control falls under national legislation. Commission Implementing Regulation (EU) 2015/1375 states that the reporting of data for domestic pigs shall, at least, provide specific information related to the number of animals tested that were raised under controlled housing conditions (RCHC) as well as the number of breeding sows, boar and fattening pigs tested. Furthermore, the regulation states that a negligible risk status for a country or region is no longer recognised.

18314732, 2022, 12, Downloaded from https://fs.aonlinelibrary.wiley.com/inchibrary.wiley.com/inchibrary.wiley.com/em-and-conditions) on Wiley Online Library on [18/12/2023], See the Terms and Conditions (http://oinineibrary.wiley.com/em-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons

<sup>&</sup>lt;sup>30</sup> Commission Implementing Regulation (EU) 2015/1375 of 10 August 2015 laying down specific rules on official controls for *Trichinella* in meat (text with EEA relevance). OJ L 212, 11.8.2015, pp. 7–34.



18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico Sperimenta

dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

### Trichinella monitoring data for animals other than domestic pigs, farmed wild boar and solipeds

MSs should monitor the circulation of these nematodes in the main natural reservoir hosts (carnivorous and omnivorous animals) to acquire information on the risk of transmission to domestic animals (and from these to humans) and on the introduction of new *Trichinella* species from non-EU countries. However, monitoring data provided by the MSs to EFSA are generated by non-harmonised monitoring schemes across MSs without mandatory reporting requirements. Wild animals are the main reservoir hosts of *Trichinella*, and their biology and ecology vary from one MS to another and from one region or habitat in the same MS to another due to the human and environmental impact on ecosystems, resulting in different transmission patterns and prevalence rates of infection. Therefore, data on *Trichinella* in wild animals are not fully comparable among MSs, as neither harmonised monitoring schemes nor mandatory reporting requirements are in place, and the reported findings must be interpreted with caution. These data allow descriptive summaries to be produced at the EU level but preclude any subsequent data analysis such as an assessment of temporal and spatial trends (Table 51).

#### 7.3. Results

#### 7.3.1. Overview of key statistics, EU, 2017–2021

Table 51 summarises EU-level statistics on human trichinellosis and on *Trichinella* in animals, for the 2017–2021 period. More detailed descriptions of these statistics are provided in the below subsections and in the chapter on foodborne outbreaks.

**Table 51:** Summary of *Trichinella* statistics related to humans and main animal species, EU, 2017–2021

2021						
	2021 <sup>(a)</sup>	2020	2019 <sup>(b)</sup>	2018 <sup>(b)</sup>	2017 <sup>(b)</sup>	Data source
Humans						
Total number of confirmed cases	77	117	97	66	168	ECDC
Total number of confirmed cases/100,000 population (notification rates)	0.02	0.03	0.02	0.01	0.03	ECDC
Number of reporting MSs	26	26	27	27	27	ECDC
Infections acquired in the EU	29	99	26	18	81	ECDC
Infections acquired outside the EU	2	2	2	1	2	ECDC
Unknown travel status or unknown country of infection	46	16	69	47	85	ECDC
Number of outbreak- related cases	2	119	44	114	199	EFSA
Total number of outbreaks	1	6	5	10	11	EFSA
Animals						
Domestic pigs RCHC						
Number of units tested <sup>(c)</sup>	55,177,802	55,989,292	73,633,900	77,794,786	72,227,074	EFSA
% of positive units	0	0	0	0	0	EFSA
Number of reporting MSs	17	16	16	15	14	EFSA
Domestic pigs NRCHC						
Number of units tested <sup>(c)</sup>	161,129,635	139,637,631	145,213,445	152,922,322	124,689,434	EFSA
% of positive units	0.0001	0.0001	0.0002	0.0003	0.0002	EFSA
Number of reporting MSs	22	22	25	25	25	EFSA



18314732, 2022, 12, Downloaded from https://disa.onlinelibrary.wiley.com/doi/10.2903/j.disa.2022.7666 by Isituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.2903/j.disa.2022.7666 by Isituto Zooprofilation Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.2903/j.disa.2022.7666 by Isituto Zooprofilation Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.2903/j.disa.2022.7666 by Isituto Zooprofilation Sperimenta dell'Umbria e delle Marche (IZSUM). Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.2903/j.disa.2022.7666 by Isituto Zooprofilation Sperimenta dell'Umbria e dell'Um

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

	2021 <sup>(a)</sup>	2020	2019 <sup>(b)</sup>	2018 <sup>(b)</sup>	2017 <sup>(b)</sup>	Data source
Farmed wild boar						
Number of units tested <sup>(c)</sup>	5,755	3,922	7,570	6,343	17,799	EFSA
% of positive units	0	0	0	0	0.74	EFSA
Number of reporting MSs	8	6	7	7	8	EFSA
Hunted wild boar						
Number of units tested <sup>(c)</sup>	1,786,892	1,470,830	1,757,383	1,465,788	1,398,905	EFSA
% of positive units	0.0664	0.0484	0.0778	0.0891	0.0878	EFSA
Number of reporting MSs	20	21	23	23	22	EFSA
Foxes		,				
Number of units tested <sup>(c)</sup>	6,776	5,764	6,696	6,612	6,486	EFSA
% of positive units	1.5939	0.8501	1.3292	1.6334	1.2180	EFSA
Number of reporting MSs	11	9	10	10	11	EFSA

ECDC: European Centre for Disease Prevention and Control; EFSA: European Food Safety Authority; MSs: Member States. RCHC: raised under controlled housing conditions; NRCHC: not raised under controlled housing conditions.

#### 7.3.2. Human trichinellosis

In 2021, 77 confirmed cases of trichinellosis were reported by 26 MSs, which was a decrease compared with 2020 (Table 51). The EU notification rate decreased by 32.5% from 0.03 per 100,000 in 2020 to 0.02 per 100,000 in 2021. This decrease was mainly due to the decrease in the number of outbreak-related cases reported by Italy, who reported for 2020 a large outbreak. Fifteen countries, including Italy, reported zero cases in 2021. Compared with the rate before the COVID-19 pandemic (2017–2019 annual mean), there were decreases of 17.5% and 28.6% with and without the data from the United Kingdom, respectively.

In 2021, Bulgaria and Croatia had the highest notification rates in the EU (0.42 cases per 100,000 in both countries), followed by Latvia (0.37 cases per 100,000) and Austria (0.11 cases per 100,000) (Table 52). Together, these four countries (Austria, Bulgaria, Croatia and Latvia) accounted for 80% of all confirmed trichinellosis cases reported at the EU level in 2021.

In 2021, 29 cases (36.7%) of trichinellosis with known travel status and with known country of infection were reported to be acquired in the EU. One MS (Austria) reported two travel-associated trichinellosis cases infected outside the EU. For 46 cases (59.7%), travel information was not reported (Table 51).

**Table 52:** Reported confirmed human cases of trichinellosis and notification rates per 100,000 population in EU MSs and non-MS countries, by country and year, 2017–2021

	2021						2020 2019		2018		2017			
Country	National coverage <sup>(a)</sup>	Data format <sup>(a)</sup>	Confirmed cases and rates		cases and				Confirmed cases and rates		Confirmed cases and rates		Confirmed cases and rates	
			Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate		
Austria	Y	С	10	0.11	6	0.07	1	0.01	2	0.02	3	0.03		
Belgium <sup>(b)</sup>	Υ	Α	0	_	_	_	_	_	0	_	0	_		
Bulgaria	Υ	Α	29	0.42	13	0.19	55	0.79	45	0.64	55	0.77		
Croatia	Υ	С	17	0.42	0	0	3	0.07	0	0	21	0.51		
Cyprus	Υ	С	0	0	0	0	0	0	0	0	0	0		
Czechia	Υ	С	0	0	0	0	0	0	0	0	0	0		

<sup>(</sup>a): Data on animals from the United Kingdom (Northern Ireland) are taken into account for 2021. In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/Northern Ireland, the EU requirements on data sampling are also applicable to Northern Ireland.

<sup>(</sup>b): Data from the United Kingdom are taken into account for 2017–2019, because the United Kingdom was an EU MS, but it became a third country on 1 February 2020.

<sup>(</sup>c): Units: animals and/or slaughter animal batches.



		2021			20	20	20	19	20	18	2017	
Country	National coverage <sup>(a)</sup>	Data format <sup>(a)</sup>	cases	rmed s and tes	Confi cases		cases	rmed s and tes	case	rmed s and tes	case	firmed es and ates
			Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate
Denmark <sup>(c)</sup>	_	_	_	_	_	_	_	_	_	_	_	_
Estonia	Υ	С	0	0	0	0	0	0	0	0	0	0
Finland	Υ	С	0	0	0	0	0	0	0	0	0	0
France	Υ	С	2	< 0.01	1	< 0.01	2	< 0.01	0	0	8	0.01
Germany	Y	С	2	< 0.01	1	< 0.01	3	< 0.01	0	0	2	< 0.01
Greece	Υ	С	0	0	0	0	0	0	0	0	1	0.01
Hungary	Υ	С	0	0	0	0	0	0	2	0.02	0	0
Ireland	Υ	С	0	0	0	0	0	0	0	0	0	0
Italy	Y	С	0	0	79	0.13	10	0.02	2	< 0.01	4	0.01
Latvia	Υ	С	7	0.37	1	0.05	1	0.05	1	0.05	1	0.05
Lithuania	Y	С	1	0.04	0	0	0	0	0	0	9	0.32
Luxembourg	Υ	С	0	0	0	0	0	0	0	0	0	0
Malta	Y	С	0	0	0	0	0	0	0	0	0	0
Netherlands	Υ	Α	0	0	0	0	1	0.01	0	0	0	0
Poland	Y	С	2	0.01	11	0.03	2	0.01	2	0.01	9	0.02
Portugal	Y	С	0	0	0	0	1	0.01	0	0	1	0.01
Romania	Y	С	6	0.03	4	0.02	6	0.03	10	0.05	48	0.24
Slovakia	Y	С	0	0	0	0	0	0	0	0	1	0.02
Slovenia	Y	С	0	0	0	0	0	0	0	0	0	0
Spain	Y	С	1	< 0.01	1	< 0.01	12	0.03	2	< 0.01	5	0.01
Sweden	Y	С	0	0	0	0	0	0	0	0	0	0
EU Total 27			77	0.02	117	0.03	97	0.02	66	0.02	168	0.04
United Kingdom	Y	С	-	_	_	_	0	0	0	0	0	0
EU Total <sup>(d)</sup>			77	0.02	117	0.03	97	0.02	66	0.01	168	0.03
Iceland	Υ	С	0	0	0	0	0	0	0	0	0	0
Norway	Y	С	0	0	0	0	0	0	0	0	0	0
Liechtenstein	Υ	С	0	0	4	0.05	3	0.03	0	0	1	0.01
Switzerland <sup>(e)</sup>	Y	С	0	0								

<sup>-:</sup> Data not reported.

The EU trend in confirmed cases of trichinellosis did not show any statistically significant decrease or increase over the period 2017–2021 (Figure 24). During the same period, only Romania reported a significant decreasing trend and none of the MSs observed a significant increasing trend. Bulgaria, which had reported most of the cases until 2019 and had the highest notification rates in the EU in 2017–2021, was not included in the EU trend calculations since monthly data were not available.

Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.c

<sup>(</sup>a): Y: yes; N: no; A: aggregated data; C: case-based data.

<sup>(</sup>b): Sentinel surveillance, disease not under formal surveillance. Notification rate not calculated.


<sup>(</sup>c): No surveillance system.

<sup>(</sup>d): Cases reported by the United Kingdom for the period 2017–2019 were also considered for this estimation (EU-28). When the United Kingdom data were collected for the period 2017–2019, the United Kingdom was an EU MS, but it became a third country on 1 February 2020.

<sup>(</sup>e): Switzerland provided data directly to EFSA. The human data for Switzerland include data from Liechtenstein for the years 2017–2020.

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilattico

Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [1812/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common.



Source: Austria, Cyprus, Czechia, Estonia, Finland, France, Germany, Greece, Hungary, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Poland, Portugal, Romania, Slovakia, Slovenia, Spain and Sweden.

Figure 24: Trend in reported confirmed human cases of trichinellosis in the EU by month, 2017–2021

Of the 10 MSs reporting confirmed cases for 2021, six provided information on hospitalisation (26 cases, 33.8% of all confirmed cases reported in the EU). Of these cases, 10 (38.5%) were hospitalised, which was a decrease compared with 2020 (72.7%). Six MSs provided information on the outcome of the cases (28 cases, 35.4% of all confirmed cases); no deaths were reported. *Trichinella spiralis* was identified as the causative agent of all the 27 (34.2%) confirmed human cases for which species identification was reported by five MSs.

#### 7.3.3. Trichinella infection in food and animals

In 2021, one MS (Croatia) reported monitoring data for four meat products (fermented sausages made with pork) samples taken at the farm level and all tested positive. *T. spiralis* was identified in three samples.

In 2021, 31 countries (27 MSs, the United Kingdom (Northern Ireland) and three non-MSs) provided information on *Trichinella* in domestic animals (pigs and/or farmed wild boar). Sixteen MSs, the United Kingdom (Northern Ireland) and one non-MS (Iceland) reported data on breeding and/or fattening pigs RCHC; no positive findings were reported in these domestic pigs RCHC. No positive findings were found in farmed wild boar (Table 53).



**Table 53:** *Trichinella* monitoring results for domestic pigs and farmed wild boar in reporting EU MSs and non-MS countries, by housing conditions<sup>(a)</sup>, 2021

		Νp	ositive/tested (% pos	sitive)	
Country	Not co	ontrolled housing co	nditions (NCHC)	Controlled hous	ing conditions
Country	Farmed wild boar	Fattening pigs	Breeding pigs	Fattening pigs	Breeding pigs
Austria	0/80 (0)	0/5,024,563 (0)	0/90,865 (0)	_	_
Belgium	_	0/498,359 (0)	0/1,598,668 (0)	0/8,213,983 (0)	_
Bulgaria	_	_	-	0/98,945 (0)	-
Croatia	_	5/152,512 (< 0.001)	_	0/1,045,327 (0)	0/70,710 (0) <sup>(j)</sup>
Cyprus	_	0/572,556 (0)	0/13,059 (0)	_	_
Czechia	_	_	_	0/2,385,961 (0)	_
Denmark	0/394 (0)	0/771,963 (0)	0/271,526 (0)	0/17,164,861 (0)	0/289,868 (0)
Estonia	_	0/149,163 (0)	. , , ,	0/224,415 (0)	_
Finland	0/198 (0)		0/34,690 (0)	0/2,165 (0)	0/108 (0)
France		1/524,576 (< 0.001) <sup>(b)</sup>		0/25,734 (0)	0/308,567 (0)
Germany	0/1,157 (0)		-	-	_
Greece	0/1,323 (0)		0/20,981 (0)	0/7,899 (0)	_
Hungary	_	0/4,973,495 (0)	0/363,435 (0)	-	_
Ireland	_	-	-	0/3,603,287 (0)	0/90,444 (0)
Italy	0/1,773 (0)	0/9,776,660 (0) <sup>(d)</sup>	_	0/37,931 (0)	0/176,313 (0)
Latvia	_	0/473,831 (0)	_	-	-
Lithuania	_	-	_	0/901,388 (0)	_
Luxembourg	_	0/144,880 (0)	_	-	_
Malta	_	0/52,797 (0)	0/1,244 (0)		_
Netherlands	_	0/32/131 (0)	0/1/211(0)	0/14,955,097 (0)	_
Poland	_	19/21,438,846	_	0/14,955,097 (0)	_
Tolaria	_	(< 0.001)	_	_	_
Portugal	_	0/4,072,097 (0)	0/29,033 (0)	_	-
Romania	_	81/211,260 (< 0.001)	_	0/3,731,300 (0)	0/9,323 (0)
Slovakia	_	0/662,881 (0) <sup>(e)</sup>	0/13,373 (0)	_	_
Slovenia	-	0/242,584 (0) <sup>(f)</sup>	_	_	-
Spain	0/375 (0)	10/52,803,353 (< 0.001) <sup>(g)</sup>	3/818,668 (< 0.001) <sup>(h)</sup>	0/50,377 (0)	
Sweden	_	0/451,423 (0)	0/18,870 (0)	0/1,746,698 (0)	0/35,819 (0)
United Kingdom (Northern Ireland)	_	-	-	0/1,282 (0) <sup>(1)</sup>	-
EU Total (27 + XI)	0/5,755 (0)	117/157,840,843 (< 0.001)	3/3,288,792 (< 0.001)	0/54,196,650 (0)	0/981,152 (0)
Iceland	_			0/78,267 (0)	
Norway	_	0/1,568,000 (0)	_	_	_
Switzerland	_	0/2,265,215 (0)	0/29,836 (0)	_	_
Total non-EU Countries	-	0/3,833,215 (0)	0/29,836 (0)	0/78,267 (0)	_
Total EU (27+ XI) + non- EU countries	0/5,755 (0)	117/161,674,058 (< 0.001)	3/3,318,628 (< 0.001)	0/54,274,917 (0)	0/981,152 (0)

MSs: Member States.

1831/472, 2022, 12. Downloaded from https://dx.oonlinelibrary.wiley.com/dx/10.2903j.efxa.202.72666 by Isitiuto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18122023]. See the Terms and Conditions (https://aninebhary.wiley.com/rems-and-conditions) on Wiley Online Library for rules of use; OA attacks are governed by the applicable Creative Commons.

<sup>(</sup>a): No pigs reported in 2021 with unspecified housing conditions.

<sup>(</sup>b): Comprising 21,832 pigs in mixed herds including one positive pig.



18314732, 2022, 12, Downloaded from https://dfsa.onlinelibrary.wiley.com/doi/10.2903/j.cfsa.2022.7666 by Istituo Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM). Wiley Online Library on [18/12/2023]. See the Terms


and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

- (c): Comprising 3,104 pigs for own consumption.
- (d): Comprising 60,509 pigs for own consumption.
- (e): Comprising 1,020 pigs for own consumption.
- (f): Pigs in mixed herds.
- (g): Comprising 9,972 pigs for own consumption and 37,066 piglets.
- (h): Comprising 35,220 piglets.
- (i): Pigs in mixed herds.
- (j): Comprising 62,393 piglets and 747 boars.

Six MSs (Croatia, Finland, France, Poland, Romania and Spain) reported positive findings for domestic pigs NRCHC for 2021. For fattening pigs NRCHC, 117 (< 0.01%) (including one pig from a mixed herd) were positive, whereas only three breeding pigs were positive (< 0.01%). Romania accounted for more than half of the positive pigs (81, 67%), followed by Poland (19, 16%), Spain (13, 11%), Croatia (five, 4.2%), Finland (one, 0.8%) and France (one, 0.8%). Species identification was reported for 60 (50%) out of 120 pigs. *T. spiralis* was detected in 50 pigs (83%), *T. britovi* in eight pigs (13%), *T. pseudospiralis* in one pig (1.6%) and *T. nativa* in one pig (1.6%). As in previous years, these *Trichinella* infections were found in free-range and backyard pigs reared in rural EU regions.

As shown in Figure 25, from 2012 to 2018 (7-year period), *Trichinella* spp. were not reported in domestic pigs (RCHC or NRCHC) or farmed wild boar in 16 MSs, unlike in the other 12 MSs. In 2019, 2020 and 2021, *Trichinella* spp. in pigs were reported by six (Bulgaria, Croatia, France, Poland, Romania and Spain), seven (Bulgaria, Croatia, France, Greece, Italy, Romania and Spain) and six (Croatia, Finland, France, Poland, Romania and Spain) MSs, respectively.





These distribution maps have been built based on data from reports (EFSA and ECDC (European Food Safety Authority and European Centre for Disease Prevention Control), 2015a,b, 2016, 2017). In 2021, Bulgaria did not report data on pigs NRCHC.

**Figure 25:** *Trichinella* spp. in domestic pigs and farmed wild boar, in EU MSs and non-MS countries, 2012–2021

In 2021, as in the previous 5-year period (2016–2020), positive findings were not reported either in 103,745 domestic solipeds tested in 20 MSs or in 8,276 domestic solipeds tested in two non-MSs (Iceland and Switzerland) (Table 54).

**Table 54:** *Trichinella* monitoring results for hunted wild boar and wild boar with unspecified habitat, bears, foxes and domestic solipeds, in reporting EU MSs and non-MS countries, 2021

Country	N positive/tested (% positive)				
	Hunted or unspecified wild boar	Bears	Foxes	Domestic solipeds	
Austria	0/26,524 (0)	-	0/3 (0)	0/440 (0)	
Belgium	0/22,305 (0)	_	_	0/7,937 (0)	
Croatia	74/56,661 (0.13)	1/108 (0.93)	_	0/579 (0)	
Cyprus	_	_	0/75 (0)	_	
Czechia	0/232,355 (0)	-	3/2,758 (0.11)	0/77 (0)	

1831/372, 2022, 12, Downloaded from https://efas.onlinelibrary.wiley.com/doi/10.2903j.efas.2022.7666by Isituo Looprofilatico Sperimenta dell'Umbria e dello Marche (IZSUM), Wiley Online Library on [18/12/203]. See the Terms and Conditions (https://onlinelibrary.wiley.com/emm-and-conditions) on Wiley Online Library for rules of use; O. Aarticles are governed by the applicable Creative Common



dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.c

	N positive/tested (% positive)				
Country	Hunted or unspecified wild boar	Bears	Foxes	Domestic solipeds	
Denmark	_	_	_	0/761 (0)	
Estonia	3/1,705 (0.18)	9/68 (13.2)		0/8 (0)	
Finland	1/931 (0.11)	11/279 (3.9)	77/161 (47.8)	0/757 (0)	
France	2/51,511 (0)	_	_	0/3,815 (0)	
Germany	40/590,369 (0.01)	_	0/732 (0)	0/3,744 (0)	
Hungary	3/48,791 (0.01)	_	_	0/223 (0)	
Ireland	_	_	_	0/1,662 (0)	
Italy	14/208,148 (0.01)	0/1 (0)	8/2,316 (0.35)	0/28,800 (0)	
Latvia	16/7,641 (0.21)	_		0/34 (0)	
Luxembourg	0/6,747 (0)	_	0/116 (0)	0/10 (0)	
Netherlands	0/5,726 (0)	_		0/1,777 (0)	
Poland	453/201,263 (0.23)	_	16/85 (18.8)		
Portugal	0/121 (0)	_	_	0/649 (0)	
Romania	46/9,207 (0.50)	1/11 (9.1)	_	0/29,946 (0)	
Slovakia	0/13,072 (0)	0/3 (0)	3/93 (3.2)		
Slovenia	0/2,274 (0)	0/86 (0)	_	0/1,057 (0)	
Spain	531/170,715 (0.31)	_	_	0/20,361 (0)	
Sweden	3/130,826 (0)	0/325 (0)	1/41 (2.4)	0/1,108 (0)	
United Kingdom (Northern Ireland)	-	-	0/396 (0)	-	
EU Total (27 + XI)	1,186/1,786,892 (0.07)	22/881 (2.5)	108/6,776 (1.6)	0/103,745 (0)	
Iceland	_	_	_	0/7,158 (0)	
Norway	0/292 (0)	_	_	_	
Republic of North Macedonia	14/1,218 (1.1)	-	_	_	
Switzerland	1/10,741 (0.01)	0/1 (0)		0/1,118 (0)	
Total non-EU Countries	15/12,251 (0.12)	0/1 (0)	_	0/8,276 (0)	
Total EU (27+ XI) + non-EU countries	1,201/1,799,143 (0.07)	22/882 (2.5)	_	0/112,021 (0)	

Twelve MSs reported positive findings for hunted wild boar (1,186 positive findings out of 1,786,892 animals tested (0.07%)). Species identification was provided for 865 wild boar (73%), of which 722 (83.4%) were infected with *T. spiralis*, 140 (16.2%) with *T. britovi*, two (0.23%) with *T. pseudospiralis* and one (0.11%) with *T. nativa* and *T. britovi* (mixed infection). For 321 (27%) animals, species identification was not reported.

Six MSs (Czechia, Finland, Italy, Poland, Slovakia and Sweden) reported positive findings for *Trichinella* in foxes (*Vulpes vulpes*) with, in total, 108 (1.6%) positive out of 6,776 tested animals in 11 MSs. *T. britovi* was identified in 26 (76%) animals, *T. spiralis* in four (12%), *T. nativa* in three (8.8%) and *T. pseudospiralis* in one (2.9%), out of 34 positive foxes with species identification provided. Four MSs (Croatia, Estonia, Finland and Romania) reported positive findings for *Trichinella* in brown bears (*Ursus arctos*) with 22 (2.5%) positive units out of 881 tested in eight MSs (Table 54). *T. nativa*, *T. britovi* and *T. pseudospiralis* were identified in 13 (59%), eight (36%) and one (4.5%) brown bear(s), respectively; one of them presented with a mixed *T. nativa* and *T. britovi* infection. Nine MSs and one non-MS reported data on *Trichinella* in wild animals other than foxes, brown bears and wild boar. Positive findings were detected for raccoon dogs (32.9%), lynxes (17.2%), wolves (16.8%), birds (0.72%), martens (0.44%), badgers (0.15%) and other animals (0.4%), as shown in Table 55. In wild animals, *T. spiralis* was reported in Poland (one beaver); *T. britovi* was identified in three MSs (Italy (18 wolves), Poland (one wolf) and Sweden (two lynxes)) and in one non-MS (Switzerland (one jackal)). *T. nativa* was documented in Finland (12 racoon dogs, one beaver and four lynxes) and



18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico

Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

Sweden (four lynxes and one wolf). *T. pseudospiralis* was documented in Finland (one goshawk and one owl) and Italy (one marsh harrier).

**Table 55:** *Trichinella* monitoring results in other wild animals in reporting EU MSs and non-MS countries, 2021

•			N positiv	e/tested (%	positive)		
Country	Badgers	Wolves	Raccoon dogs	Birds	Lynxes	Martens	Other animals
Austria	0/64 (0)						0/4 (0) <sup>(j)</sup>
Croatia	0/2 (0)						0/4 (0) <sup>(j)</sup>
Cyprus							0/1 (0) <sup>(k)</sup>
Finland	1/11 (9.1) <sup>(a)</sup>	19/41 (46.3) <sup>(b)</sup>	91/231 (39.4) <sup>(e)</sup>	2/71 (2.8) <sup>(f)</sup>	20/40 (50.0) <sup>(h)</sup>	1/2 (50.0) <sup>(b)</sup>	1/51 (1.9) <sup>(1)</sup>
Germany	0/56 (0)					0/46 (0)	0/331 (0) <sup>(m)</sup>
Hungary							0/3 (0) <sup>(n)</sup>
Italy	0/521 (0)	40/279 (14.3) <sup>(c)</sup>	0/46 (0)	1/344 (0.29) <sup>(g)</sup>		0/179 (0)	0/89 (0) <sup>(o)</sup>
Poland		1/1 (100) <sup>(d)</sup>					1/1 (100) <sup>(p)</sup>
Sweden	0/9 (0)	1/42 (2.4) <sup>(a)</sup>			6/111 (5.4) <sup>(i)</sup>		0/15 (0) <sup>(q)</sup>
EU Total (27 + XI)	1/663 (0.15)		91/277 (32.9)	3/415 (0.72)	26/151 (17.2)	1/227 (0.44)	2/499 (0.40)
Switzerland	0/9 (0)	2/22 (9.1) <sup>(d)</sup>			1/34 (2.9) <sup>(d)</sup>		1/4 (25.0) <sup>(r)</sup>
Total EU (27+ XI) + non-EU countries	1/672 (0.15)	63/385 (16.4)	91/277 (32.9)	3/415 (0.72)	27/185 (14.6)	1/227 (0.44)	3/503 (0.59)

- (a): T. nativa.
- (b): Unknown species.
- (c): Eighteen infected with *T. britovi*.
- (d): T. britovi.
- (e): Twelve infected with *T. nativa*.
- (f): Comprising three crows, seven eagles, 21 white-tailed eagles, one falcon, 30 goshawks (one positive for *T. pseudospiralis*) and nine owls (one positive for *T. pseudospiralis*).
- (g): Comprising one marsh harrier (positive for *T. pseudospiralis*) and 343 unspecified birds.
- (h): Four positive for *T. nativa*.
- (i): Four positive for *T. nativa* and two for *T. britovi*.
- (j): All deer.
- (k): Bats.
- (I): Comprising four minks, 45 otters and two wolverines (one positive for unknown species).
- (m): Comprising 12 coypus, 181 racoons and 138 unspecified animals.
- (n): Rats.
- (o): Comprising five roe deer, 19 hedgehogs, 33 jackals, three polecats, five weasels and 24 wildcats (Felis silvestris).
- (p): One T. spiralis-infected beaver.
- (q): Comprising six beavers, two lions, four seals and three wolverines.
- (r): Comprising one coypu, one pet dog, one jackal (positive for *T. britovi*) and one wildcat (*Felis silvestris*).

### 7.4. Discussion

Trichinellosis is a rare but serious human disease that is still present in some EU MSs. Fifteen out of 26 MSs reported zero cases including three MSs (Cyprus, Luxembourg and Malta) that have never reported any trichinellosis cases since the beginning of EU-level surveillance in 2007.

The COVID-19 pandemic did not appear to impact the reporting of human cases of trichinellosis in 2020 and 2021, whereas the number of reported FBOs and outbreak-related human cases dropped in 2021. One MS (Portugal) reported a decrease in the number of wild boar hunted in 2020 and 2021 compared with that in the previous years (2017–2010), due to lockdowns associated with this pandemic.

In general, *Trichinella* infections in humans are often linked to FBOs; therefore, the EU trend for trichinellosis has been affected by the number and size of FBOs. The EU notification rate was not higher than 0.03 per 100,000 population in the last 5 years, from 2017 to 2021, with the highest rate



(0.03) reported in 2017 and 2020, and the lowest rate (0.01) reported in 2018; this was the lowest rate ever reported since the beginning of EU-level trichinellosis surveillance in 2007. In 2021, Bulgaria and Croatia accounted for more than half (58%) of all confirmed cases in the EU and did not report any trichinellosis FBOs. Romania, which had experienced the most *Trichinella* outbreaks in the previous years, showed a significant decrease in the 5-year trend from 2017 to 2021. In 2021, only one weak-evidence outbreak was reported in the EU, by France.

Pigs represent the largest livestock category in the EU. More than 216 million pigs were tested for Trichinella in MSs and non-MSs in 2021 out of more than 246 million pigs reared in the EU (Marguer et al., 2014; European Parliament, 2020), with only 120 positive animals, i.e. about 0.49 per million reared pigs. Only six out of 22 MSs reported Trichinella in pigs in 2021, with an overall prevalence rate of 0.00005%. All positive findings were for pigs NRCHC. Most pigs at risk for this infection are backyard or free-range pigs, which are usually slaughtered at home, where veterinary control or recording can be easily evaded. In the EU, infected pigs are usually clustered in five MSs (Bulgaria, Croatia, Poland, Romania and Spain) and sporadic infections are documented in other MSs (Pozio, 2014). Bulgaria did not report any 2021 data for pigs NRCHC and reported zero positive out of all the tested pigs RCHC, confirming that RCHC is a key condition to prevent infection with this zoonosis. Finland reported one outdoor-reared T. nativa-positive Mangalica pig; this MS had not reported any positive findings in domestic pigs since 2012, although Trichinella spp. circulate among wild animals in this country. EFSA has identified that not raising domestic pigs under controlled housing conditions is a major risk factor for Trichinella infections (EFSA and ECDC, 2011; EFSA BIOHAZ Panel et al., 2011). Identification of Trichinella larvae at species level in 2021 confirmed that T. spiralis was more prevalent than T. britovi in pigs (83% vs. 13%) (Pozio et al., 2009). However, T. spiralis was patchily distributed. T. pseudospiralis was reported in only one out of 60 pigs (1.8%) for which the Trichinella species was available; these data confirmed the low prevalence of this species in target animals (Pozio, 2016).

Hunted wild boar are a major source of trichinellosis infections in humans. Human behaviour can strongly influence sylvatic cycles, both favouring and reducing the transmission of *Trichinella* spp. For example, carcases of *Trichinella*-infected animals left by hunters, or of such animals that have died in road accidents, may be scavenged by other wild animals, thereby contributing to transmission.

No positive findings were reported for solipeds in 2021. Over the last 12 years, only four horses tested positive out of more than one million tested, in 2008, 2010 and 2012 (EFSA and ECDC, 2009, 2010, 2011, 2012, 2013, 2014). This extremely low (< 0.001%) prevalence may have been related to effective control which, according to the EFSA BIOHAZ Panel (2013b), should be maintained if there is no full and reliable traceability system in place, especially since meat from solipeds can be eaten raw in some EU countries.

Trichinella spp. circulate among wild animals in large parts of Europe. The reporting of negative findings in MSs could be explained by an insufficient number of surveys, inadequate sample sizes or investigations in regions whose environmental conditions do not favour the transmission of these zoonotic nematodes among wildlife. Some MSs (Austria, Croatia and Italy) tested non-susceptible wild animals (deer) for Trichinella infection. Apart from horses, other herbivores are considered accidental hosts of Trichinella spp. (Kärssin et al., 2021). High endemicity for Trichinella in the sylvatic cycles of these countries and/or convenience sampling might justify surveillance in these animals. Foxes (indicator animals), having a large and widespread population, can be considered as the main natural reservoir of Trichinella in Europe. In 2021, the prevalence of Trichinella infection in this animal species was 1.59%, which was the highest level observed in the last 3 years. This increase was mainly due to the positive findings reported by Poland, which had not reported data on this animal species in 2020, and Finland. In 2021, this MS reported 77 positive animals out of 161 tested (47.8%), whereas in 2020, it had documented 35 positive findings out of 210 (16.7%). In this case, the prevalence of Trichinella infection in animals may vary among the different sampling areas. In 2021, as in 2020, the proportion of positive samples from wildlife was higher in raccoon dogs (32.9%), wolves (16.8%) and lynxes (17.2%) than in other animals sampled but their population sizes and distributions in Europe are generally limited to a few countries.

In some MSs (Bulgaria, Croatia, etc.), there has been an increasing number of pigs RCHC and increased control at slaughter of pigs NRCHC during the last few years. These measures, in combination with trichinellosis awareness-raising and farmers' education activities, may have contributed to a reduction in the parasite biomass in domestic habitats and in the probability of acquiring an infection for humans.



18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilattico

Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

Farming practices at risk of transmitting *Trichinella* spp. (rearing backyard or free-range pigs) occur, in general, in disadvantaged and poor areas where veterinary services do not exist or are unable to control many small pig units, or where veterinary supervision can be circumvented (Pozio, 2014). There are examples from the past, where countries had suitable controls in place for parasite management in domestic pigs, but where changes in pork production affected by socioeconomic conflicts resulted in the re-emergence of trichinellosis as a serious public health problem (Djordjevic et al., 2003; Cuperlovic et al., 2005). The increasing number of wild boar and foxes and the spread of the raccoon dog population from eastern to western Europe and of the jackal population from southern–eastern to northern–western Europe may increase the prevalence of *Trichinella* circulating among wild animals (Alban et al., 2011; Széll et al., 2013).



e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

### 8. Echinococcus

Summary data substantiating this chapter, as well as additional information on related projects and internet sources are published for this report on the EFSA Knowledge Junction on Zenodo here. Summary statistics on human surveillance data with downloadable files are retrievable using the ECDC Surveillance Atlas of Infectious Diseases here.

# **Echinococcus** in the EU, 2021

Human cases	Notification rate (per 100,000 population)	0.15	<b>Trend</b> (2017-2021)	Increasing Decreasing Stable
<b>529</b> Cases of illness				
128 Infections acquired in the EU		51	Hospitalisations	
81 Infections acquired outside the EU		0	Deaths	
320 Unknown travel status or unknown country of infect	ion			

# **Human cases in foodborne outbreaks**



# 8.1. Key facts

- In 2021, the number of confirmed cases of human echinococcosis was 529, corresponding to an EU notification rate of 0.15 per 100,000 population. This is a decrease of 7.5% compared with 2020 (0.16 per 100,000 population). The notification rate in 2021 was the lowest since EU surveillance of echinococcosis began in 2007.
- Compared with the rate before the COVID-19 pandemic (2017–2019 annual mean), the notification rate in humans decreased by 23.0% and 30.4% with and without the data from the United Kingdom, respectively.
- Echinococcus granulosus sensu lato (s.l.) accounted for 67.0% (N = 278) of human cases reported with species information for 2021 while Echinococcus multilocularis accounted for 33.0% (N = 137).
- In total, 22 Member States and two non-Member States provided monitoring data for *Echinococcus* spp. in animals in 2021.
- For *E. multilocularis*, 14 Member States and two non-Member States reported data on 6,318 and 513 tested foxes, respectively, with nine Member States and one non-Member State reporting positive findings and an overall proportion of test-positives of 15.7%.
- Data for 2020 from Finland, Ireland, Malta, the United Kingdom and mainland Norway confirmed the free status of these countries for *E. multilocularis* in accordance with Commission Delegated Regulation (EU) No 2018/772 (EFSA and Zancanaro, 2021).
- For E. granulosus s.l., 19 Member States and two non-Member States reported data from around 96 million animals, primarily domestic livestock (> 99%), with an overall proportion of test-positives of 0.06%. Positive samples came mainly from small ruminants (sheep and goats; 57.2%, mostly from Spain, Greece, Italy and Slovakia), with cattle (bovine animals) accounting



for 20.5% of total positives (mostly from Spain, Italy and Greece) and pigs for 22.1% (mostly from Poland and Spain).

# 8.2. Surveillance and monitoring of cystic and alveolar echinococcosis in humans and animals in the EU

### 8.2.1. Humans

In 2021, 25 MSs reported information on echinococcosis infections in humans. Surveillance is mandatory in 22 MSs and voluntary in three MSs (Belgium, France and the Netherlands), and two MSs (Denmark and Italy) do not have surveillance systems for echinococcosis. The surveillance systems for echinococcosis cover the whole population in all MSs where surveillance systems are in place. All countries reported case-based data, except Bulgaria and the Netherlands, which reported aggregated data. The EU case definition was used by 23 countries; Germany reported using a different case definition, and France did not specify which case definition was used.

Alveolar echinococcosis (AE) caused by the tapeworm *Echinococcus multilocularis* and cystic echinococcosis (CE) caused by *E. granulosus sensu lato (s.l.)* are listed under the common disease name 'echinococcosis' in the EU case definition, thus making no distinction between these two diseases. AE and CE can be reported by species, and as of 2019 (2018 data), by clinical presentation of the disease in the ECDC TESSy database.


### **8.2.2.** Animals

Surveillance of E. multilocularis in Europe is usually carried out on a voluntary basis, except in the reporting countries claiming to be free of this parasite in accordance with the Commission Delegated Regulation (EU) 2018/772 supplementing Regulation (EU) No 576/2013<sup>31</sup>. Surveillance is mainly carried out in red fox, which is Europe's main definitive host. In 2020, Finland, Ireland, mainland Norway (Svalbard archipelago excluded) and the United Kingdom demonstrated the absence of E. multilocularis through the implementation of an annual surveillance programme in compliance with Regulation (EU) 2018/772 (EFSA and Zancanaro, 2021). In accordance with said Regulation, Malta is not required to implement a surveillance programme, due to the absence of the definitive red fox host on its territory. In all other MSs, data on E. multilocularis depend on whether findings are notifiable, monitoring is in place or studies on E. multilocularis are performed. Given that data on E. multilocularis in animals vary geographically (and also within countries) and over time, depending on the sampling effort, it is difficult to compare reported cases of E. multilocularis within and between countries. According to a meta-analysis based on studies published between 1900 and 2015, E. multilocularis has been documented in red foxes from 21 countries (Oksanen et al., 2016) (Figure 26). Since 2018, this parasite has also been found in golden jackals from Croatia, Hungary, Serbia and Slovenia (Dušek et al., 2020; Balog et al., 2021).

In accordance with CIR (EU) 2020/2002, disease-free EU MSs must notify outbreaks of infection with *E. multilocularis* in Canidae to the EU ADIS.<sup>32</sup>

<sup>&</sup>lt;sup>31</sup> Commission Delegated Regulation (EU) 2018/772 of 21 November 2017 supplementing Regulation (EU) No 576/2013 of the European Parliament and of the Council with regard to preventive health measures for the control of *Echinococcus multilocularis* infection in dogs and repealing Delegated Regulation (EU) No 1152/2011 C/2017/7619 OJ L 130, 28.5.2018, pp. 1–10.

<sup>&</sup>lt;sup>32</sup> EU Animal Diseases Information System (ADIS). More information is available at: https://food.ec.europa.eu/animals/animal-diseases/animal-disease-information-system-adis\_en




**Figure 26:** Pooled prevalence of *Echinococcus multilocularis* in red and Arctic foxes within the EU and adjacent countries, depicting the current epidemiological situation in Europe (Oksanen et al., 2016)

Surveillance of *E. granulosus s.l.* is carried out in livestock intermediate hosts during slaughterhouse inspections. In particular, necropsy on sheep liver and lungs is used to detect the presence of parasitic cysts, while molecular PCR-based methods are used to confirm and identify genotype/species belonging to the *Echinococcus* genus (Siles-Lucas et al., 2017). Although *Regulation (EU) 2018/772* is in force for *E. multilocularis*, no specific EU regulation is in place for detecting *E. granulosus s.l.* in animals or humans. Surveillance of the latter parasite therefore depends on national regulations. Approximate geographical distribution of *E. granulosus sensu stricto, E. canadensis* and *E. ortleppi* species causing human cystic echinococcosis in Europe are reported in Figure 27 (Casulli et al., 2022).

18314732, 2022, 12 Downloaded from https://eis.an.oim.elibrary.wiley.com/in/10/2003/j.efs.2022/7666 by Isituo Zooprofilatios Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on (18/12/2023). See the Terms and Conditions (http://oinineibrary.wiley.com/ems-and-conditions) on Wiley Online Library for rules of use; OA article are governed by the applicable Creative Commons of the policies of the applicable o





**Figure 27:** Approximate geographical distribution of the *Echinococcus granulosus sensu lato* species causing human cystic echinococcosis in Europe (2000–2021) (Casulli et al., 2022)

### 8.3. Results

# 8.3.1. Overview of key statistics, EU, 2017–2021

Table 56 summarises EU-level statistics aggregated by year for CE and AE in humans for *E. granulosus s.l.* and *E. multilocularis* in their most relevant definitive and intermediate animal hosts in 2017–2021. More detailed descriptions of these statistics are provided in the below subsections.

**Table 56:** Summary of echinococcosis in humans, caused by *Echinococcus multilocularis* or *Echinococcus granulosus sensu lato (s.l.)* in the most relevant definitive and intermediate animal hosts in the EU, 2017–2021

	2021 <sup>(a)</sup>	2020	2019 <sup>(b)</sup>	2018 <sup>(b)</sup>	2017 <sup>(b)</sup>	Data source
Humans						
Total number of confirmed cases	529	544	769	815	851	ECDC
Total number of confirmed cases/100,000 population (notification rates)	0.15	0.16	0.17	0.21	0.19	ECDC
Number of reporting MSs	25	25	26	25	26	ECDC
Infection acquired in the EU	128	62	176	149	169	ECDC
Infection acquired outside the EU	81	77	96	83	77	ECDC

dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.



18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilattico

Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [1812/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA article are governed by the applicable Creative Common

	2021 <sup>(a)</sup>	2020	2019 <sup>(b)</sup>	2018 <sup>(b)</sup>	2017 <sup>(b)</sup>	Data source
Unknown travel status or unknown country of infection	320	405	497	583	605	ECDC
Animals <sup>(c)</sup>						
Echinococcus multilocularis	in foxes					
Number of animals tested	6,318	5,506	6,326	6,566	7,148	EFSA
% positive animals	17.0	16.1	13.7	18.4	16.9	EFSA
Number of reporting MSs	14	10	13	13	11	EFSA
Echinococcus spp. in dogs						
Number of animals tested	2,942	2,515	2,113	2,605	2,538	EFSA
% positive animals	0.07	0.08	0.24	0.08	0	EFSA
Number of reporting MSs	5	5	6	6	7	EFSA
Echinococcus granulosus s.	<i>l</i> . in cattle (b	ovine anima	ls)			
Number of animals tested	7,065,934	7,035,066	10,956,688	9,920,327	9,833,614	EFSA
% positive animals	0.21	0.21	0.17	0.23	0.22	EFSA
Number of reporting MSs	16	15	16	17	15	EFSA
Echinococcus granulosus s.	<i>l</i> . in sheep a	nd goats				
Number of animals tested	10,817,922	11,089,043	36,890,847	38,870,491	38,278,556	EFSA
% positive animals	0.38	0.96	0.38	0.37	0.40	EFSA
Number of reporting MSs	14	12	15	15	14	EFSA

ECDC: European Centre for Disease Prevention and Control; EFSA: European Food Safety Authority; MSs: Member States.

### 8.3.2. Human echinococcosis

In 2021, 529 confirmed echinococcosis cases were reported by 25 MSs, corresponding to an EU notification rate of 0.15 cases per 100,000 population (Table 57). This was a decrease of 7.5% compared with 2020 (0.16 per 100,000 population). A mean notification rate of 0.19 was reported over 2017–2019. Compared with the rate before the COVID-19 pandemic, the notification rate in humans decreased by 23.0% and 30.4% with and without the data from the United Kingdom, respectively. In 2021, 22 MSs reported at least one confirmed case, while three MSs reported zero cases (Cyprus, Malta and the Netherlands). In 2021, the highest notification rate was observed in Bulgaria, with 1.3 cases per 100,000 population, followed by Lithuania, Slovenia and Austria with 0.72, 0.52 and 0.47 cases per 100,000 population, respectively. Germany, Bulgaria and France reported the highest numbers of cases, with 152 (28.7%), 89 (16.8%) and 75 (14.2%) cases out of 529, respectively.

Most echinococcosis cases (60.5%; 320/529) were reported without data on their importation and probable country of infection; 61.2% (128/209) of cases reported with such information were domestic or related to travel within the EU, while 38.8% (81/209) were associated with travel outside the EU. In 2021, 14 MSs out of the 25 reporting MSs notified that all their *Echinococcus* spp. infections were domestically acquired. Considering data with information on travel-related status, the highest proportions of cases (N = 107) were reported by Germany (100%; 59 vs. 0 non-travel-related cases), Sweden (100%; 18 vs. 0 cases), Belgium (72%; 13 vs. 5 cases) and Austria (48%; 11 vs. 12 cases). At the species level, human *E. granulosus s.l.* infections were more often reported as travel-associated than human *E. multilocularis* infections, accounting for 95.2% (N = 79) and 4.8% (N = 4) of cases reported with this information, respectively. Of the 81 travel-associated cases of *Echinococcus* spp. for which the origin of infection is known, most (83.5%) were reported as originating from outside the EU, mainly from Syria (37%), followed by Turkey (11.9%), Iraq (9.9%) and Morocco (7.4%). Within the EU, Bulgaria (45%), Romania (40%), Luxembourg (10%) and Austria (5%) were reported as the probable countries of infection in 20 cases.

<sup>(</sup>a): Data on animals from the United Kingdom (Northern Ireland) are taken into account for 2021. In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/Northern Ireland, the EU requirements on data sampling are also applicable to Northern Ireland.

<sup>(</sup>b): Data from the United Kingdom are taken into account for 2017–2019, because the United Kingdom was an EU MS, but it became a third country on 1 February 2020.

<sup>(</sup>c): Single tested animals.



**Table 57:** Reported human cases of cystic and alveolar echinococcosis and notification rates per 100,000 population in EU MSs and non-MS countries, by country and year, 2017–2021

		2021			20	20	20	19	20	18	20	17
Country	National coverage <sup>(a)</sup>	Data format <sup>(a)</sup>	Confi cases rat	and	Confi cases rat	and	Confi cases rat		Confi cases rat	and	case	irmed s and ites
			Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate
Austria	Y	С	42	0.47	34	0.38	36	0.41	46	0.52	50	0.57
Belgium	Υ	С	18	0.16	19	0.16	22	0.19	15	0.13	13	0.11
Bulgaria	Y	Α	89	1.29	95	1.37	193	2.76	206	2.92	218	3.07
Croatia	Y	С	3	0.07	3	0.07	3	0.07	4	0.10	15	0.36
Cyprus	Υ	С	0	0	1	0.11	0	0	0	0	0	0
Czechia	Υ	С	1	0.01	4	0.04	1	0.01	4	0.04	1	0.01
Denmark <sup>(b)</sup>	_	_	_	_	_	_	_	_	_	_	_	_
Estonia	Υ	С	4	0.30	1	0.08	2	0.15	0	0	1	0.08
Finland <sup>(c)</sup>	Υ	С	6	0.11	4	0.07	8	0.14	1	0.02	5	0.09
France	Υ	С	75	0.11	53	0.08	45	0.07	62	0.09	53	0.08
Germany	Y	С	152	0.18	170	0.20	150	0.18	176	0.21	141	0.17
Greece	Υ	С	4	0.04	7	0.07	7	0.07	11	0.10	15	0.14
Hungary	Y	С	7	0.07	4	0.04	10	0.10	9	0.09	14	0.14
Ireland <sup>(c)</sup>	Υ	С	1	0.02	0	0	0	0	2	0.04	0	0
Italy <sup>(b)</sup>	_	_	_	_	-	_	-	_	-	_	_	_
Latvia	Y	С	6	0.32	5	0.26	6	0.31	10	0.52	6	0.31
Lithuania	Υ	С	20	0.72	37	1.32	81	2.90	50	1.78	53	1.86
Luxembourg	Υ	С	1	0.16	3	0.48	1	0.16	0	0	2	0.34
Malta <sup>(c)</sup>	Υ	С	0	0	0	0	0	0	0	0	0	0
Netherlands	Υ	Α	0	0	48	0.28	48	0.28	42	0.24	38	0.22
Poland	Y	С	26	0.07	18	0.05	70	0.18	51	0.13	75	0.20
Portugal	Υ	С	2	0.02	1	0.01	5	0.05	9	0.09	2	0.02
Romania	Y	С	1	0.01	0	0	1	0.01	4	0.02	14	0.07
Slovakia	Υ	С	2	0.04	3	0.05	11	0.20	10	0.18	7	0.13
Slovenia	Y	С	11	0.52	3	0.14	6	0.29	6	0.29	7	0.34
Spain <sup>(d)</sup>	Υ	С	33	_	8	_	34	0.07	68	0.15	83	0.18
Sweden	Υ	С	25	0.24	23	0.22	26	0.25	29	0.29	34	0.34
EU Total 27			529	0.15	544	0.16	766	0.20	815	0.21	847	0.22
United Kingdom <sup>(c)</sup>	Υ	С	_	_	_	_	3	0	_	_	4	0.01
EU Total <sup>(e)</sup>			529	0.15	544	0.16	769	0.17	815	0.21	851	0.19
Iceland	Y	С	0	0	0	0	0	0	0	0	0	0
Norway <sup>(c)</sup>	Υ	С	11	0.20	6	0.11	7	0.13	7	0.13	6	0.11
Liechtenstein <sup>(b)</sup>	_	_	_	_	_	_	_	_	_	_	-	_
Switzerland <sup>(b)</sup>	_	_	_	_								

<sup>-:</sup> Data not reported.

18314732, 2022, 12, Downloaded from https://eftsaonlinelibrary.wiley.com/doi/10.2903j.eftsa.2022.7666 by Islituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18122023]. See the Terms and Conditions, wiley convierms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Ceravive Commons.

<sup>(</sup>a): Y: yes; N: no; A: aggregated data; C: case-based data.

<sup>(</sup>b): No surveillance system.

<sup>(</sup>c): Finland, Ireland, Malta, the United Kingdom and mainland Norway have been declared free of Echinococcus multilocularis.

<sup>(</sup>d): Data not complete for 2020–2021, rate not estimated.

<sup>(</sup>e): Cases reported by the United Kingdom for the period 2017–2019 were also considered for this estimation (EU-28). When the United Kingdom data were collected for the period 2017–2019, the United Kingdom was an EU MS, but it became a third country on 1 February 2020.



18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilattico

dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

In 2021, species information was provided for 415 confirmed echinococcosis cases (78.4%) out of 529 confirmed cases reported by 19 MSs (Table 58). In 2021, human infections caused by *E. multilocularis* accounted for 137 cases (33.0% of cases with known species information), more than in 2020 and slightly fewer than in 2017–2019. In 2021, Germany and France reported the highest number of human cases caused by *E. multilocularis*, accounting for 38.7% and 29.2% of all reported *E. multilocularis* cases, respectively. Human infections caused by *E. granulosus s.l.* accounted for 67.0% (278 cases) of the cases with species information available. In 2021, Bulgaria and Germany reported the highest number of human cases caused by *E. granulosus s.l.*, accounting in both cases for 32.0% of all reported *E. granulosus s.l.* cases.

In 2021, 13 MSs provided information on hospitalisations, covering 22.9% (121/529) of all confirmed cases of echinococcosis in the EU. The overall hospitalisation rate was 42.1%. Information on the outcome of the cases was provided by 15 MSs for 51.0% of confirmed cases with no deaths reported. Information on age was reported by 19 MSs with the highest proportion of cases (83.1%) occurring in over 30-year-olds.

**Table 58:** Reported human cases of cystic and alveolar echinococcosis in EU MSs and non-MS countries, by country, year and *Echinococcus* species, 2017–2021

	2	2021		2020			2019			2	2018			2017	
Country	Total Cases	Eg	Em	Total Cases	Eg	Em	Total Cases	Eg	Em	Total Cases	Eg	Em	Total Cases	Eg	Em
Austria	42	24	7	34	18	4	36	16	13	46	29	12	50	37	8
Belgium	18	11	7	19	10	8	22	12	10	15	10	5	13	9	4
Bulgaria	89	89	0	95	95	0	193	193	0	206	206	0	218	218	0
Croatia	3	_	_	3	_	_	3	_	_	4	_	_	15	_	_
Cyprus	0	0	0	1	_	_	0	0	0	0	0	0	0	0	0
Czechia	1	0	1	4	1	2	1	_	_	4	1	2	1	_	_
Denmark <sup>(a)</sup>	_	_	-	-	_	_	_	_	_	_	_	_	_	_	_
Estonia	4	_	_	1	1	0	2	_	_	0	0	0	1	0	1
Finland <sup>(b)</sup>	6	5	_	4	3	_	8	8	0	1	1	0	5	5	0
France	75	22	53	53	11	42	45	10	35	62	21	41	53	5	48
Germany	152	89	40	170	80	54	150	87	41	176	93	59	141	86	35
Greece	4	_	_	7	7	0	7	_	_	11	_	_	15	_	_
Hungary	7	_	_	4	1	_	10	_	_	9	_	_	14	1	1
Ireland <sup>(b)</sup>	1	1	0	0	0	0	0	0	0	2	_	_	0	0	0
Italy <sup>(a)</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
Latvia	6	6	0	5	5	0	6	4	_	10	5	1	6	4	_
Lithuania	20	2	14	37	_	_	81	30	21	50	11	17	53	19	20
Luxembourg	1	_	_	3	3	0	1	1	0	0	0	0	2	2	0
Malta <sup>(b)</sup>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Netherlands	0	0	0	48	_	_	48	_	_	42	_	_	38	_	_
Poland	26	12	11	18	8	6	70	21	25	51	17	19	75	27	31
Portugal	2	2	0	1	1	0	5	5	0	9	9	0	2	_	_
Romania	1	_	_	0	0	0	1	_	_	4	_	_	14	_	_
Slovakia	2	0	2	3	1	2	11	3	8	10	3	3	7	2	3
Slovenia	11	2	2	3	1	_	6	1	_	6	3	_	7	_	_
Spain <sup>(c)</sup>	33	1	_	8	1	_	34	6	_	68	12	_	83	4	_
Sweden	25	12	0	23	8	3	26	17	2	29	5	2	34	11	4
EU Total 27	529	278	137	544	255	121	766	414	155	815	426	161	847	430	155
United Kingdom <sup>(b)</sup>	_	_	_	_	_	_	3	3	0	_	_	_	4	4	0
EU Total <sup>(d)</sup>	529	278	137	544	255	121	769	417	155	815	426	161	851	434	155
Iceland	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Norway <sup>(b)</sup>	11	5	_	6	1	1	7	2	_	7	5	_	6	3	1



/efsa.onlinelbrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023], See the Terms and Conditions (https://onlinelbrary.wiley.com/terms

-and--conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

•	2021			2020		2019			2018			2017			
Country	Total Cases	Eg	Em	Total Cases	Eg	Em	Total Cases	Eg	Em	Total Cases	Eg	Em	Total Cases	Eg	Em
Liechtenstein <sup>(a)</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	-
Switzerland <sup>(a)</sup>	_	_	_	_											

Eg: Echinococcus granulosus sensu lato; Em: Echinococcus multilocularis.

- -: Data not reported.
- (a): No surveillance system.
- (b): Finland, Ireland, Malta, the United Kingdom and mainland Norway have been declared free of E. multilocularis.
- (c): Data not complete for 2020-2021.
- (d): Cases reported by the United Kingdom for the period 2017–2019 were also considered for this estimation (EU-28). When the United Kingdom data were collected for the period 2017–2019, the United Kingdom was an EU MS, but it became a third country on 1 February 2020.

### 8.3.3. *Echinococcus* spp. in animals and food

### Monitoring data for Echinococcus multilocularis

Table 59 summarises the most relevant definitive and intermediate host species tested for *E. multilocularis*. In 2021, results were reported by 18 MSs and two non-MSs (Norway and Switzerland). Austria, Bulgaria, Croatia, Lithuania and Portugal reported no animal monitoring data for *E. multilocularis* or *E. granulosus s.l.* 

In 2021, 13 MSs and the United Kingdom (Northern Ireland) reported monitoring data from 6,318 foxes examined for *E. multilocularis*. Also, two non-MSs (Norway and Switzerland) reported on monitoring data from a total of 513 foxes examined for *E. multilocularis*. Nine MSs and one non-MS (Switzerland) reported a total of 15.7% positive samples: Czechia (23.4% positive samples), France (13.7%), Germany (29.5%), Luxembourg (20.6%), the Netherlands (1.7%), Poland (35.3%), Slovakia (5.2%), Sweden (4.7%) and Switzerland (50%). Czechia (N = 645) reported the largest number of infected foxes in Europe, accounting for 60.2% of positive findings.

In addition to its presence in foxes (as definitive hosts), *Echinococcus* spp. have been reported in one beaver (France), one cat (Switzerland), 16 dogs (France, Poland and Switzerland), one golden jackal (Switzerland), five wild boars (Germany and Switzerland), three wolves (Switzerland) and one lynx (Switzerland). Moreover, 13,086 and 34 positive pigs were reported by two MSs (Poland and Slovakia), co-endemic for both *Echinococcus* species and five pigs from one non-MS, endemic only for *E. multilocularis* (Switzerland). Poland reported 10 cases as *E. multilocularis* and 13,076 as *Echinococcus* unspecified; Slovakia reported 43 cases as *Echinococcus* unspecified, while Switzerland reported five cases as *E. multilocularis*. It should also be emphasised that pigs are good hosts for *E. granulosus s.l.* while *E. multilocularis* metacestodes in pigs are abortive, and their presence is often used as a sentinel for the circulation of this parasite in animal hosts. For this reason, the presence of both *E. multilocularis* and *E. granulosus s.l.* may be overestimated in co-endemic countries with unknown species identification. In this context, it should also be noted that positive samples from pigs, dogs and wolves without species identification were mentioned in Tables 59 and/or 60 only for countries with known circulation of both *E. granulosus s.l.* and *E. multilocularis. E. multilocularis* was also detected in one sample batch of fresh leaf vegetables in France.



**Table 59:** Echinococcus multilocularis monitoring results in wild and domestic animals in reporting EU MSs and non-MSs, 2021

	Presence of				N Positive/N tes	sted (% positive)		
Country	Em/Eg <sup>(a)</sup>	Beavers	Cats	Coypu	Dogs	Foxes	Jackals	Pigs
Czechia	Em/Eg	_	_	_	_	645/2,758 (23.4)	-	_
Denmark	Em	_	_	_	_	_	_	0/18,592,853 (0)
Estonia	Em/Eg	_	_	_	_	_	_	0/548,849 (0) <sup>(e)</sup>
Finland	Eg	_	_	_	_	0/244 (0)	_	_
France	Em/Eg	1/1 (100)	0/1 (0) <sup>(e)</sup>	_	1/52 (1.9) <sup>(e)</sup>	32/233 (13.7)	_	_
Germany	Em	_	0/2 (0)	_	0/322 (0)	179/607 (29.5)	_	0/1 (0)
Hungary	Em/Eg	_	_	_	_	0/1 (0)	-	0/61 (0) <sup>(e)</sup>
Ireland <sup>(b)</sup>	Eg	_	_	_	_	0/398 (0)	_	_
Italy	Em/Eg	_	_	_	0/58 (0) <sup>(e)</sup>	0/139 (0)	_	0/5,087,079 (0) <sup>(e)</sup>
Latvia	Em/Eg	_	_	_	_	_	_	0/473,831 (0) <sup>(e)</sup>
Luxembourg	Em	_	_	_	_	22/107 (20.6)	_	0/122,548 (0)
Netherlands	Em	_	_	_	_	3/176 (1.7)	_	_
Poland	Em/Eg	_	_	_	1/90 (1.1) <sup>(e)</sup>	120/340 (35.3) <sup>(e)</sup>	_	13,086/21,439,046 (0.06) <sup>(e)</sup>
Romania	Em/Eg	_	_	_	_	_	_	0/53 (0) <sup>(e)</sup>
Slovakia	Em/Eg	_	0/768 (0) <sup>(e)</sup>	_	0/2,420 (0) <sup>(e)</sup>	5/96 (5.2) <sup>(e)</sup>	_	43/675,234 (0.01) <sup>(e)</sup>
Slovenia	Em/Eg	_	_	_	_	49/184 (26.6)	_	0/242,584 (0) <sup>(e)</sup>
Sweden	Em/Eg	_	_	_	_	16/343 (4.7)	_	0/2,651,110 (0) <sup>(e)</sup>
United Kingdom (Northern Ireland) <sup>(c)</sup>	Eg	_	_	_	_	0/692 (0)	_	_
EU Total (27 + XI)		1/1 (100)	0/771 (0)	_	2/2,942 (0.07)	1071/6,318 (17)	-	13,129/49,833,249 (0.03)
Norway <sup>(d)</sup>	Eg	_	_	_	_	0/511 (0)		_
Switzerland	Em	0/2 (0)	1/7 (14.3)	1/1 (100)	14/42 (33.3)	1/2 (50)	1/1 (1)	5/5 (100)
Total non-EU countries		0/2 (0)	1/7 (14.3)	1/1 (100)	14/42 (33.3)	1/513 (0.2)	1/1 (1)	5/5 (100)
Total EU (27+ XI) + non-EU countries		1/3 (33.3)	1/778 (0.13)	1/1 (100)	16/2,984 (0.54)	1,072/6,831 (15.7)	1/1 (1)	13,134/49,833,254 (0.03)



				N Positive	e/N tested (% positiv	re)		
Country	Presence of Em/ Eg <sup>(a)</sup>	Raccoon dogs	Raccoons	Wild boars	Wild cats (Felis silvestris)	Wolves	Voles	Lynx
Czechia	Em/Eg	_	-	_	_	_	-	_
Denmark	Em	_	_	-	-	_	_	_
Estonia	Em/Eg	_	_	_	_	_	_	_
Finland	Eg	0/409 (0)	_	_	_	_	0/618 (0)	_
France	Em/Eg	_	_	_	_	_	_	_
Germany	Em	_	0/3 (0)	4/4 (100)		0/1 (0)	_	_
Hungary	Em/Eg	_	_	_	_	_	_	_
Ireland <sup>(b)</sup>	Eg	_	_	_	_	_	_	_
Italy	Em/Eg	_	_	0/55,673 (0)		0/55 (0) <sup>(e)</sup>	_	_
Latvia	Em/Eg	_	_	_	_	_	_	_
Luxembourg	Em	_	_	_	_	_	_	_
Netherlands	Em	_	_	_	_	_	_	_
Poland	Em/Eg	_	_	_	_	_	_	_
Romania	Em/Eg	_	_	_	_	_	_	_
Slovakia	Em/Eg	_	_	_	0/1 (0)	0/9 (0) <sup>(e)</sup>	_	_
Slovenia	Em/Eg	_	_	_	_		_	_
Sweden	Em/Eg	_	_	0/16,569 (0)	_	0/51 (0) <sup>(e)</sup>	_	_
United Kingdom (Northern Ireland) <sup>(c)</sup>	Eg	_	_	_	_	_	_	_
EU Total (27 + XI)		0/409 (0)	0/3 (0)	4/72,246 (0.01)	0/1 (0)	0/116 (0)	0/618 (0)	_
Norway <sup>(d)</sup>	Eg	_	-	_	_	_	_	_
Switzerland	Em	_	_	1/1 (100)	0/1 (0)	3/5 (60)	_	1/1 (100)
Total non-EU countries		_	-	1/1 (100)	0/1 (0)	3/5 (60)	_	1/1 (100)
Total EU (27+ XI) + non-EU countries		0/409 (0)	0/3 (0)	5/72,247 (0.01)	0/2 (0)	3/121 (2.5)		1/1 (100)

<sup>-:</sup> Data not reported.

Em: Echinococcus multilocularis; Eq: Echinococcus granulosus sensu lato.

<sup>(</sup>a): Presence in the country of Echinococcus multilocularis (Em) and/or Echinococcus granulosus sensu lato (Eg).

<sup>(</sup>b): Member States listed in the Annex to Commission Implementing Regulation (EU) 2018/878 on the application of preventive health measures for the control of E. multilocularis infection in dogs.

<sup>(</sup>c): Data on animals from the United Kingdom (Northern Ireland) are taken into account for 2021. In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the/Northern Ireland protocol, the EU requirements on data sampling are also applicable to Northern Ireland.

<sup>(</sup>d): Mainland Norway (excluding the Svalbard archipelago where E. multilocularis was documented).

<sup>(</sup>e): Positive samples reported from dogs, cats, wolves and pigs without *Echinococcus* species information were mentioned in the table only for countries with known circulation of both *E. multilocularis* and *E. granulosus sensu lato*.



18314732, 2022, 12, Downloaded from https://disa.onlinelibrary.wiley.com/doi/10.2903/j.disa.2022.7666 by Isituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.2903/j.disa.2022.7666 by Isituto Zooprofilation Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.2903/j.disa.2022.7666 by Isituto Zooprofilation Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.2903/j.disa.2022.7666 by Isituto Zooprofilation Sperimenta dell'Umbria e delle Marche (IZSUM). Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.2903/j.disa.2022.7666 by Isituto Zooprofilation Sperimenta dell'Umbria e dell'Um


and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

### Monitoring data for Echinococcus granulosus sensu lato

Table 60 summarises the most relevant definitive and intermediate host species tested for  $E.\ granulosus\ s.l.$  In 2021, 19 MSs and two non-MSs (Norway and Switzerland) reported monitoring data for  $E.\ granulosus\ s.l.$  The data reported were from around 93 million domestic and wild animals tested for  $E.\ granulosus\ s.l.$ , of which more than 99% were domestic animals (sheep, cattle, goats, pigs, horses, water buffalos, dogs and cats). A large proportion of these data was obtained from domestic livestock during meat inspections at the slaughterhouse. Wild animals tested included Cantabrian chamois, deer, moose, mouflons, reindeer, wild boars and wolves. A total of 71,869 (0.07%) positive samples were reported by 11 MSs, mainly from domestic animals. These positive samples came mainly from small ruminants (sheep and goats; N = 41,081), accounting for 57.2% of positive results. Positive tests in small ruminants were mainly reported by Spain (79.8%), followed by Greece, Italy and Slovakia. Results showed 14,769 positive cattle (20.5% of animals positive for  $E.\ granulosus\ s.l.$ ), mainly reported by Spain (76.52%), followed by Italy and Greece, and 15,874 positive pigs (22.1% of animals positive for  $E.\ granulosus\ s.l.$ ), of which 82.4% were reported by Poland, followed by Spain.

Belgium, Denmark, Estonia, Ireland, Latvia, Malta and Slovenia among MSs and Norway among non-MSs, reported no positive findings for *E. multilocularis* or *E. granulosus s.l.* Austria, Bulgaria, Croatia, Lithuania and Portugal reported no animal monitoring data for *E. multilocularis* or *E. granulosus s.l.* 

Figures 28 and 29 show the cumulative proportion of positive samples from different intermediate hosts of *E. granulosus s.l.* and their geographical distribution in EU MSs and other European countries for the period between 2017 and 2021. Small ruminants (sheep and goats) accounted for 75.5% (2017–2021) of all positive samples and were reported by a few countries with large animal populations (Greece, Italy and Spain and the United Kingdom). Positive results in cattle (11.8%; 2017–2021) were also mainly reported by Greece, Italy and Spain. Positive results in pigs (12.6%; 2017–2021) were mainly reported by Poland, followed by Spain.




**Figure 28:** Cumulative proportion (%) of test-positive animals for *Echinococcus granulosus sensu lato*, by intermediate host species, in EU MSs and non-MSs, 2017–2021



onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico

Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.



Intermediate hosts included on the map are cattle, deer, goats, horses, moose, mouflons, pigs, reindeer, sheep, water buffalos and wild boars. It should be stressed that pigs are intermediate hosts for both *E. multilocularis* and *E. granulosus s.l.* For this reason, data from pigs were excluded from co-endemic countries – Poland, Romania, Slovakia and Slovenia – where *Echinococcus* species information was not reported. Spain (N = 302,433), Italy, (N = 248,530), United Kingdom<sup>33</sup> (N = 71,861), Greece (N = 53,140), Bulgaria (N = 12,214), Poland (N = 2,630) and Slovakia (N = 995) were the countries with the highest endemicity for *E. granulosus s.l.* in the EU in 2017–2021.

**Figure 29:** Map of the cumulative proportion (%) of test-positive animals for *Echinococcus granulosus* sensu lato in MSs and non-MSs, by country, 2017–2021

33

<sup>&</sup>lt;sup>33</sup> Data from the United Kingdom are taken into account for 2017–2019, because the United Kingdom was an EU MS over this period but it became a third country on 1 February 2020. United Kingdom (Northern Ireland) did not report monitoring data for 2021 from intermediate hosts for *Echinococcus granulosus sensu lato*.



**Table 60:** Echinococcus granulosus sensu lato monitoring results in wild and domestic animals in reporting EU MSs and non-MSs, 2021

				N Pos	itive/N test	ed (% posi	itive)		
Country	Presence of Em/Eg <sup>(a)</sup>	Cantabrian chamois	Cats	Cattle (bovine animals)	Deer	Dogs	Moose	Mouflons	Pigs
Belgium <sup>(b)</sup>	Em	_	_	0/770,235 (0)	_	_	_	_	_
Cyprus	Eg	_	_	_	_	_	_	2/43 (4.7)	_
Denmark <sup>(b)</sup>	Em	_	_	0/452,600 (0)	_	_	_	_	_
Estonia	Em/Eg	_	_	0/34,011 (0)	_	_	_	_	0/548,849 (0) <sup>(c)</sup>
Finland	Eg	-	_	0/257,680 (0)	0/2,765 (0)	-	1/212 (0.47)	-	0/1,942,376 (0)
France	Em/Eg	_	0/1 (0) <sup>(c)</sup>	0/1 (0)	_	0/51 (0) <sup>(c)</sup>	_	_	
Germany	Em	-	_	0/12 (0)	0/2 (0)	-	_	_	
Greece	Eg	_	_	930/45,462 (2.0)	_	_	_	_	29/250,835 (0.01)
Hungary	Em/Eg	_	_	4/8 (50.0)	_	_	_	_	1/62 (1.6) <sup>(c)</sup>
Italy	Em/Eg	0/1 (0)	_	2,515/2,493,943 (0.10)	0/629 (0)	0/58 (0) <sup>(c)</sup>	_	0/12 (0)	10/5,087,089 (< 0.001) <sup>(c)</sup>
Latvia	Em/Eg	_	_	0/73,531 (0)	_	_	_	_	0/473,831 (0) <sup>(c)</sup>
Luxembourg <sup>(b)</sup>	Em	_	_	0/27,326 (0)	_	_	_	_	
Malta	Eg	-	_	_	_	-	_	_	
Poland	Em/Eg	-	_	_	_	0/89 (0) <sup>(c)</sup>	-	_	13,076/21,439,036 (0.06) <sup>(c)</sup>
Romania	Em/Eg	_	_	18/21 (85.7)	_	_	_	_	0/53 (0) <sup>(c)</sup>
Slovakia	Em/Eg	_	0/768 (0) <sup>(c)</sup>	0/34,771 (0)	_	0/2,420 (0) <sup>(c)</sup>	-	_	43/675,234 (0.01) <sup>(c)</sup>
Slovenia	Em/Eg	_	_	0/123,961 (0)	_	_	_	_	0/242,584 (0) <sup>(c)</sup>
Spain	Eg	-	_	11,302/2,340,722 (0.48)	2/185,894 (0)	-	_	0/7,412 (0)	2,714/41,148,859 (0.01)
Sweden	Em/Eg	_	_	0/411,650 (0)	0/6,631 (0)	-	_	_	0/2,651,110 (0) <sup>(c)</sup>
EU Total (27 + XI)		0/1 (0)	0/769 (0)	14,769/7,065,934 (0.21)	2/195,921 (0)	0/2,618 (0)	1/212 (0.47)	2/7,467 (0.03)	15,873/74,459,918 (0.02)
Norway	Eg	_	_	0/301,000 (0)	_	_	_	-	0/1,568,000 (0)
Switzerland <sup>(b)</sup>	Em	_	_	0/2 (0)	0/1 (0)	_	_	-	
Total non-EU countries		_	_	0/301,002 (0)	0/1 (0)	_	_	-	0/1,568,000 (0)
Total EU (27+ XI) + non-EU countries		0/1 (0)	0/769 (0)	14,769/7,366,936 (0.20)	2/195,922 (0)	0/2,618 (0)	1/212 (0.47)	2/7,467 (0.03)	15,873/76,027,918 (0.02)



				N Positive/N te	sted (% posit	ive)		
Country	Presence of Em/Eg <sup>(a)</sup>	Reindeers	Goats	Sheep	Solipeds, domestic	Water buffalos	Wild boars	Wolves
Belgium <sup>(b)</sup>	Em	-	-	-	_	-	_	-
Cyprus	Eg	_	_	_	_	_	_	_
Denmark <sup>(b)</sup>	Em	_	_	-	_	_	_	_
Estonia	Em/Eg	_	0/171 (0)	0/8,378 (0)	0/8 (0)	_		
Finland*	Eg	7/56,079 (0.01)	0/748 (0)	0/62,724* (0)	0/757 (0)	_	0/183 (0)	7/41 (17.1)
France	Em/Eg	_	_	0/2 (0)	_	_	_	_
Germany	Em	_	_	0/2 (0)	_	_	_	_
Greece	Eg	_	1,477/259,488 (0.57)	4,490/621,068 (0.72)	_	-	_	-
Hungary	Em/Eg	_	_	3/3 (100)	_	_	_	_
Italy	Em/Eg	_	272/48,039 (0.57)	1,292/107,2250 (0.12)	0/537 (0)	114/33,912 (0.34)	7/55,680 (0.01)	0/55 (0) <sup>(c)</sup>
Latvia	Em/Eg	_	0/510 (0)	0/26,943 (0)	0/34 (0)	_	_	_
Luxembourg <sup>(b)</sup>	Em	_	_	-	_	_	_	_
Malta	Eg	-	_	0/7,373 (0)	_	_	_	_
Poland	Em/Eg	_	_		_	_	_	_
Romania	Em/Eg	_	0/3 (0)	0/30 (0)	_	_	_	_
Slovakia	Em/Eg	_	0/91 (0)	775/6,448 (12.0)	_	_	_	0/9 (0) <sup>(c)</sup>
Slovenia	Em/Eg	_	0/1,172 (0)	0/10860 (0)	0/1,057 (0)	_	_	_
Spain	Eg	_	3,803/783,974 (0.49)	28,969/7,679,758 (0.38)	0/6,637 (0)	-	5/138,599 (< 0.001)	_
Sweden	Em/Eg	0/40,528 (0)	0/817 (0)	0/227,070 (0)	1/1,200 (0.08)	_	0/16,569 (0)	0/51 (0) <sup>(c)</sup>
EU Total (27 + XI)		7/96,607 (0.01)	5,552/1,095,013 (0.51)	35,529/9,722,909 (0.37)	1/10,230 (0.01)	114/33,912 (0.34)	12/211,031 (0.01)	7/156 (4.5)
Norway	Eg	_	0/28,600 (0)	0/1,204,600 (0)	_	_	_	
Switzerland <sup>(b)</sup>	Em	_	_	_	_	_	_	
Total non-EU countries		_	0/28,600 (0)	0/1,204,600 (0)	_	_	_	-
Total EU (27+ XI) + non-EU countries		7/96,607 (0.01)	5,552/1,123,613 (0.49)	35,529/10,927,509 (0.33)	1/10,230 (0.01)	114/33,912 (0.34)	12/211,031 (0.01)	7/156 (4.5)



- -: Data not reported; Em: Echinococcus multilocularis; Eg: Echinococcus granulosus sensu lato.
- \*: During the finalisation of this report, Finland indicated a reporting error: only 51,221 sheep were tested.
- (a): Presence in the country of E. multilocularis (Em) and/or E. granulosus sensu lato (Eg).
- (b): Reporting countries with known circulation of *E. multilocularis* only, and which tested suitable hosts for *E. granulosus sensu lato*.
- (c): Positive samples from dogs, cats, wolves and pigs without *Echinococcus* species information reported, were mentioned in the table only for MSs with known circulation of both *E. multilocularis* and *E. granulosus sensu lato*.



### 8.4. Discussion

The EU notification rate of confirmed human echinococcosis cases decreased slightly from 2017 to 2019. In 2020 and 2021, the EU notification rates for infections caused by *Echinococcus* species fell sharply compared with the previous 3 years, probably due to the COVID-19 pandemic and partly due to the withdrawal of the United Kingdom from the EU. In 2021, although the COVID-19 pandemic was still ongoing, a slight decrease was observed in the number of confirmed echinococcosis cases with respect to 2020.

In a few countries, the increase in case numbers over the last few years may be explained by increased human surveillance activities and improved notification systems for these diseases. Increased awareness of the diseases among clinicians, as well as immigration from endemic countries, may also have influenced the number of diagnosed cases in some countries (Richter et al., 2019). It should be emphasised that the true prevalence of these diseases is extremely difficult to estimate due to the long incubation period, high proportion of asymptomatic or paucisymptomatic carriers who never seek medical attention, non-specific symptoms and under-reporting/misdiagnosed cases. The abovementioned factors contribute to the neglected status of these diseases (Casulli, 2020). For these reasons, the patchy data reported by MSs on the number of people with echinococcosis, currently represent the 'tip of the iceberg' for infections, with asymptomatic carriers and misdiagnosed cases of CE/AE making up the invisible portion. It has been estimated that the official figures from hospital records should be far higher, with true values 10 and 700 times greater for Bulgaria and Romania, respectively (Tamarozzi et al., 2018). The prospective, observational, multicentre and online 'European Register of Cystic Echinococcosis' (ERCE) (Rossi et al., 2020) is currently seeking to establish a voluntary system to collect harmonised clinical data in the EU. The European (Alveolar) Echinococcosis Registry (Kern et al., 2003) organised a similar initiative in the past.

In 2021, 22 MSs reported monitoring data on E. granulosus s.l. and/or E. multilocularis in animals. The highest numbers of animals infected with E. granulosus s.l. were reported in Spain, Greece, Italy and Poland, and were mainly observed in sheep intermediate hosts, and secondarily in cattle and pigs. Most of the animals (mainly red foxes) infected with E. multilocularis were reported by Czechia, followed by France, Germany, Poland and Slovenia. The surveillance of E. multilocularis in foxes is important to assess the prevalence of AE in Europe, given that its geographical distribution seems to have widened over the last decades. It is difficult to establish whether the increased geographical distribution of E. multilocularis is due to a growing fox population in Europe (Oksanen et al., 2016) and the expansion of their habitat to urban areas (Deplazes et al., 2004), or whether it reflects greater efforts in surveillance, as there is a general lack of baseline data and standardised detection methods. One growing source of concern is the detection in recent years of E. multilocularis in golden jackals, which are increasing their geographical range from southern-eastern to northern-western Europe, and therefore contributing to the dispersion of this parasite over long distances (Dušek et al., 2020; Balog et al., 2021). Moreover, in animals, notification is necessary to obtain reliable data, and information on parasite speciation is essential for risk management efforts, since E. granulosus s.l. and E. multilocularis have different epidemiology and pose different health risks for humans (Possenti et al., 2016; Conraths et al., 2017; Casulli, 2020). For E. granulosus s.l., a notification requirement would ensure that comparable data between MSs are obtained from meat inspections of foodproducing animals. For E. multilocularis, while the need for general notification by all MSs may raise questions, it is nevertheless required in countries that are free from this parasite, in accordance with Regulation (EU) No 2018/772.

In general terms, it should be emphasised that animal findings from most endemic countries fluctuated from year to year over the period 2017–2021, although positive findings in animals and humans were reported in most years. The fluctuations over the last 5 years probably reflect the investigational efforts performed during a particular year, rather than a change in the true prevalence of the disease. For instance, Italy reported 1,292 (0.12%) test-positive sheep in 2021, but documented 60,608 (6.5%) in 2020. Moreover, these fluctuations in animals can be partially explained by the United Kingdom leaving the EU: in 2019 the United Kingdom accounted for 68.8% and 37.2% of all EU-tested sheep and goats, and cattle, respectively.



# Foodborne outbreaks (in accordance with Directive 2003/99/EC)

Summary data and figures substantiating this chapter, as well as additional information on related projects and internet sources are published for this report on the EFSA Knowledge Junction on Zenodo here. Summary statistics on foodborne outbreaks reported in the framework of Directive 2003/99/EC are retrievable using the EFSA dashboard on foodborne outbreaks available here. Summary statistics on human surveillance data with downloadable files are retrievable using the ECDC Surveillance Atlas of Infectious Diseases here.

# 1. Key facts

- In 2021, 27 EU MSs and the United Kingdom (Northern Ireland) reported 4,005 foodborne outbreaks, 32,543 cases of illness, 2,495 hospitalisations and 31 deaths. At the same time, 83 outbreaks, 1,270 cases of illness, 65 hospitalisations and 2 deaths were reported by seven non-MSs.
- In 2021, foodborne outbreaks in the EU increased by 29.8% compared with the previous year (3,086 in 2020). Human cases and hospitalisations also increased, by 62.6% (20,017 cases in 2020) and 49.0% (1,675 hospitalisations in 2020), respectively. However, the total number of outbreaks, cases and hospitalisations reported in 2021 was lower on average than in the most recent pre-pandemic years (2017–2019), resulting in a relative decrease of 28.5% for outbreaks (5,601 outbreaks in 2017–2019, on average), 34.2% for cases (49,444 cases reported in 2017–2019, on average) and 44.3% for hospitalisations (4,482 hospitalisations in 2017–2019, on average). The number of reported deaths decreased compared with both 2020 and the period 2017–2019, by 8.8% (34 deaths in 2020) and 30.1% (44 deaths/year in 2017–2019, on average), respectively.
- In 2021, the foodborne outbreak reporting rate in the EU was 0.89% per 100,000 population. This represents an increase of 29.0% compared with 2020 (0.69 per 100.000 population) and a decrease of 18.3% compared with the pre-pandemic years (1.09 per 100.000 population in 2017–2019).
- These findings suggest that the impact of the COVID-19 pandemic was still considerable in 2021 in the EU, in relation to both the occurrence of foodborne outbreaks and their detection, investigation and reporting. However, the rise in the number of foodborne outbreaks, compared with 2020, might indicate a progressive return of foodborne outbreak surveillance to pre-pandemic stability for most MSs, although higher consumer exposure to contaminated foods may have contributed to this rise in 2021.
- The fall in foodborne outbreaks in the second year of the COVID-19 pandemic compared with pre-pandemic levels, did not affect all causative agents equally. For example, the number of outbreaks caused by Listeria monocytogenes was the highest since EFSA started collecting data.
- Salmonella was identified as causative agent in most foodborne outbreaks in the EU (N = 773), making up 19.3% of total outbreaks. This pathogen was also associated with the highest number of cases (20.8% of outbreak-associated cases) and hospitalisations (45.0% of outbreak-associated hospitalisations). S. Enteritidis was the predominant serovar (N = 350; 79.7% of all Salmonella outbreaks).
- A total of 31 deaths among outbreak cases were notified to EFSA by six MSs and the United Kingdom (Northern Ireland). *L. monocytogenes* was associated with 12 of these deaths (36.4% of all deaths), while *Campylobacter* was associated with six (18.2% of all deaths).
- A total of 355 strong-evidence outbreaks were reported in 2021 (8.9% of all outbreaks). Among these, food vehicles of animal origin (i.e. 'meat and meat products', 'fish and fishery products', 'eggs and egg products' and 'milk and milk products') were implicated in most FBOs (56.9%). Salmonella was the most frequent agent paired with various food items (e.g. eggs and egg products, mixed foods, vegetables and juices), in terms of the number of outbreaks, human cases and hospitalisations.
- 'Composite foods or multi-ingredient foods' caused the highest number of cases in strongevidence outbreaks (39.3% of all cases), with 'mixed foods' being the most frequently reported cause. Outbreaks implicating these foods were associated with a wide range of causative agents.

18314732, 2022, 12, Downloaded from https://efs.aonlinelibrary.wiley.com/doi/10.2903/jefs.a.2022.7666 by Istituto Zopprofilative Sperimenta dell'Umbria e delle Marche (ISSUM), Wiley Online library.wiley.com/rems-and-conditions) on Wiley Online library for rules of use. O. A articles are governed by the applicable Creative Common

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (thps://onlinelibrary.wiley.com/terms

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons

- Outbreaks associated with the consumption of 'vegetables and juices and other products thereof' rose considerably compared with both 2020 and the pre-pandemic years. This foodstuff was second only to 'mixed food' in the overall number of cases reported in FBOs in 2021.
- As in recent years, most outbreaks identified in 2021 took place on domestic premises (121 outbreaks; 34.1%. of all strong-evidence outbreaks). 'Restaurant or cafe or pub or bar or hotel or catering service' were the most frequently reported places of exposure for outbreaks occurring in other types of setting (77 outbreaks; 21.7%). 'School or kindergarten' was the place of exposure associated with the highest number of FBO cases (2,104 cases; 30.0% of all cases in strong-evidence outbreaks). These findings highlight the importance of proper implementation of HACCP in public catering as well as the need to improve the awareness of both consumers and food business operators on correct procedures for handling and consuming food.
- Alongside the present report, data on foodborne outbreaks may be explored through the EFSA interactive communication tools on foodborne outbreaks: the story map (available here), addressed to the general public, and the dashboard (available here) for an interactive data visualisation.

# 2. Surveillance and monitoring of foodborne outbreaks in the EU

Every year, EU Member States (MSs) and non-MSs report to EFSA data concerning the foodborne outbreaks (FBOs) occurring in their country, in compliance with Directive 2003/99/EC. The reporting of data is based on the standard set out in the guidance documents, published annually by EFSA (EFSA, 2021a,b).

EFSA is assigned the task of describing the causative agents and foodstuffs implicated in FBOs, along with their time trends. The aim is to assess the health impact of FBOs in Europe and to characterise the food production and distribution chains most frequently implicated. Outbreaks are categorised as having 'strong evidence' or 'weak evidence' based on the strength of evidence implicating a suspected food vehicle as the cause of the outbreak (EFSA, 2014). Data on the circumstances and risk factors contributing to consumer exposure to contaminated food are also analysed. The analysis takes account of any uncertainty around the evidence implicating a given food as the vehicle of the outbreak, by looking only at strong-evidence outbreaks for some analyses.

The current data reporting system is known as the European Union Foodborne Reporting System (EU-FORS). It applies to outbreaks caused by bacteria, viruses, parasites, algae, fungi and their products, such as toxins and biological amines (e.g. histamine), either typical foodborne agents or agents for which foodborne transmission is usually accidental. Outbreaks caused by the ingestion of drinking water are also considered in FBO reporting, since drinking water is defined as a food in Regulation (EC) No. 178/2002.

FBO data reporting is mandatory for EU MSs and the key findings are summarised in this report.

A description of the national systems in place for foodborne outbreak surveillance and reporting can be found in the national zoonoses reports submitted in accordance with Directive 2003/99/EC, which are published on the EFSA website and available here.

More details on the surveillance and monitoring of foodborne outbreaks in the EU are available on the EFSA story map (available here; see sections on 'What foodborne outbreaks are and how they are classified', 'Who investigates foodborne outbreaks' and 'EU regulatory framework and the role of EFSA').

# 3. Data analyses

Basic indicators used to describe the impact of FBOs on human health include the total number of outbreaks and cases, the number and proportion of cases (%) leading to hospitalisations or deaths, the mean outbreak size (cases per outbreak) and the range of cases per outbreak (minimum and maximum). Outbreak and case reporting rates (per 100,000 population) are used as relative measures of occurrence in the population, allowing a direct comparison among MSs, independently of the size of the population and any variations over time. However, due to the lack of full harmonisation of FBO surveillance among MSs, any direct comparison of findings between countries should be interpreted with caution. The case reporting rate is a new measure of the impact of FBOs, introduced in this report for the first time. Counts of hospitalisations and deaths and the proportion (%) of hospitalisations and deaths among outbreak cases are used as indicators for outbreak severity, However, since the total number of hospitalisations and deaths can be reported as 'unknown' by MSs, these values might be underestimated and should be analysed with caution.



The causative agents in FBOs in 2021 have been grouped in accordance with the following criteria:

- 'E. coli other than STEC' includes any pathogenic E. coli other than 'Shiga Toxin-producing E. coli (STEC)'. In 2021, this group included 'Enteroinvasive E. coli' (EIEC), 'Enterotoxigenic E. coli' (ETEC) and 'E. coli, unspecified'.
- 'Bacillus cereus toxins' includes 'B. cereus' and 'B. cereus enterotoxins'.
- 'Staphylococcus aureus toxins" includes 'S. aureus', 'Staphylococcus unspecified' and 'Staphylococcal enterotoxins'.
- 'Clostridium perfringens toxins' includes 'Clostridium, unspecified' and 'C. perfringens'.
- 'Norovirus (and other caliciviruses)' includes 'calicivirus, unspecified', 'norovirus' and 'sapovirus'.

To optimise the description of the findings and avoid data sparsity, information on 'food vehicle' and 'place of exposure' have been grouped, where necessary. Details concerning single entities in the group are described in the footnotes to the graphs or tables. Food vehicles have been grouped according to the general criteria set out in the EFSA data catalogues (EFSA, 2022). Places of exposure have been grouped according to the general characteristics and level of risk associated with the setting, as well as the process behind food preparation. For more information concerning the grouping criteria for causative agents, food vehicles and places of exposure, see 'Addendum foodborne outbreaks, 2021' from Section 1 to Section 3, published on the EFSA Knowledge Junction on Zenodo here.

Key statistics at EU level included the United Kingdom for 2019 and previous years since it was a MS. For 2021, according to the Protocol on Ireland/Northern Ireland, FBOs reported by the United Kingdom (Northern Ireland) were included in EU statistics.

Eurostat data on the resident population (updated on 1 January 2022) were used to calculate reporting rates at national and EU level, while data from the NISRA<sup>34</sup> Census 2021 were used for the United Kingdom (Northern Ireland).

Short-term variations over time were described using three main time frames to emphasise the impact of the COVID-19 pandemic on data reporting. In the main tables displaying key statistics at EU level (Tables 61, 62, 63 and 68), FBO and case reporting rates were described separately for 2021, 2020 and 2017–2019 (annual mean). The purpose was to enable an easier comparison between the trends of 2021, the first year of the pandemic and the pre-pandemic period.

Long-term variations were also described, taking 2012–2021 as the reference time period. Frequency distributions and trends are shown at EU level. Trends were analysed only at MS level, according to the rationale described in Boelaert et al. (2016) for data quality (Table 1).

Key statistics on FBOs for 2020 and previous years may differ from those published in the European Union One Health 2020 Zoonoses Report, as MSs may have updated their data from previous years.

Time trends were tested for statistical significance over the period 2012–2021 using the Cox-Stuart sign test, a nonparametric test appropriate for limited numbers of observations. A p-value of < 0.05 was considered to identify a statistically significant trend, beyond chance. The detection of significant trends at national level should be interpreted with caution, following changes in the reporting specifications for FBO, introduced in 2014 (EFSA, 2014). MSs with incomplete datasets for the 2012–2021 period were excluded from the trend analysis.

## 4. Results and discussion

# 4.1. Overview of countries reporting foodborne outbreak data in 2021

In 2021, 27 MSs and the United Kingdom (Northern Ireland) reported a total of 4,005 FBOs with 32,543 human cases, 2,495 hospitalisations and 31 deaths. At the same time, seven non-MSs (Bosnia and Herzegovina, Iceland, Montenegro, Norway, Republic of North Macedonia, Serbia, Switzerland) reported 83 outbreaks, 1,270 human cases, 65 hospitalisations and two deaths (Table 61).

The total number of FBOs reported by each MS in 2021 varied substantially, with most outbreaks concerning a small number of MSs. Belgium, France, the Netherlands and Poland accounted for most of the total number of FBOs (2,970 outbreaks or 74.2% of the total number) and for the majority of human cases reported in the EU in 2021 (19,936 cases or 61.3% of the total number).

Table 61 shows the substantial variations among MSs with respect to the numbers of outbreaks and cases reported in 2021. Outbreak reporting rates (per 100,000 population) ranged from 0.01

<sup>&</sup>lt;sup>34</sup> Northern Ireland Statistics and Research Agency (NISRA) https://www.nisra.gov.uk/

(Romania) to 5.0 (Malta) outbreaks per 100,000 population, corresponding to a 504-fold variation. The variation for case reporting rates (per 100,000 population) was lower but still considerable, ranging from 0.14 (Slovenia) to 25.0 (Finland).

The 'mean outbreak size' (i.e. the mean number of cases per outbreak) and the 'range of cases per outbreak' complete the characterisation of the FBOs occurring in 2021 in MSs and non-MSs. Taken together, the indicators in Table 61 describe the considerable differences among countries in terms of the occurrence of FBOs and their impact on health. These differences are largely influenced by the characteristics of the surveillance put in place by the MSs, and in particular the type of FBOs and causative agents under surveillance (Figure 32).

In 2021, FBOs classified as 'general outbreaks' (N = 1,550; 38.7%) outnumbered 'household outbreaks' (N = 794; 19.8%) even though not all MSs reported 'household outbreaks'. Overall, 17 MSs (Austria, Belgium, Croatia, Estonia, France, Germany, Greece, Ireland, Italy, Malta, the Netherlands, Poland, Romania, Slovakia, Slovenia, Spain, Sweden) provided data on 'household outbreaks' (i.e. FBOs involving single households) in 2021. The reporting of 'household' outbreaks, most of which are small and occur in domestic settings (N = 745; 93.8% of household outbreaks), has a direct consequence on the total count of FBOs occurring in a country and on the relative measures of FBO occurrence. This explains why the MSs reporting 'household' outbreaks generally have higher FBO reporting rates than MSs that do not notify this type of FBO to EFSA.

The case reporting rate is a sensitive measure of the health impact of FBOs as it is largely influenced by the size of the outbreaks. In 2021, the MSs with the highest case reporting rates (i.e. Finland, Latvia and Denmark) did not report 'household' outbreaks but provided information on several large (i.e.  $\geq 50$  cases) or very large (i.e.  $\geq 100$  cases) outbreaks. Finland reported a single FBO caused by S. Typhimurium in pre-cut vegetables involving 728 cases, and four other large FBOs caused by norovirus, STEC and S. Typhimurium. In Latvia, norovirus was responsible for two large FBOs. Denmark also reported four large FBOs caused by norovirus, and two other large FBOs caused by Enteroinvasive E. Coli (EIEC) and S. Typhimurium, respectively, implicating food of non-animal origin.

Information regarding the number of FBOs and cases, hospitalisations and deaths at EU and single MS level during 2016–2021, can be viewed dynamically on the dashboard (available here, see page 'reporting countries' for information on FBOs by reporting country). In 2021, the severity of outbreaks did not significantly change compared with 2020. The proportion of hospitalisations and deaths at EU-level was 7.7% and 0.1%, respectively, based on the total number of outbreak cases. Six MSs (Austria, France, Germany, the Netherlands, Spain, Sweden) and the United Kingdom (Northern Ireland) and one non-MS (Switzerland) reported a number of deaths among FBO cases. The number of deaths in the EU did not substantially change compared with 2020 (N = 34), although the distribution among MSs showed a considerable imbalance, with France (17 deaths) accounting for over half of the total deaths reported in the EU. This is the highest number of deaths among FBO cases reported by a single MS in 1 year, since 2012. At EU level, 15 deaths – including 13 in France – were reported as general outbreaks in 'health care and residential facilities', thus emphasising the risks of foodborne hazards for vulnerable populations.

Strong-evidence outbreaks accounted for 355 FBOs (8.9% of all FBOs) and were reported by 21 MSs (all MSs except Bulgaria, Cyprus, Estonia, Ireland, Latvia, Slovenia, United Kingdom (Northern Ireland)). For five MSs (Finland, Lithuania, Luxembourg, Romania, Sweden) strong-evidence outbreaks accounted for more than a third of total reported FBOs, while for four MSs (Belgium, France, the Netherlands, Slovakia), this proportion did not exceed 10%.



Table 61: Number of foodborne outbreaks, human cases, hospitalisations and deaths, in reporting EU MSs and non-MS countries, 2021

		utbreak	s				Cases	Hospit	talised	Deaths					
Country	Total (Strong- evidence)	% of	Outbro	eaks pei	100,000		% of	Mean outbreak	Case	s per 10	0,000		% of		% of
	N	total	2021	2020	2017– 2019 <sup>(a)</sup> (mean)	N	total	size (N) and range (min-max)	2021 <sup>(a)</sup>	2020	2017– 2019 <sup>(b)</sup> (mean)	N	total	N	total
Austria	20 (4)	0.50	0.22	0.24	0.64	92	0.30	4.6 (2–31)	1.0	0.75	4.7	27	1.1	2	6.5
Belgium	547 (3)	13.7	4.7	2.9	3.7	2,070	6.4	3.8 (2–60)	17.9	10.9	17.8	78	3.1	0	_
Bulgaria	0 (–)	_	0	0.09	0.26	0	_	_	0	2.1	3.0	0	_	0	_
Croatia	9 (1)	0.20	0.22	0.32	1.0	102	0.30	11.3 (2–71)	2.5	2.3	11.4	4	0.20	0	_
Cyprus	0 (-)	_	0	0.11	0.15	0	_	_	0	1.4	2.3	0	_	0	_
Czechia	42 (7)	1.0	0.39	0.20	0.25	1,976	6.1	47.0 (6–230)	18.5	7.9	8.4	260	10.4	0	_
Denmark	63 (12)	1.6	1.1	0.60	1.0	1,257	3.9	20.0 (2–85)	21.5	25.1	27.2	155	6.2	0	_
Estonia	8 (0)	0.20	0.60	1.1	0.93	20	0.10	2.5 (2–4)	1.5	2.2	10.0	8	0.30	0	_
Finland	48 (20)	1.2	0.87	0.65	1.0	1,385	4.3	28.9 (2–728)	25.0	10.8	20.9	28	1.1	0	-
France	1,286 (93)	32.1	1.9	1.5	2.4	10,836	33.3	8.4 (2–329)	16.0	10.1	22.0	561	22.5	17	54.8
Germany <sup>(b)</sup>	168 (21)	4.2	0.20	0.23	0.48	1,179	3.6	7.1 (2–98)	1.4	1.4	2.7	196	7.9	2	6.5
Greece	6 (1)	0.10	0.06	0.04	0.06	55	0.20	9.2 (2–30)	0.52	1.7	3.1	5	0.20	0	_
Hungary	19 (6)	0.50	0.20	0.11	0.40	564	1.7	29.7 (2–104)	5.8	2.8	17.0	43	1.7	0	_
Ireland	6 (0)	0.10	0.12	0.46	0.47	73	0.20	12.2 (2–35)	1.5	0.97	2.4	1	0.04	0	_
Italy	94 (21)	2.3	0.16	0.12	0.20	1,142	3.5	12.1 (2–150)	1.9	0.92	1.8	115	4.6	0	_
Latvia	12 (0)	0.30	0.63	0.89	1.5	454	1.4	37.8 (10–102)	24.0	6.0	19.8	24	1.0	0	_
Lithuania	7 (5)	0.20	0.25	0.18	1.7	71	0.20	10.1 (2–43)	2.5	1.1	9.4	39	1.6	0	_
Luxembourg	1 (1)	0.02	0.16	0.16	0.11	3	0.01	3.0 (-)	0.47	0.32	0.40	0	_	0	_
Malta	26 (3)	0.60	5.0	4.9	9.0	77	0.20	3.0 (2–19)	14.9	33.8	50.0	8	0.30	0	_
Netherlands	838 (7)	20.9	4.8	3.2	4.2	3,517	10.8	4.2 (2–402)	20.1	11.0	17.1	21	0.80	4	12.9
Poland	299 (39)	7.5	0.79	0.40	1.3	3,513	10.8	11.7 (2–152)	9.3	3.2	14.9	509	20.4	0	_
Portugal	17 (4)	0.40	0.17	0.04	0.14	495	1.5	29.1 (2–68)	4.8	0.55	4.9	80	3.2	0	_
Romania	2 (2)	0.05	0.01	0.02	0.08	48	0.10	24 (4–44)	0.25	0.21	2.4	6	0.20	0	_
Slovakia	214 (17)	5.3	3.9	5.9	11.7	844	2.6	3.9 (2–41)	15.5	13.0	43.3	172	6.9	0	_
Slovenia	1 (0)	0.02	0.05	0	0.05	3	0.01	3.0 (-)	0.14	0	2.2	0	_	0	_



Country		0	utbreak	s				Cases	Hospitalised		Deaths				
	Total (Strong- evidence)	% of	Outbreaks per 100,000				% of	Mean outbreak	Cases per 100,000				% of		% of
	N	total	2021	2020	2017– 2019 <sup>(a)</sup> (mean)	N	total	size (N) and range (min-max)	2021 <sup>(a)</sup>	2020	2017– 2019 <sup>(b)</sup> (mean)	N	total	N	total
Spain	221 (64)	5.5	0.47	0.34	1.1	2,052	6.3	9.3 (2–114)	4.3	2.8	11.7	129	5.2	2	6.5
Sweden	49 (24)	1.2	0.47	0.48	1.6	698	2.1	14.2 (2–90)	6.7	9.0	25.0	12	0.50	1	3.2
United Kingdom (Northern Ireland) <sup>(c)</sup>	2 (0)	0.05	0.11	_	-	17	0.10	8.5 (7–10)	0.89	_	_	14	0.60	3	9.7
EU Total (27 + XI) <sup>(b)</sup>	4,005 (355)	100	0.89	0.69	1.1	32,543	100	8.1 (2–728)	7.3	4.5	9.7	2,495	100	31	100
Bosnia and Herzegovina	2 (0)	-	0.06	0.03	0.10	66	-	33.0 (10–56)	2.0	1.9	1.4	1	-	0	_
Iceland	6 (0)	_	1.6	0.27	1.6	47	_	7.8 (3–13)	12.7	12.4	50.5	4	_	0	_
Montenegro	6 (1)	_	0.97	0.32	1.1	43	_	7.2 (2–20)	6.9	2.6	21.8	1	_	0	_
Norway	25 (9)	_	0.46	0.43	0.79	327	_	13.1 (3–30)	6.1	9.3	26.6	4	_	0	_
Republic of North Macedonia	3 (2)	-	0.15	0.05	0.40	195	_	65.0 (12–93)	9.4	0.48	24.0	12	-	0	_
Serbia	4 (2)	_	0.06	0.13	0.74	47	_	11.8 (5–24)	0.68	0.77	7.9	3	_	0	_
Switzerland <sup>(b)</sup>	37 (12)	_	0.43	0.15	0.21	545	_	15.1 (2–126)	6.3	1.9	3.4	40	_	2	_

<sup>(</sup>a): Data on FBOs from the United Kingdom are taken into account for 2017–2019, because the United Kingdom was an EU MS, but it became a third country on 1 February 2020.

<sup>(</sup>b): For one outbreak, information on cases was not available.


<sup>(</sup>c): Data on FBOs from the United Kingdom (Northern Ireland) are taken into account for 2021. In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/Northern Ireland, the EU requirements on data sampling are also applicable to Northern Ireland.

18314732, 2022, 12, Downloaded from https:

.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilattico

dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.c

-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commo



- \* 2021/2020 relative variation for Portugal was 325% and for Iceland was 500%.
- \*\* The mean annual total of FBOs reported by Bosnia and Herzegovina was calculated considering only the years 2017–2018, since no FBO data were reported in 2019.

**Figure 30:** Relative variations (%) in foodborne outbreaks reported in 2021 compared with 2020, and mean annual totals for the pre-COVID-19 pandemic period 2017–2019, by reporting country


In 2021, although the number of FBOs reported in the EU was higher than in 2020, it did not generally return to the level of pre-pandemic years (Figure 30). EU MSs reported overall 919 FBOs more than in 2020 (a 29.8% relative increase) but 1,596 fewer than the mean annual total for the period 2017–2019 (a 28.5% relative decrease). A similar pattern can be seen for cases of illness and hospitalisation, with 12,526 cases and 820 hospitalisations more than in 2020 (a relative increase of 62.6% and 49.0%, respectively) and 16,901 cases and 1,987 hospitalisations fewer, on average, than in 2017–2019 (a relative decrease of 34.2% and 44.3%). The number of deaths decreased both compared with 2020 (three fewer deaths; a relative decrease of 8.8%) and the pre-pandemic years (13 fewer deaths; a 30.1% relative decrease compared with 2017–2019).

Details of the annual variations in FBOs reported for 2021 as compared with 2020 and the prepandemic years (mean total of FBOs per year for the period 2017–2019) are shown in Figure 30, by country. Details on relative variations by causative agent and country are available in the 'Addendum foodborne outbreaks, 2021', published on the EFSA Knowledge Junction on Zenodo here, in sections 4 and 5.

Eight MSs (Finland, France, Hungary, Italy, Lithuania, Malta, Poland, Spain) and four non-MSs (Bosnia and Herzegovina, Montenegro, Norway, Republic of North Macedonia) reported more FBOs in 2021 than in 2020 but fewer than in the pre-pandemic years 2017–2019, on average. Eleven MSs (Austria, Bulgaria, Croatia, Cyprus, Estonia, Germany, Ireland, Latvia, Romania, Slovakia, Sweden) and one non-MS (Serbia) reported fewer FBOs in 2021 on average than in both 2020 and the period 2017–2019. Taken together, these results suggest that the COVID-19 pandemic and the associated control measures continued to have a major impact in 2021 on the occurrence of FBOs and their reporting in European countries. Six MSs (Belgium, Czechia, Denmark, Greece, the Netherlands, Portugal) and two non-MSs (Iceland and Switzerland) reported more FBOs in 2021 than in 2020 and 2017–2019, on average. Both an improvement in surveillance sensitivity and an increase in FBOs taking place in a domestic setting were mentioned by some countries as a possible reason for this increase.




Over the long term (2012–2021), the number of FBOs reported annually to EFSA increased significantly for the Netherlands and Switzerland, while decreased significantly for Austria, Slovenia and Norway (Figure 31).



18314732, 2022, 12, Downloaded from https://elsa.onlinelibrary.wiley.com/io/i0.2003/j.elsa.2022.7666 by Islituto Zoporellatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18122023]. See the Terms and Conditions (theta/continentibrary.wiley.com/rems-and-conditions) on Wiley Online Library for rules of use; OA stateles are governed by the applicable Cerative Commons





1831/4732, 2022, 12, Downloaded from https://efs.aonlinelibrary.wiley.com/oi/0.02903/j.cfs.a.2022.7266 by Istituto Zopopfulation Speriments dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [1812/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/erms-and-conditions) on Wiley Online Library for rules of use; OA arches are governed by the applicable Creative Commons Licensea

Note: \* indicates countries with a statistically significant trend (p < 0.05) over the period.

Dark red and light red show strong- and weak-evidence outbreaks, respectively. Black dots and lines show FBO reporting rates. The dots, lines and secondary Y-axis in blue showing the outbreak reporting rates have been used for Latvia, Lithuania, Malta and Slovakia, in order to draw attention to a scale that is different to that of the other countries.

- (a): Data on FBOs from the United Kingdom (Northern Ireland) are taken into account for 2021. In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/Northern Ireland, the EU requirements on data sampling are also applicable to Northern Ireland
- (b): Data on FBOs from the United Kingdom are taken into account for 2017–2019, because the United Kingdom was an EU MS, but it became a third country on 1 February 2020.

**Figure 31:** Trends in the number of strong-evidence and weak-evidence outbreaks (left axis) and the outbreak reporting rate (per 100,000) (right axis) in the EU and reporting EU MSs and non-MS countries, 2012–2021

## 4.2. Overview of causative agents in foodborne outbreaks in 2021

In 2021, a causative agent was identified in 2,174 FBOs (54.3% of the total) reported in the EU, causing 22,381 cases (68.8% of the total), 2,146 hospitalisations (86.0% of the total) and 28 deaths (90.3% of the total).

Table 62 shows the key statistics for FBOs reported in the EU, by causative agent. More information on the agents most frequently implicated in FBOs are shown on the dashboard (available here, see the page on 'causative agents' for more information on FBOs per causative agent).

Most outbreaks implicated bacteria, which were also responsible for the highest number of cases, hospitalisations and deaths. *Salmonella* was responsible for the majority of FBOs (about one in five), human cases and hospitalisations, by far. *Listeria monocytogenes* was associated with the highest number of deaths. After bacteria, the most frequently reported causative agents were bacterial toxins and viruses, in particular norovirus. Parasites and other causative agents including histamine were much less frequently reported. It is noteworthy that in 2021, reports listed one severe FBO caused by *Cronobacter sakazakii* and one by non-toxigenic *Vibrio cholerae*, a first since EFSA began collecting data on FBOs.

The breakdown of causative agents by country is shown in Figure 32. The MSs of Belgium, Denmark, Finland, France, Germany, Italy and Spain and one non-MS (Switzerland) were the countries with the highest number (more than 10) and variety of causative agents reported in FBOs in 2021. Salmonella, Campylobacter, norovirus (and other calicivirus), Staphylococcus aureus toxins and Clostridium perfringens toxins were the pathogens reported by most countries.

For a further interactive look at FBO data: see the dashboard here (different filters can be applied; outbreaks by causative agent are shown on different pages of the dashboard).

See the EFSA story map (available here) on FBOs, section on 'what organisms and symptoms'.

### 4.2.1. Bacteria

### Salmonella

In 2021, 26 MSs and six non-MSs (Table 61 and Figure 32) reported FBOs caused by Salmonella. France (N = 176), Poland (N = 165), Slovakia (N = 154) and Spain (N = 93) were the main contributors. In 15 MSs (Austria, Belgium, Croatia, Estonia, Greece, Hungary, Italy, Lithuania, Luxembourg, the Netherlands, Poland, Romania, Slovakia, Slovenia, Spain) and the United Kingdom (Northern Ireland), and in five non-MSs (Iceland, Montenegro, Republic of North Macedonia, Serbia, Switzerland), Salmonella was either the leading cause or the only cause of FBOs.

Information on *Salmonella* serovars was available for 439 FBOs (56.8% of all *Salmonella* FBOs). **S. Enteritidis** was the predominant serovar (N = 350; 79.7%), followed by **S. Typhimurium** (N = 50; 11.4%), **S. Braenderup** (N = 9; 2.1%) and **S. Typhimurium monophasic** (N = 6; 1.4%).

The number of *Salmonella* FBOs in the EU fell considerably in 2021 (N = 773) and 2020 (N = 694) compared with the pre-pandemic years (N = 1,371, on average), after several years of relative stagnation. This decline can be primarily attributed to the impact of the COVID-19 pandemic and was mainly driven by a decrease in the number of FBOs caused by S. Enteritidis, S. Typhimurium and S. Typhimurium monophasic, which together were responsible for 92.5% of all *Salmonella* FBOs in

18314732, 2022, 12, Downloaded from https://fs.aonlinelibrary.wiley.com/continelibrary.wiley.com

2021, with information on the serovar provided. The number of FBOs caused by **S. Enteritidis** gradually fell in 2020 and 2021, accounting for 452 and 504 fewer FBOs than over the period 2017–2019 (N = 854 reported FBOs on average), respectively. Slovakia (N = 146) and Poland (N = 125) reported most of the FBOs caused by **S.** Enteritidis in 2021, accounting for 77.4% of the total FBOs linked to this serovar in the EU. The results of a source attribution and trend analysis exercise recently carried out in Europe as part of the DiSCoVeR project showed an increased number of *Salmonella* outbreaks in the pre-pandemic years, which was mostly driven by **S.** Enteritidis outbreaks identified in Eastern Europe (Chanamé Pinedo et al., 2022). In 2020 and 2021, a relative drop could also be seen for FBOs caused by **S. Typhimurium** and **S. Typhimurium monophasic**, with 60 and 55 fewer reported FBOs in 2020 and 2021 than for the period 2017–2019 (N = 111 reported FBOs on average), respectively.

Over the same period, a similar relative decrease was observed for the number of cases and hospitalisations arising from FBOs caused by the main *Salmonella* serovars except *S.* **Typhimurium**. In 2021, this serovar caused 1,222 cases (1,062 cases more than in 2020; a 663.8% increase) mainly due to a single FBO in Finland with 728 cases. The outbreak (a strong-evidence FBO) was associated with the consumption of pre-cut vegetables and was not only the largest FBO reported in 2021 in the EU, but also the largest FBO caused by *S.* Typhimurium ever reported since data collection on FBOs began in 2004.

In 2021, a major outbreak caused by **S. Braenderup** leading to hundreds of cases was reported by multiple European countries (Austria, Belgium, Denmark, France, Luxembourg, the Netherlands and Switzerland). This FBO is noteworthy since this is a serovar rarely reported in outbreaks. According to the ECDC and EFSA rapid outbreak assessment (ECDC and EFSA, 2021a), this multi-country outbreak was probably associated with the consumption of contaminated 'Galia melons' imported from Honduras. More information on this outbreak is available in the 'ECDC and EFSA rapid outbreak assessment of multi-country foodborne outbreaks' section (see Section 4.8. ECDC and EFSA rapid outbreak assessment of multi-country foodborne outbreaks).

Altogether, the other 17 Salmonella serovars were responsible for 24 outbreaks (5.5% of the total with known serovars). They included S. Agona, S. Bareilly, S. Bovismorbificans, S. Chester, S. Coeln, S. Corvallis, S. Haifa, S. Infantis, S. Kedougou, S. Litchfield, S. Mikawasima, S. Montevideo, S. Muenchen, S. Oranienburg, S. Stanley, S. Strathcona and S. Virchow. Cases of infection caused by S. Chester and S. Montevideo were reported as being 'Part of a multicountry outbreak'.

# Campylobacter

In 2021, Campylobacter was the fourth most frequently reported causative agent in FBOs in the EU (Table 62). It was reported by 17 MSs (Austria, Belgium, Croatia, Czechia, Denmark, Estonia, Finland, France, Germany, Italy, Lithuania, Malta, the Netherlands, Poland, Slovakia, Spain and Sweden) and three non-MSs (Norway, Serbia, Switzerland) (Figure 32). Germany (N = 64), France (N = 55) and Slovakia (N = 55) reported most of these FBOs, making up 69.9% of the total. Campylobacter was the main causative agent in two MSs (Germany, Malta) and in Norway. Of the 112 FBOs reported by MSs with known Campylobacter species, C. jejuni was the causative agent in 106 FBOs (94.6%) and C. coli in six (5.4%). For 137 FBOs, species information was missing. Campylobacter FBOs are generally medium or small size outbreaks characterised by relatively mild illnesses. However, six severe outbreaks leading to deaths or a high proportion of hospitalisations were reported in 2021 by four MSs (Denmark, France, Spain, Sweden). The number of deaths observed in 2021 was the highest ever reported involving Campylobacter FBOs since 2007. Nevertheless, the number of FBOs and cases fell gradually in both 2020 and 2021. In 2021, the number of FBOs and cases reported in the EU decreased by 21.5% and 20.3% respectively, compared with 2020 (68 fewer outbreaks and 268 fewer cases). Campylobacter FBOs and cases halved compared with the pre-pandemic period of 2017-2019 on average (242 and 1,530 fewer FBOs and cases respectively).

18314732, 2022, 12, Downloaded from https://fs.aonlinelibrary.wiley.com/inchibrary.wiley.com/inchibrary.wiley.com/em-and-conditions) on Wiley Online Library on [18/12/2023], See the Terms and Conditions (http://oinineibrary.wiley.com/em-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons



Table 62: Number of foodborne outbreaks, human cases, hospitalisations and deaths, by causative agents, in reporting EU MSs, 2021

Outbre							Cases of illness							
Type of agent		Total (Strong- evidence)		Repor	ting rate	per 100,000	Human	cases	Mean outbreak size (cases)	Hospit	alisations	[	Deaths	
		N	<b>total</b> <sup>(a)</sup>	2021 <sup>(b)</sup>	<b>2020</b>	2017–2019 <sup>(c)</sup> (Mean) < 0.01	<b>N</b>	% of total <sup>(a)</sup>	and range (min-max)	N	% of cases <sup>(d)</sup>	N	% of cases <sup>(d)</sup>	
Bacteria	Aeromonas	1 (1) 1 (0)		< 0.01				0.10	19.0 (–)	0	0	0	0	
	Brucella	1 (0)	0.02	< 0.01	< 0.01	< 0.01	2	0.01	2.0 (-)	2	100	0	0	
	Campylobacter <sup>(e)</sup>	249 (20)	6.2	0.06	0.07	0.10	1,051	3.2	4.2 (2–39)	134	12.7	6	0.57	
	Cronobacter sakazakii	1 (1)	0.02	< 0.01	0	< 0.01	4	0.01	4.0 (-)	4	100	1	25.0	
	Escherichia coli other than STEC	27 (4)	0.70	0.01	< 0.01	< 0.01	327	1.0	12.1 (2–85)	44	13.5	0	0	
	Listeria monocytogenes	23 (8)	0.60	0.01	< 0.01	< 0.01	104	0.30	4.5 (2–11)	48	46.2	12	11.5	
	Salmonella	773 (143)	19.3	0.17	0.16	0.27	6,755	20.8	8.7 (2–728)	1,123	16.6	1	0.01	
	Shiga toxin- producing <i>E. coli</i> (STEC)	31 (5)	0.80	0.01	0.01	0.01	275	0.80	8.9 (2–76)	47	17.1	0	0	
	Shigella	11 (1)	0.30	< 0.01	< 0.01	0.01	63	0.20	5.7 (2–21)	4	6.3	0	0	
	Vibrio cholera (non- toxigenic)	1 (1)	0.02	< 0.01		< 0.01	47	0.10	47.0 (-)	1	2.1	0	0	
	Vibrio parahaemolyticus	3 (1)	0.10	< 0.01	< 0.01	< 0.01	10	0.03	3.3 (2–6)	0	0	0	0	
	Yersinia	21 (4)	0.50	< 0.01	< 0.01	< 0.01	125	0.40	6.0 (2–26)	14	11.2	0	0	
	Other bacteria/ unspecified	1 (0)	0.02	< 0.01	< 0.01	< 0.01	16	0.05	16.0 (-)	0	0	0	0	
	Subtotal <sup>(e)</sup>	1,143 (189)	28.5	0.25	0.24	0.39	8,798	27.0	7.7 (2–728)	1,421	16.2	20	0.23	
Bacterial toxins	Bacillus cereus toxins	87 (15)	2.2	0.02	0.02	0.04	679	2.1	7.8 (2–93)	9	1.3	1	0.15	
	Clostridium botulinum toxins	7 (4)	0.20	< 0.01	< 0.01	< 0.01	24	0.10	3.4 (2–8)	15	62.5	0	0	
	Clostridium perfringens toxins	40 (20)	1.0	0.01	0.01	0.02	778	2.4	19.5 (2–69)	25	3.2	4	0.51	



				Outbreak	(S		Cases of illness							
Type of agent		Total (Strong- evidence)	% of	Repor	ting rate	per 100,000	Human (	cases	Mean outbreak size (cases)	Hospit	alisations	C	Deaths	
		N	total <sup>(a)</sup>	2021 <sup>(b)</sup>	2020	2017–2019 <sup>(c)</sup> (Mean)	N	% of total <sup>(a)</sup>	and range (min-max)	N	% of cases <sup>(d)</sup>	N	% of cases <sup>(d)</sup>	
	Staphylococcus aureus toxins	61 (20)	1.5	0.01	0.01	0.04	640	2.0	10.5 (2–62)	51	8.0	0	0	
	Bacterial toxins, unspecified	484 (13)	12.1	0.11	0.08	0.13	4,257	13.1	8.8 (2–329)	210	4.9	2	0.05	
	Subtotal	679 (72)	17.0	0.15	0.12	0.18	6,378	19.6	9.4 (2–329)	310	4.9	7	0.11	
Viruses	Adenovirus	1 (0)	0.02	< 0.01	0	< 0.01	2	0.01	2.0 (-)	0	0	0	0	
	Flavivirus (including tick-borne Encephalitis virus)	1 (1)	0.02	< 0.01	< 0.01	< 0.01	5	0.02	5.0 (-)	5	100	0	0	
	Hepatitis A	13 (0)	0.30	< 0.01	< 0.01	0.01	264	0.80	20.3 (2–199)	209	79.2	0	0	
	Hepatitis E	1 (0)	0.02	< 0.01	< 0.01	< 0.01	3	0.01	3.0 (-)	1	33.3	0	0	
	Norovirus (and other Calicivirus)	251 (41)	6.3	0.06	0.03	0.07	6,545	20.1	26.1 (2–230)	156	2.4	1	0.02	
	Other viruses, unspecified	4 (0)	0.10	< 0.01	< 0.01	0.01	23	0.10	5.8 (2–12)	2	8.7	0	0	
	Subtotal	271 (42)	6.8	0.06	0.03	0.10	6,842	21.0	25.2 (2–230)	373	5.5	1	0.01	
Parasites	Cryptosporidium	2 (1)	0.05	< 0.01	< 0.01	< 0.01	25	0.10	12.5 (2–23)	0	0	0	0	
	Giardia	5 (0)	0.10	< 0.01	< 0.01	< 0.01	11	0.03	2.2 (2–3)	0	0	0	0	
	Trichinella	1 (0)	0.02	< 0.01	< 0.01	< 0.01	2	0.01	2.0 (-)	0	0	0	0	
	Subtotal	8 (1)	0.20	< 0.01	< 0.01	0.01	38	0.10	4.8 (2–23)	0	0	0	0	
Other causative agents	Histamine and Scombrotoxin	47 (15)	1.2	0.01	0.01	0.02	209	0.60	4.4 (2–14)	16	7.7	0	0	
	Marine biotoxins	17 (5)	0.40	< 0.01	0.01	0.01	83	0.30	4.9 (2–38)	3	3.6	0	0	
	Mushroom toxins	6 (4)	0.10	< 0.01	0	< 0.01	23	0.10	3.8 (2–6)	21	91.3	0	0	
	Other agents (incl. unspecified)	3 (1)	0.10	< 0.01	< 0.01	< 0.01	10	0.03	3.3 (2–6)	2	20.0	0	0	
	Subtotal	73 (25)	1.8	0.02	0.02	0.03	325	1.0	4.5 (2–38)	42	12.9	0	0	



Type of agent  Unknown Unknown/ Unspecified				Outbreak	(S		Cases of illness							
		Total (Strong- evidence)	% of	Repor	ting rate	per 100,000	Human cases		Mean outbreak size (cases)	Hospitalisations		С	Deaths	
		N	total <sup>(a)</sup>	2021 <sup>(b)</sup>	2020	2017–2019 <sup>(c)</sup> (Mean)	N	% of total <sup>(a)</sup>	and range (min-max)	N	% of cases <sup>(d)</sup>	N	% of cases <sup>(d)</sup>	
		1,831 (26)	45.7	0.41	0.27	0.39	10,162	31.2	5.5 (2–402)	349	3.4	3	0.03	
EU Total <sup>(e)</sup>		4,005 (355)	100	0.89	0.69	1.1	32,543	100	8.1 (2–728)	2,495	7.7	31	0.10	

<sup>&#</sup>x27;Escherichia coli' other than STEC' includes Enteroinvasive Escherichia coli (EIEC) (5), Enteropathogenic Escherichia coli (EPEC) (2) and other unspecified Escherichia coli (20).

- (a): Percentage out of the total number of cases reported in the EU.
- (b): Data on FBOs from the United Kingdom (Northern Ireland) are taken into account for 2021. In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/Northern Ireland, the EU requirements on data sampling are also applicable to Northern Ireland.
- (c): Data on FBOs from the United Kingdom are taken into account for 2017–2019, because the United Kingdom was an EU MS, but it became a third country on 1 February 2020.
- (d): Percentage out of the total number of cases caused by the causative agent.
- (e): For one outbreak, information on cases was not available. This outbreak was excluded from the calculation of the mean outbreak size.

<sup>&#</sup>x27;Marine biotoxins' includes ciguatoxin (8) and other unspecified toxins (9).

<sup>&#</sup>x27;Other causative agents' includes atropine (1) lectin (2).



### Listeria monocytogenes

Key statistics on FBOs caused by *Listeria monocytogenes* are available in Table 62. In 2021, this agent was reported in FBOs by eight MSs (Austria, Denmark, Finland, France, Germany, Italy, the Netherlands, Sweden) and the United Kingdom (Northern Ireland) (Figure 32). As in recent years, *L. monocytogenes* was associated with one of the highest numbers of deaths among outbreak cases and case fatality rates. The number of FBOs reported for 2021 was the highest since EFSA first started collecting data even though the United Kingdom is no longer included in the analysis since it left the EU in 2020. However at EU level, fewer cases, hospitalisations and deaths were reported in 2021 compared with both 2020 and 2017–2019, on average. One possible reason for this could be an increased detection of small listeriosis outbreaks in the population. This could be related to more sensitive laboratory-based surveillance of listeriosis with the widespread implementation of fine-tuning characterisation methods, in particular Whole Genome Sequencing (WGS), for *L. monocytogenes* (EFSA BIOHAZ Panel, 2019). This is a promising perspective since WGS data collection in humans and food, along with interoperable systems to support surveillance of *L. monocytogenes* outbreaks, have been recently implemented at EU level under the coordination of ECDC and EFSA (EFSA et al., 2022d).

## Shiga toxin-producing E. coli (STEC)

In 2021, **Shiga toxin-producing** *E. coli* (STEC) was the third most frequently reported bacterial agent detected in FBOs in the EU (Table 62). STEC were reported by 10 MSs (Austria, Belgium, Denmark, Finland, Germany, Ireland, Malta, the Netherlands, Spain, Sweden) and two non-MSs (Norway, Switzerland) (Figure 32). STEC was confirmed as the leading causative agent of FBOs in Ireland. The number of FBOs and cases of illness caused by STEC has fallen gradually and steadily over the past 4 years. FBOs fell from 50 outbreaks reported in 2018 to 31 in 2021 and in parallel cases decreased from 390 in 2018 to 275 in 2021. Conversely, the number of hospitalisations and deaths did not show a similar variation over the same period. Information on the STEC serogroup was available for 22 FBOs reported by MSs (all countries except Germany, Malta and Spain) and two FBOs reported by one non-MS (Norway). Overall, the identified serogroups among STEC FBOs were **Q157** (nine outbreaks), **Q26** (six outbreaks), **Q103** (five outbreaks) and **Q12**, **Q145**, **Q146**, **Q91** (one outbreak each).

### Other bacteria

Among the FBOs caused by other bacteria (Table 62), the number of outbreaks caused by **E. coli other than STEC** rose considerably in 2021 compared with recent years. **Enteroinvasive E. coli (EIEC)** were reported in five FBOs in France and Denmark, leading to 159 cases and 27 hospitalisations. The largest outbreak was associated with two different EIEC strains (O136 and O96) and was identified in Denmark. It involved 85 cases with infections taking place around the country. The implicated food was spring onion. An **Enterotoxigenic E. coli (ETEC)** strain was reported in 2021 in two single FBOs reported by Finland (29 cases) and Denmark (16 cases), linked to the consumption of vegetables and multi-ingredient foods, respectively. Both ETEC and EIEC were previously reported as causative agents of FBOs only in 2016, 2017 and 2018. Unfortunately, for most of these FBOs no information was available on the characteristics of the *E. coli* (20 outbreaks).

Summary statistics on the FBOs caused by **Yersinia enterocolitica** are shown in Table 62. In 2021, this pathogen was detected in FBOs reported by 11 MSs (Belgium, Denmark, Estonia, Finland, France, Germany, Lithuania, the Netherlands, Slovakia, Spain, Sweden) and one non-MS (Norway) (Figure 32). For three FBOs, the serotype involved was 0:3. The number of FBOs reported in 2021 was relatively stable compared with 2020 and the preceding pre-pandemic years.

Six MSs (Denmark, France, Germany, Malta, Spain, Sweden) reported 11 FBOs caused by **Shigella**. **S. sonnei** was identified in six FBOs, while the information on species was missing for the remaining FBOs.

In 2021, for the first time since EFSA began collecting data in 2004, one FBO caused by *Cronobacter sakazakii* was reported. The outbreak notified by Germany involved four newborns (premature) and caused one death. The implicated food vehicle was a hospital-mixed probiotic formula for infants. This report is not surprising since one of the main routes of transmission of this pathogen is powdered infant formula (PIF), which is not a sterile product (Forsythe, 2015). Food safety criteria for *Cronobacter* spp. are set out in Reg. (EC) No 2073/2005 for infant formulae and dried dietary foods intended for infants below 6 months of age. This report highlights the importance of good hygiene practices in reconstituting the product (Forsythe, 2015).



**V. cholerae** (non-toxigenic) was also reported for the first time in 2021 as the causative agent of an FBO. The FBO was detected in Spain, in a 'residential institution (nursing home or prison or boarding school)' and was linked to the consumption of mixed food. Usually, non-toxigenic serogroups cause self-limiting gastroenteritis and have a limited enteroinvasive potential (Sack et al., 2004). In literature, human infection is mainly associated with the intake of contaminated water or food, such as bivalve molluscs (Sack et al., 2004).

Other bacterial agents causing FBOs in the EU in 2021 included **Brucella**, **Aeromonas** and **Vibrio parahaemoloyticus**.

### 4.2.2. Bacterial toxins

In 2021, 16 MSs (Belgium, Croatia, Czechia, Denmark, Finland, France, Germany, Greece, Hungary, Italy, Poland, Portugal, Romania, Slovakia, Spain, Sweden) and three non-MSs (Norway, Republic of North Macedonia, Switzerland) reported FBOs caused by bacterial toxins (Figure 32). Bacterial toxins were the leading cause of FBOs in France and, as in recent years, France was also by far the country reporting the most FBOs of this type in the EU (611 outbreaks; 90.0% of all FBOs caused by bacterial toxins). In France, three different major FBOs caused by bacterial toxins (unspecified) led to 708 cases and 97 hospitalisations following the consumption of food in 'Canteen or catering at workplace, school, etc.'. All the seven deaths in cases of FBOs caused by bacterial toxins took place in France and involved people living in 'Health care and residential facilities'.

Although the number of FBOs involving bacterial toxins increased in 2021 (152 FBOs more than in 2020), it was still lower on average than for the period 2017–2019 (242 fewer FBOs; a 26.3% relative fall compared with 2017–2019).

In 2021, information on the toxigenic bacteria involved in FBOs was available only for 195 outbreaks (28.0% of FBOs caused by bacterial toxins). Although FBOs implicating **Bacillus cereus** toxins were associated with the highest number of FBOs among bacterial toxins, **Clostridium perfringens** toxins caused the highest number of cases and deaths and **Staphylococcus aureus** toxins the highest number of hospitalisations (Table 62). **S. aureus** toxins were the leading causative agent of FBOs in Portugal, Romania and Republic of North Macedonia. The number of FBOs caused by **Clostridium botulinum** toxins did not substantially change, compared with recent years, with two fewer outbreaks being reported in 2021, compared with 2020.

## **4.2.3.** Viruses

### **Norovirus (and other calicivirus)**

In 2021, **norovirus** (and other calicivirus) were the third most frequently reported agent causing FBOs in the EU (Table 62), and were reported by 14 MSs (Belgium, Czechia, Denmark, Finland, France, Germany, Ireland, Italy, Latvia, the Netherlands, Poland, Portugal, Spain, Sweden) and two non-MSs (Norway, Switzerland) (Figure 32). Norovirus (and other calicivirus) were the leading causative agent in six MSs (Belgium, Czechia, Denmark, Finland, Latvia, Sweden). France contributed the most to the total number of FBOs with 112 outbreaks (43.6% of all FBOs caused by norovirus (and other calicivirus)). In 2021, the FBOs caused by norovirus had the highest mean outbreak size (cases per outbreak) with MSs reporting 24 large (≥ 50 cases) and 13 very large (≥ 100 cases) outbreaks.

At EU level, although the reporting of FBOs caused by norovirus (and other calicivirus) increased last year compared with 2020 (121 FBOs more than 2020), the number of FBOs was still below that of the pre-pandemic years 2017–2019, on average (107 fewer outbreaks; a 29.9% relative decrease). The number of cases of illness, hospitalisations and deaths followed a similar downward trend.

### **Hepatitis A**

Overall, six MSs (Czechia, Finland, Germany, the Netherlands, Poland, Sweden) notified EFSA of FBOs caused by **hepatitis A** in 2021 (Figure 32). None of the FBOs were strong-evidence outbreaks (Table 62). One strong-evidence outbreak was reported by Norway, involving 20 cases. This outbreak was associated with the consumption of 'fruit, berries and juices and other products thereof'. Czechia reported a major weak-evidence outbreak of hepatitis A, involving 199 cases, almost all of which required hospitalisation (N = 195). Mixed food was the suspected food vehicle. Other statistics on hepatitis A FBOs are shown in Table 62.

18314732, 2022, 12, Downloaded from https://fs.aonlinelibrary.wiley.com/inchibrary.wiley.com/inchibrary.wiley.com/em-and-conditions) on Wiley Online Library on [18/12/2023], See the Terms and Conditions (http://oinineibrary.wiley.com/em-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons

#### Other viruses

Within the EU, FBOs caused by **hepatitis E** virus have been reported continuously since the first report in 2017, albeit in small numbers. In 2021, hepatitis E FBOs were reported from Belgium and Switzerland (Figure 32). The Swiss FBO was the largest ever reported to EFSA (105 cases) and was particularly severe in terms of the number of hospitalisations (N = 29) and deaths (N = 2). The outbreak was declared to be linked to the hepatitis E virus subtype 3 h, which is highly prevalent in the Swiss pig population (Office fédérel de la santé publique, 2022). The FBO was classified as a weak-evidence outbreak and no information on the suspect food vehicle was provided.

Of the other viruses reported in FBOs in 2021 (Table 62), Slovakia reported a single strongevidence FBO caused by the **tick-borne encephalitis virus**, associated with the consumption of raw goat's milk, while Poland reported one FBO caused by **adenovirus**.

#### 4.2.4. Parasites

### Trichinella

The number of countries reporting FBOs caused by *Trichinella*, as well as the total number of outbreaks and cases of **trichinellosis** has fallen steadily across the EU in the last 10 years, reaching the lowest value ever reported in 2021, with only one FBO and two cases (Table 62) reported by France. The outbreak was a weak-evidence FBO and neither information on the *Trichinella* species nor details on the implicated 'meat and meat product' was available.

### Cryptosporidium

FBOs caused by *Cryptosporidium* were reported by two MSs (Malta and Sweden) in 2021, and accounted for two outbreaks (Table 62 and Figure 32). The outbreak notified by Sweden was the largest (23 cases) and was a strong-evidence FBO associated with the consumption of kale, a food that has been recurrently reported in various outbreaks in Sweden over the past year. The species involved was *C. parvum*. Kale is a cruciferous vegetable belonging to the *Brassicaceae* family (Šamec et al., 2019), which has gained great popularity in Europe due its nutritional properties, cheapness and high availability on the market (Kapusta-Duch et al., 2011; Šamec et al., 2019).

#### Other parasites


All FBOs caused by other parasites in 2021 (Table 62) were associated with *Giardia intestinalis* (*lamblia*) and were reported by Poland. No detail on the suspect food vehicles was available.

# 4.2.5. Other causative agents

This group of outbreaks includes mainly 'histamine and scombrotoxin', 'marine biotoxins', 'mushroom toxins' and a few other chemical agents of biological origin that may accidentally contaminate food or its ingredients (Table 62). The data collected on FBOs caused by 'other causative agents' are the least harmonised among MSs, since these agents are not regularly covered by national outbreak surveillance programmes. This type of food poisoning is therefore likely to be underestimated at EU level.

In 2021, FBOs caused by 'histamine and scombrotoxin' were reported by six EU MSs (Finland, France, Germany, Italy, Spain, Sweden) (Figure 32). Most of the cases concerned France (33 outbreaks; 70.2%). No substantial variations in reporting were observed, compared with recent years. 'Fish and fish products' were the food implicated in all strong-evidence FBOs caused by 'histamine and scombrotoxin', except one, which was linked to the consumption of 'cheese'. Fish and fish products and ripened cheeses are considered the most common sources of histamine intoxication (EFSA BIOHAZ Panel, 2011).





The table may be read by column (country) or by row (causative agent). The number at the end of each row is the number of countries reporting a given causative agent for outbreaks in 2021, while the number at the top of each column indicates the number of causative agents identified in outbreaks by a given country in 2021. 'Escherichia coli' other than STEC includes Enteroinvasive E. coli (EIEC), Enterotoxigenic E. coli (ETEC) and E. coli, unspecified. 'Bacillus toxins' include Bacillus cereus, Bacillus cereus enterotoxins. 'Staphylococcus aureus toxins' include staphylococcal enterotoxins. 'Norovirus (and other calicivirus)' include norovirus (Norwalk-like virus) and calicivirus

Figure 32: Overview of EU MSs and non-MS countries reporting data on foodborne outbreaks, by causative agent, 2021

unspecified. 'Marine biotoxins' include ciquatoxin and other unspecified marine toxins. 'Other agents' include atropine and lectin.



Two MSs (France and Spain) reported FBOs caused by **marine biotoxins** (Figure 32), which are mainly produced by algae or phytoplankton accumulating in fish and filter-feeding molluscan shellfish. **Ciguatoxin** was responsible for eight FBOs (nine in 2020 and 26 over the period 2017–2019, on average). The increasing threat of ciguatoxin has been highlighted by a recent publication under the EuroCigua Project, as findings confirmed the presence of toxin-producing microalgae (e.g. *Gambierdiscus*) and fish with ciguatoxins in Europe (Canals et al., 2021).

In 2021, **mushroom toxin** poisoning was reported as the cause of FBOs by three MSs (Italy, Poland, Spain) and by Switzerland (Table 62 and Figure 32).

Two weak-evidence FBOs caused by **lectin** poisoning were reported in 2021, by Denmark, while one strong-evidence FBO associated with **atropine** was notified by Italy. This last outbreak was linked to the consumption of vegetables. Atropine poisoning can occur following incidental ingestion of contaminated food if toxic plants enter the food chain during harvest or processing (Adamse et al., 2014).

### 4.2.6. Outbreaks caused by unknown/unspecified agents

In 2021, FBOs of unknown aetiology (Table 62) accounted for 45.7% of all outbreaks in the EU, and were reported by 18 MSs and six non-MSs (Figure 32). In six MSs, unknown agents were the most frequent category of causative agents of FBOs. These countries were: the Netherlands (N = 810; 96.7% of all FBOs reported by this country), Belgium (N = 512; 93.6%), Portugal (N = 10; 58.8%), Hungary (N = 9; 47.4%), Italy (N = 35; 37.2%) and Sweden (N = 12; 24.5%).

FBOs of unknown aetiology tend to be small outbreaks, supporting the hypothesis that these outbreaks occur mainly in confined environments such as domestic settings or small groups, where it is relatively easy to identify a link between cases. However, 22 large ( $\geq$  50 cases) FBOs were reported by 11 MSs (Croatia, Czechia, Denmark, France, Hungary, Italy, the Netherlands, Poland, Portugal, Spain, Sweden). Several reasons may explain the reporting of unknown/unspecified agents, including the late reporting of illness, failure to detect the causative agents in patients or food, the unavailability of clinical or food samples (e.g. leftovers), etc.

Short-term relative variations (%) in the number of FBOs reported in 2021 compared with 2020 and the pre-pandemic period (2017–2019), by causative agents and by countries, are shown in the 'Addendum foodborne outbreaks, 2021', published on the EFSA Knowledge Junction on Zenodo here, in sections 4 and 5.

## 4.3. Overview of food vehicles implicated in foodborne outbreaks

The description of the food vehicles most frequently implicated in FBOs provides useful indications on the sources to be targeted by control policies, either at primary production level or in the various food preparation sectors, in order to reduce the public health impact of FBOs.

## 4.3.1. Food vehicles implicated in strong-evidence outbreaks

The summary statistics described in this section refer only to strong-evidence FBOs, in order to minimise uncertainty around the reported findings. In 2021, 355 strong-evidence FBOs (8.9% of total FBOs) were reported by 21 MSs (all MSs except Bulgaria, Cyprus, Estonia, Ireland, Latvia, Slovenia). In addition, 26 strong-evidence FBOs were reported by five non-MSs (Montenegro, Norway, Republic of North Macedonia, Serbia, Switzerland). Details of the food vehicles implicated in strong-evidence FBOs in MSs and non-MSs can be interactively searched using the EFSA dashboard (available here, see page 'food vehicles' for information on the FBOs per implicated food vehicle).

Table 63 describes the food vehicles implicated in the FBOs reported by MSs in 2021. Overall, three MSs (France, Spain, Poland) accounted for 55.2% of all strong-evidence FBOs (N = 196) reported in the EU. The remaining outbreaks (N = 159) were notified by 18 MSs.

#### Food of animal origin

In 2021, FBOs in the EU were mainly associated with the consumption of foods of animal origin (i.e. 'meat and meat products', 'fish and fishery products', 'eggs and egg products' and 'milk and milk products'). Foods of animal origin were implicated in 202 outbreaks (56.9% of all strong-evidence FBOs), 2,221 cases (31.7%), 316 hospitalisations (42.6%) and 11 deaths (84.6%).

'Meat and meat products' were the most frequently reported food group in 2021 (Table 63). Sixteen MSs (Austria, Czechia, Denmark, Finland, France, Germany, Greece, Hungary, Italy, Lithuania,

18314732, 2022, 12, Downloaded from https://fs.aonlinelibrary.wiley.com/inchibrary.wiley.com/enthated (UZSUM), Wiley Online Library on [18/12/2023], See the Terms and Conditions (http://oininelibrary.wiley.com/em-sand-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Cerative Commons

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (thps://onlinelibrary.wiley.com/terms

-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons



Malta, the Netherlands, Poland, Slovakia, Spain, Sweden) and four non-MSs (Norway, Republic of North Macedonia, Serbia, Switzerland) notified EFSA of FBOs associated with this food. 'Pig meat and products thereof' ranked first among all the items included in this group. The causative agents associated with this vehicle were Salmonella (14 FBOs caused by S. Typhimurium monophasic, S. Enteritidis, S. Bovismorbificans, Salmonella unspecified), Clostridium perfringens toxins (three FBOs), Bacillus cereus toxins, Staphylococcus aureus toxins, Yersinia and unspecified bacterial toxins (one FBO each). In one FBO, the agent was unknown. 'Broiler meat (Gallus gallus) and products thereof' were implicated in more FBOs than in 2020 (14 FBOs more than in 2020) but fewer than the mean total of FBOs reported for the period 2017-2019 (5 FBOs fewer). Salmonella (10 FBOs including S. Enteritidis, S. Oranienburg, S. Typhimurium and Salmonella unspecified), Campylobacter (seven FBOs), Staphylococcus aureus toxins (two FBOs), Clostridium perfringens toxins and Listeria monocytogenes (one FBO each) were the agents identified in these FBOs. In 2021, 'broiler meat (Gallus gallus) and products thereof' ranked first among the food vehicles implicated in FBOs caused by Campylobacter. The number of FBOs associated with 'bovine meat and products thereof' was the highest for the last 5 years (seven FBOs more than in 2020 and two more than in the period 2017–2019, on average). The causative agents of these FBOs included Clostridium perfringens toxins (five FBOs), Campylobacter (three FBOs), Salmonella (three FBOs including S. Enteritidis and Salmonella unspecified) and STEC 0157 (two FBOs). In 2021, 'bovine meat and products thereof' ranked first among the food vehicles implicated in FBOs caused by STEC. 'Other, mixed and/or unspecified poultry meat and products thereof' and 'other or mixed red meat and products thereof' were each identified as the implicated vehicle in two FBOs, caused by Campylobacter (two FBOs), Listeria monocytogenes and Staphylococcus aureus toxins.

In 2021, 'fish and fishery products' were reported by 10 MSs (Croatia, Denmark, Finland, France, Germany, Italy, the Netherlands, Poland, Spain, Sweden) (Table 63). The number of FBOs associated with this foodstuff was lower than in 2020 (10 FBOs fewer) and also lower than the mean total of FBOs for the period 2017-2019 (83 FBOs fewer). Among the foods included in this group, 'fish and fish products' were the fourth most frequently reported vehicles in strong-evidence FBOs in the EU, with notifications by nine MSs (Denmark, Finland, France, Germany, Italy, the Netherlands, Poland, Spain, Sweden). Note that 'fish and fish products' had a considerable health impact in 2021, leading to 190 cases, 41 hospitalisations and four deaths, the highest number of deaths in strongevidence outbreaks. The agents implicated in these FBOs were histamine and scombrotoxin (14 FBOs), marine biotoxins (five FBOs), Listeria monocytogenes (four FBOs), unspecified bacterial toxins (three FBOs), Bacillus cereus toxins, Clostridium botulinum toxins, Salmonella and Staphylococcus aureus toxins (one FBO each). All the deaths concerned two listeriosis FBOs identified in the Netherlands and associated with the consumption of different smoked fishes (smoked salmon, eel and mackerel). In 2021, 'fish and fish products' ranked first among the food vehicles implicated in FBOs caused by Listeria monocytogens, histamine and scombrotoxin and marine biotoxins. 'Crustaceans, shellfish, molluscs and products thereof', were implicated in FBOs reported by five MSs (Croatia, France, Italy, Spain, Sweden). Norovirus (and other calicivirus) were the agents most frequently reported in these FBOs (19 FBOs), followed by Bacillus cereus toxins, Staphylococcus aureus toxins and Vibrio parahaemolyticus (one outbreak each). In three FBOs, the agent was unknown. In 2021, 'crustaceans, shellfish, molluscs and products thereof' ranked first among the food vehicles implicated in FBOs caused by norovirus (and other caliciviruses).

**'Eggs and egg products**' were the second food vehicle group most frequently implicated in FBOs in the EU in 2021 (Table 63), reported by eight MSs (Denmark, France, Hungary, Lithuania, Poland, Slovakia, Spain, Sweden) and one non-MS (Switzerland). Almost all the FBOs linked to 'eggs and egg products', were associated with *Salmonella* (39 outbreaks, including *S*. Enteritidis, *S*. Typhimurium, *S*. Oranienburg, *S*. Infantis and *Salmonella* unspecified). One FBO was caused by norovirus and involved patients in a 'residential institution (nursing home or prison or boarding school)', causing one death. For two FBOs the causative agent was unknown. In 2021, 'eggs and egg products' ranked first among the food vehicle implicated in FBOs caused by *Salmonella* spp., *S*. Typhimurium including the monophasic variant.

**'Milk and milk products'** were reported by seven MSs (Finland, France, Germany, Lithuania, Poland, Slovakia, Spain) and one non-MS (Norway). The number of FBOs notified to EFSA by MSs concerning the consumption of 'milk and milk products' was higher than in 2020 (12 FBOs more) but lower on average than during the period 2017–2019, (8 FBOs fewer). This increase can be attributed to the reporting of more FBOs associated with the consumption of **'cheese'** in 2021 than in 2020 (14 more FBOs). The number of FBOs associated with the consumption of 'cheese' reported in 2021 was



18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (thps://onlinelibrary.wiley.com/terms

-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons

even higher than in the pre-pandemic years on average (5 more FBOs). The causative agents of FBOs linked to cheese included *Salmonella* (eight FBOs, including *S*. Enteritidis and *Salmonella* unspecified), *Staphylococcus aureus* toxins (four FBOs), histamine and unspecified bacterial toxins (one FBO each). For four FBOs, the information was not available. FBOs caused by 'dairy products' were also associated with *Salmonella* (*S*. Enteritidis) and *Staphylococcus aureus* toxins. In addition, one severe FBO caused by *Cronobacter sakazakii* in contaminated probiotic formula caused four human cases, four hospitalisations and one death in Germany. Among the FBOs linked to the consumption of 'milk', three FBOs were associated with raw milk from cows (two FBOs) and from goats (one FBO), contaminated with either *Campylobacter*, STEC or tick-borne encephalitis virus (one outbreak each), consumed on the farm or in a domestic setting.

## Food of non-animal origin

In 2021, FBOs associated with the consumption of foods of non-animal origin were reported by 14 MSs (Austria, Belgium, Denmark, Finland, France, Germany, Hungary, Italy, Lithuania, Luxembourg, the Netherlands, Poland, Spain, Sweden) and two non-MSs (Switzerland, Norway). In 2021, the number of FBOs associated with foods of non-animal origin doubled compared with 2020 (22 FBOs more), almost reaching the mean annual total reported during the period 2017-2019 (5 FBOs fewer, on average) (Table 63). This increase was mainly driven by FBOs implicating 'vegetables and juices and other products thereof'. This food group was associated with the largest variety of causative agents. Among the bacteria: Salmonella (11 FBOs, including S. Braenderup, S. Typhimurium, S. Enteritidis, S. Coeln, S. Kedougou and Salmonella unspecified), Enteroinvasive E. coli (EIEC) (two FBOs), Shiga toxin-producing E. coli (one FBO) and Yersinia enterocolitica (two FBOs), Enterotoxigenic E. coli (ETEC) (one FBO); bacterial toxins including Bacillus cereus (two FBOs), Clostridium botulinum (two FBOs), Staphylococcus aureus (one FBO) and bacterial toxins, unspecified (two FBOs); mushroom toxins (four FBOs); viruses including Norovirus (and other calicivirus) (three FBOs): parasite: Cryptosporidium parvum (one FBO). Nine large (≥ 50 cases) or very large (≥ 100 cases) FBOs associated with 'vegetables and juices and other products thereof' were reported in 2021 in the EU. This explains why the mean size of the FBOs linked to this foodstuff (50 cases/outbreak) was one of the highest, outnumbering the FBOs associated with foods of animal origin (11 cases/outbreak). Multiple mechanisms leading to the primary contamination of crops and the cross-contamination of vegetables during processing have been described in literature. Pre-cut vegetables were implicated in five FBOs, all reported by Finland. Of this total, one was associated with S. Typhimurium, causing 728 cases. Of the single outbreaks linked to this product in 2021, we should mention the large FBO associated with 'Alfalfa sprouts' and caused by S. Coeln in Sweden. 'Galia melons' imported from Honduras were responsible for a large multi-country outbreak of infections by S. Braenderup (see Section 4.8. ECDC and EFSA rapid outbreak assessment of multi-country foodborne outbreaks). Other foods of non-animal origin implicated in FBOs in 2021 were 'cereal products including rice and seeds/pulses (nuts, almonds)', reported in outbreaks caused mainly by Bacillus cereus (four FBOs), Salmonella, unspecified (three FBOs), Clostridium botulinum (one FBO) and unspecified bacterial toxins (one FBO, each) and 'fruit, berries and juices and other products thereof', reported in outbreaks caused by Salmonella unspecified and Yersinia enterocolitica. In 2021, among non-MSs, Switzerland reported one large general outbreak caused by norovirus and associated with the consumption of berries and small fruits.

## Composite foods, multi-ingredient foods and other foods

This food group includes foods resulting from the assembly of multiple ingredients, or highly processed or manipulated foods. A range of foodstuffs belong to this category, namely **'bakery products'**, **'buffet meals'**, **'mixed foods'**, **'sweets and chocolate'** and **'other foods'**. Eighteen MSs and four non-MSs (Switzerland, Republic of North Macedonia, Montenegro, Norway) reported FBOs associated with **'composite foods, multi-ingredient foods and other'**. The number of FBOs reported was higher than in 2020 (50 FBOs more) but still lower than the total mean reported annually during the period 2017–2019 (85 fewer on average). As was the case for foods of non-animal origin, the mean size of FBOs implicating this food group was 26 cases/outbreak, larger than those associated with foods of animal origin.



**Table 63:** Frequency distribution of strong-evidence foodborne outbreaks, by food vehicle, in reporting EU MSs, 2021

			Stron	g-eviden	ce outb	reaks			Repo	rting rate	per 100,000	Rank	
Type of vehicle	Out	breaks	Ca	ises	Hospit	talisations	Deat	ths			2017 2010(b)		
Type of venicle	N	% of total	N	% of total	N	% of total	N	% of total	2021 <sup>(a)</sup>	2020	2017–2019 <sup>(b)</sup> (mean)	2021 <sup>(a)</sup>	2017– 2020 <sup>(b)</sup>
Composite foods, multi-ing	gredien	ts foods	and othe	r foods									
Mixed foods	76	21.4	2,156	30.8	181	24.4	2	15.4	0.017	0.006	0.017	1	3
Bakery products	18	5.1	258	3.7	52	7.0	0	0	0.004	0.002	0.008	8	5
Buffet meals	5	1.4	205	2.9	33	4.5	0	0	0.001	0.001	0.003	14	13
Sweets and chocolate	2	0.60	34	0.50	8	1.1	0	0	< 0.001	0.001	0.001	17	21
Other foods	5	1.4	103	1.5	2	0.30	0	0	0.001	0.002	0.007	14	7
Subtotal	106	29.9	2,756	39.3	276	37.2	2	15.4	0.024	0.013	0.037	_	-
Meat and meat products													
Pig meat and products thereof	22	6.2	347	5.0	69	9.3	3	23.1	0.005	0.004	0.007	6	6
Broiler meat ( <i>Gallus gallus</i> ) and products thereof	21	5.9	202	2.9	42	5.7	0	0	0.005	0.002	0.005	7	10
Meat and meat products, unspecified	17	4.8	237	3.4	18	2.4	2	15.4	0.004	0.002	0.007	10	8
Bovine meat and products thereof	13	3.7	201	2.9	12	1.6	0	0	0.003	0.001	0.002	11	17
Other or mixed red meat and products thereof	2	0.60	11	0.20	0	0	0	0	< 0.001	0.001	0.002	17	14
Other, mixed or unspecified poultry meat and products thereof	2	0.60	14	0.20	0	0	0	0	< 0.001	< 0.001	0.002	17	19
Subtotal	77	21.7	1,012	14.4	141	19	5	38.5	0.017	0.010	0.026	_	_
Fish and fishery products													
Fish and fish products	30	8.5	190	2.7	41	5.5	4	30.8	0.007	0.006	0.01	4	4
Crustaceans, shellfish, molluscs and products thereof	25	7.0	171	2.4	13	1.8	0	0	0.006	0.008	0.017	5	2
Subtotal	55	15.5	361	5.2	54	7.3	4	30.8	0.012	0.015	0.027	_	_



			Stron	g-eviden	ce outb	reaks			Repo	rting rate	per 100,000	Rank	
Type of vehicle	Out	breaks	Са	ises	Hospit	alisations	Deat	ths			2017 2010(b)		2017
Type of vehicle	N	% of total	N	% of total	N	% of total	N	% of total	2021 <sup>(a)</sup>	2020	2017–2019 <sup>(b)</sup> (mean)	2021 <sup>(a)</sup>	2017– 2020 <sup>(b)</sup>
Food of non-animal origin													
Vegetables and juices and products thereof	34	9.6	1,700	24.3	131	17.7	0	0	0.008	0.003	0.006	3	9
Cereal products including rice and seeds/pulses	9	2.5	194	2.8	17	2.3	0	0	0.002	0.001	0.002	12	18
Fruit, berries and juices and products thereof	2	0.60	15	0.20	0	0	0	0	< 0.001	0.001	0.002	17	20
Subtotal	45	12.7	1,909	27.3	148	20.0	0	0	0.010	0.005	0.01	_	_
Eggs and egg products	42	11.8	439	6.3	90	12.1	1	7.7	0.009	0.009	0.024	2	1
Milk and milk products													
Cheese	18	5.1	235	3.4	11	1.5	0	0	0.004	0.001	0.002	8	15
Dairy products (other than cheeses)	6	1.7	119	1.7	7	0.90	1	7.7	0.001	0.001	0.001	13	22
Milk	4	1.1	55	0.80	13	1.8	0	0	0.001	0.002	0.004	16	11
Subtotal	28	7.9	409	5.8	31	4.2	1	7.7	0.006	0.004	0.007	-	_
Water (and other beverage	es)												
Water	2	0.60	119	1.7	1	0.10	0	0	< 0.001	0.001	0.002	17	16
Subtotal	2	0.60	119	1.7	1	0.10	0	0	< 0.001	0.001	0.002	_	_
EU Total <sup>(b)</sup>	355	100	7,005	100	741	100	13	100	0.079	0.055	0.266	_	_

Note: Single food items are consolidated into major groups according to their origin. The 'Outbreak Reporting Rate' columns include the mean outbreak reporting rate per 100,000 for 2021 and for the previous years (2017–2020) for trend watching. The ranking of each food item provides a visual demonstration of the relative importance of the item, among all food vehicles implicated in foodborne outbreaks, for the same year and period.

<sup>(</sup>a): Data on FBOs from the United Kingdom (Northern Ireland) are taken into account for 2021. In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/Northern Ireland, the EU requirements on data sampling are also applicable to Northern Ireland.

<sup>(</sup>b): Data on FBOs from the United Kingdom are taken into account for 2017–2019, because the United Kingdom was an EU MS, but it became a third country on 1 February 2020.

'Bakery products' includes 'Bakery products' 'Bakery pro

<sup>&#</sup>x27;Bakery products' includes 'Bakery products', 'Bakery products – cakes', 'Bakery products – cakes – containing heat-treated cream', 'Bakery products – desserts – containing raw eggs', 'Bakery products – pastry'.

<sup>&#</sup>x27;Bovine meat and products thereof' include 'Bovine meat and products thereof', 'Cooked cured (or seasoned) bovine meat', 'Meat from bovine animals – meat products', 'Meat from bovine animals – meat products – raw and intended to be eaten raw', 'Meat from bovine animals – minced meat'.

<sup>&#</sup>x27;Broiler meat (*Gallus gallus*) and products thereof' include 'Broiler meat (*Gallus gallus*) and products thereof', 'Meat from broilers (*Gallus gallus*)', 'Meat from broilers (*Gallus gallus*) – meat products'. 'Cheese' includes 'Cheese', 'Cheeses made from sheep's milk – fresh'.

<sup>&#</sup>x27;Crustaceans, shellfish, molluscs and products thereof' includes 'Crustaceans, shellfish, molluscs and products thereof', 'Molluscan shellfish'.



'Eggs and egg products' include 'Eggs', 'Eggs and egg products'.

'Fish and fish products' include 'Fish', 'Fish – cooked', 'Fish – Fishery products from fish species associated with a high amount of histidine – not enzyme maturated', 'Fish – raw', 'Fish – smoked', 'Fish and fish products', 'Roe – chilled'.

'Milk' includes 'Milk', 'Milk, cows' – raw milk', 'Milk, goats' – raw milk'.

'Mixed food' includes 'Foodstuffs intended for special nutritional uses – dietary foods for special medical purposes', 'Foodstuffs intended for special nutritional uses – other food for infants and children', 'Mixed food', 'Other processed food products and prepared dishes – fish and seafood based dishes', 'Other processed food products and prepared dishes – meat based dishes', 'Other processed food products and prepared dishes – mushroom based dishes', 'Other processed food products and prepared dishes – pasta based dishes', 'Other processed food products and prepared dishes – sandwiches – non-meat', 'Other processed food products and prepared dishes – sushi', 'Other processed food products and prepared dishes – unspecified', 'Ready-to-eat salads', 'Sauce and dressings', 'Sauce and dressings', 'Soups'.

Other or mixed red meat and products thereof includes 'Meat, mixed meat – meat products – ready-to-eat', 'Other or mixed red meat and products thereof'.

Other, mixed or unspecified poultry meat and products thereof includes 'Meat from duck – meat products – ready-to-eat', 'Meat from poultry, unspecified'.

'Pig meat and products thereof' includes 'Cooked cured (or seasoned) pork meat', 'Meat from pig – meat products', 'Meat from pig – meat products – meat specialities', 'Meat from pig – meat products – ready-to-eat', 'Meat from pig – offal', 'Pig meat and products thereof'.

'Sweets and chocolate' include 'Confectionery products and pastes - hard candy', 'Sweets and chocolate'.

'Vegetables and juices and other products thereof' include 'Alfalfa sprouts', 'Lettuce', 'Melons (except watermelon)', 'Mushrooms', 'Spring onion', 'Vegetables – pre-cut', 'Vegetables – products – cooked', 'Vegetables and juices and other products thereof'.

'Water' includes 'Tap water, including well water'.

In 2021, 'mixed foods' were the foodstuffs most frequently reported in FBOs. Food items in this group are highly heterogeneous. Details are provided in (Table 63). Many different causative agents were linked to 'mixed foods', mainly bacteria: Salmonella (24 FBOs, including S. Enteritidis, S. Typhimurium, S. Typhimurium monophasic, S. Corvallis, S. Bareilly, Salmonella, unspecified), Campylobacter (five FBOs), Aeromonas, E. coli unspecified, Enterotoxigenic E. coli (ETEC), Shigella, non-toxigenic Vibrio cholerae (one FBO, each); viruses: norovirus (13 FBOs); bacterial toxins: Bacillus cereus (six FBOs), Clostridium perfringens (seven FBOs), Staphylococcus aureus (seven FBOs) and unspecified (five FBOs). For five FBOs, the causative agent was unknown. Interestingly, the majority of FBOs were characterised as general outbreaks, accounting for 80.3% of all FBOs caused by 'mixed foods'. In 2021, mixed foods' ranked first among the food vehicles implicated in FBOs caused by S. Enteritidis, Bacillus cereus toxins, Staphylococcus aureus toxins and Clostridium perfringens toxins.

**'Bakery products'** were reported in FBOs mainly caused by *Salmonella* (15 FBOs, including *S*. Enteritidis and *Salmonella*, unknown). Norovirus (and other calicivirus) and unknown agents were also reported (one and two FBOs, respectively). **'Buffet meals'** were associated with *Salmonella* (two FBOs, including *S*. Typhimurium monophasic, *S*. Enteritidis), norovirus (two FBOs) and *Campylobacter* (one FBO). Finally, **'sweets and chocolate'** led to two FBOs caused by *S*. Enteritidis, while **'other foods'** were linked to FBOs caused by *S*. Enteritidis (one FBO) and *Clostridium perfringens* toxins (two FBOs) and an unknown agent (three FBOs).

For more information concerning the causative agent associated with the consumption of different types of food implicated in strong-evidence FBOs, see the dashboard (available here, see page 'Food vehicles and causative agents' for information on FBOs by food vehicle and by causative agent).

#### 4.3.2. Food vehicles implicated in weak-evidence outbreaks

In 2021, a total of 1,873 weak-evidence FBOs with a suspected food vehicle (46.8% of all FBOs) were reported by 20 MSs. No major discrepancies were observed in the ranking of food types for weak-evidence outbreaks, compared with strong-evidence outbreaks, in terms of causative agents, with a few exceptions. One of the main discrepancies observed in 2021 concerned STEC FBOs. Although 'bovine meat and products thereof' was the main food category reported in strong-evidence outbreaks (two strong-evidence outbreaks, one weak-evidence outbreak), 'water' was actually the most frequently suspected source, even though the evidence is weak (four weak-evidence outbreaks). This situation is consistent with the observations of previous years within the EU, and once again draws attention to environmental pathways in the transmission of STEC infections to humans.

Information regarding the foodstuffs implicated in strong- and weak-evidence foodborne outbreaks by reporting MSs in 2021, for each causative agent, are available on the dashboard (available here, see page 'Food vehicles and causative agents' for information on FBOs by food vehicle and by causative agent).

# 4.3.3. Top-10 agent/food pairs in strong-evidence outbreaks associated with the highest impact on health in the EU, 2021

Tables 64–67 show the top-10 pairs of causative agents and food vehicles for the strong-evidence outbreaks with the highest health impact in 2021 in the EU, in terms of total outbreaks, cases, hospitalisations and deaths, respectively. The number of MSs that reported FBOs implicating each food/agent pair is also included in the tables, indicating how common these types of outbreaks were in EU MSs. When looking at the tables, it is important to remember that the ranking of the pairs may be influenced by the scale of the contribution made by each MS to data collection. The same information is provided for the 2017–2019 period, for trend watching purposes.

18314732, 2022, 12, Downloaded from https://fs.aonlinelibrary.wiley.com/continelibrary.wiley.com



Table 64: Top 10 pathogen/food vehicle pairs causing the highest number of strong-evidence outbreaks in reporting EU MSs, 2021

		2	021 <sup>(a)</sup>			Evaluation		
Rank <sup>(c)</sup>	Causative agent	Food vehicle	Outbreaks (N)	Reporting MSs (N outbreaks)	Rank <sup>(c)</sup>	Outbreaks (N/year) (range) <sup>(d)</sup>	Reporting MSs (N/year) <sup>(d)</sup>	2021 vs. 2017-2020 <sup>(e)</sup>
1	Salmonella	Eggs and egg products <sup>(f)</sup>	39	Poland (13), Spain (12), France (8), Sweden (2), Slovakia (1), Hungary (1), Denmark (1), Lithuania (1)		97.2 (37–135)	10.0	<b>↓</b> ↓
2	Salmonella	Mixed foods <sup>(g)</sup>	24	Slovakia (10), Poland (3), Finland (3), Spain (3), Belgium (1), Denmark (1), Malta (1), Germany (1), Italy (1)	5	22.2 (3–34)	7.7	-
3	Norovirus (and other calicivirus)	Crustaceans, shellfish, molluscs and products thereof <sup>(h)</sup>	19	France (15), Sweden (2), Spain (1), Italy (1)	2	55.7 (8–144)	5.75	<b>↓</b> ↓
4	Salmonella	Bakery products <sup>(i)</sup>	15	Poland (7), Spain (3), Romania (1), Slovakia (1), Malta (1), Italy (1), Czechia (1)	3	28.7 (9–45)	2.7	<b>\</b>
5	Histamine and scombrotoxin	Fish and fish products <sup>(j)</sup>	14	Sweden (5), France (4), Italy (3), Finland (1), Germany (1)	4	28.2 (14–55)	7.2	$\downarrow\downarrow$
5	Salmonella	Pig meat and products thereof <sup>(k)</sup>	14	France (9), Germany (3), Italy (2)	6	16.5 (11–26)	7.7	-
6	Norovirus (and other calicivirus)	Mixed foods <sup>(I)</sup>	13	Finland (3), France (3), Spain (2), Sweden (2), Portugal (1), Belgium (1), Italy (1)	11	8.5 (5–11)	4.5	$\uparrow \uparrow$
7	Salmonella	Vegetables and juices and other products thereof <sup>(m)</sup>	11	Finland (3), Belgium (1), Netherlands (1), Luxembourg (1), Denmark (1), Spain (1), Sweden (1), Austria (1), Germany (1)	29	4.0 (1–6)	3.0	<b>↑</b> ↑
8	Salmonella	Broiler meat (Gallus gallus) and products thereof	10	Spain (3), Poland (2), France (2), Sweden (1), Hungary (1), Malta (1)	16	7.0 (2–13)	3.7	1
9	Salmonella	Cheese	8	France (7), Poland (1)	26	4.2 (1–8)	1.5	$\uparrow \uparrow$

<sup>(</sup>a): Data on FBOs from the United Kingdom (Northern Ireland) are taken into account for 2021. In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/Northern Ireland, the EU requirements on data sampling are also applicable to Northern Ireland.

<sup>(</sup>b): Data on FBOs from the United Kingdom are taken into account for 2017–2019, because the United Kingdom was an EU MS, but it became a third country on 1 February 2020.

<sup>(</sup>c): Ranking of the food vehicle based on the number of strong-evidence FBOs in which the combination (causative agent/food vehicle) was implicated (rank 1 is the highest rank, meaning the most commonly implicated). Strong-evidence outbreaks with unknown causative agents are not included.

<sup>(</sup>d): Outbreaks reported by the United Kingdom are also included.

<sup>(</sup>e): A single arrow indicates variations of 25% and 50% in the number of outbreaks; double arrows indicate variations > 50%; a 'stable' value indicates variations of between -25% and +25%.

<sup>(</sup>f): 'Eggs and egg products' include 'Eggs', 'Eggs and egg products'.



- (g): 'Mixed food' includes 'Foodstuffs intended for special nutritional uses dietary foods for special medical purposes', 'Mixed food', 'Other processed food products and prepared dishes fish and seafood-based dishes', 'Other processed food products and prepared dishes pasta', 'Other processed food products and prepared dishes pasta', 'Sauce and dressings mayonnaise'.
- (h): 'Crustaceans, shellfish, molluscs and products thereof' includes 'Crustaceans, shellfish, molluscs and products thereof', 'Molluscan shellfish'.
- (i): 'Bakery products' include 'Bakery products', 'Bakery products cakes', 'Bakery products cakes containing heat-treated cream', 'Bakery products desserts containing raw cream', 'Bakery products desserts containing raw eggs'.
- (j): 'Fish and fish products' include 'Fish', 'Fish cooked', 'Fish Fishery products from fish species associated with a high amount of histidine not enzyme maturated', 'Fish raw', 'Fish and fish products'.
- (k): 'Pig meat and products thereof' includes 'Meat from pig meat products', 'Meat from pig meat products ready-to-eat', 'Meat from pig offal', 'Pig meat and products thereof'.
- (I): 'Mixed food' includes 'Mixed food', 'Other processed food products and prepared dishes', 'Other processed food products and prepared dishes sushi', 'Ready-to-eat salads'.
- (m): 'Vegetables and juices and other products thereof' include 'Alfalfa sprouts', 'Melons (except watermelon)', 'Vegetables pre-cut', 'Vegetables and juices and other products thereof.

Table 65: Top-10 pathogen/food vehicle pairs causing the highest number of cases in strong-evidence outbreaks in reporting EU MSs, 2021

			2021 <sup>(a</sup>	)		2017–2020 <sup>(t</sup>	<b>)</b>	Evaluation
Rank <sup>(c)</sup>	Causative agent	Food vehicle	Cases (N)	Reporting MSs (N cases)	Rank <sup>(c)</sup>	Cases (N/year) (range) <sup>(d)</sup>	Reporting MSs (N/year) <sup>(d)</sup>	2021 vs. 2017–2020 <sup>(e)</sup>
1	Salmonella	Vegetables and juices and other products thereof <sup>(f)</sup>	1,103	Finland (797), Germany (82), Belgium (55), Sweden (53), Denmark (41), Netherlands (37), Spain (24), Austria (11), Luxembourg (3)	24	108.2 (7–244)	3.0	<b>↑</b> ↑
2	Norovirus (and other calicivirus)	Mixed foods <sup>(g)</sup>	487	France (227), Portugal (68), Italy (62), Finland (49), Sweden (44), Spain (31), Belgium (6)	7	377.7 (223–735)	4.5	<b>↑</b>
3	Salmonella	Mixed foods <sup>(h)</sup>	486	Slovakia (187), Spain (134), Denmark (52), Italy (37), Finland (25), Germany (19), Malta (19), Poland (9), Belgium (4)	2	777.8 (91–1,595)	8.0	<b>\</b>
4	Salmonella	Eggs and egg products <sup>(i)</sup>	403	France (228), Poland (60), Spain (59), Denmark (26), Sweden (14), Hungary (9), Slovakia (5), Lithuania (2)	1	1,140.8 (303–1,989)	10.0	<b>↓</b> ↓
5	Norovirus (and other calicivirus)	Vegetables and juices and other products thereof <sup>(j)</sup>	263	France (118), Germany (98), Poland (47)	14	227.0 (151–332)	3.0	_
6	Salmonella	Pig meat and products thereof <sup>(k)</sup>	236	France (104), Italy (67), Germany (65)	12	235.3 (69–341)	7.8	_
7	Salmonella	Bakery products <sup>(I)</sup>	178	Poland (110), Czechia (29), Spain (14), Slovakia (14), Malta (4), Romania (4), Italy (3)	8	352.5 (56–621)	3.3	<b>\</b>



			2021 <sup>(a</sup>	)		Evaluation		
Rank <sup>(c)</sup>	Causative agent	Food vehicle	Cases (N)	Reporting MSs (N cases)	Rank <sup>(c)</sup>	Cases (N/year) (range) <sup>(d)</sup>	Reporting MSs (N/year) <sup>(d)</sup>	2021 vs. 2017–2020 <sup>(e)</sup>
8	Staphylococcus aureus toxins	Mixed foods <sup>(m)</sup>	166	Portugal (93), Romania (44), Spain (15), Denmark (11), Italy (3)	25	107.0 (3–212)	4.5	<b>↑</b> ↑
9	Clostridium perfringens toxins	Mixed foods <sup>(n)</sup>	161	Italy (69), France (39), Portugal (20), Germany (15), Finland (12), Denmark (6)	6	380.8 (292–507)	5.3	<b>↓</b> ↓
10	Norovirus (and other calicivirus)	Crustaceans, shellfish, molluscs and products thereof <sup>(o)</sup>	147	France (53), Italy (49), Sweden (37), Spain (8)	4	658.0 (104–1,152)	5.8	††

- (a): Data on FBOs from the United Kingdom (Northern Ireland) are taken into account for 2021. In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/Northern Ireland, the EU requirements on data sampling are also applicable to Northern Ireland.
- (b): Data on FBOs from the United Kingdom are taken into account for 2017–2019, because the United Kingdom was an EU MS, but it became a third country on 1 February 2020.
- (c): Ranking of the food vehicle based on the number of strong-evidence FBOs in which the combination (causative agent/food vehicle) was implicated (rank 1 is the highest rank meaning the most commonly implicated). Strong-evidence outbreaks with unknown causative agents are not included.
- (d): Outbreaks reported by the United Kingdom are also included.
- (e): A single arrow indicates variations of 25% and 50% in the number of outbreaks; double arrows indicate variations > 50%; a 'stable' value indicates variations of between -25% and +25%.
- (f): 'Vegetables and juices and other products thereof' include 'Alfalfa sprouts', 'Melons (except watermelon)', 'Vegetables pre-cut', 'Vegetables and juices and other products thereof'.
- (g): 'Mixed food' includes 'Mixed food', 'Other processed food products and prepared dishes', 'Other processed food products and prepared dishes sushi', 'Ready-to-eat salads'.
- (h): 'Mixed food' includes 'Foodstuffs intended for special nutritional uses dietary foods for special medical purposes', 'Mixed food', 'Other processed food products and prepared dishes fish and seafood-based dishes', 'Other processed food products and prepared dishes pasta', 'Other processed food products and prepared dishes pasta', 'Sauce and dressings mayonnaise'.
- (i): 'Eggs and egg products' include 'Eggs', 'Eggs and egg products'.
- (i): 'Vegetables and juices and other products thereof' include 'Lettuce', 'Vegetables and juices and other products thereof'.
- (k): 'Pig meat and products thereof' includes 'Meat from pig meat products', 'Meat from pig meat products ready-to-eat', 'Meat from pig offal', 'Pig meat and products thereof'.
- (I): 'Bakery products' include 'Bakery products', 'Bakery products cakes', 'Bakery products cakes containing heat-treated cream', 'Bakery products desserts containing raw cream', 'Bakery products desserts containing raw eggs'.
- (m): 'Mixed food' includes 'Mixed food', 'Other processed food products and prepared dishes', 'Other processed food products and prepared dishes meat-based dishes', 'Other processed food products and prepared dishes mushroom-based dishes'.
- (n): 'Mixed food' includes 'Mixed food', 'Other processed food products and prepared dishes,' 'Other processed food products and prepared dishes unspecified', 'Sauce and dressings', 'Soups'.
- (o): 'Crustaceans, shellfish, molluscs and products thereof' includes 'Crustaceans, shellfish, molluscs and products thereof', 'Molluscan shellfish'.



Table 66: Top-10 pathogen/food vehicle pairs causing the highest number of hospitalisations in strong-evidence outbreaks in reporting EU MSs, 2021

		2	021 <sup>(a)</sup>		2017–2020 <sup>(b)</sup>			Evaluation
Rank <sup>(c)</sup>	Causative agent	Food vehicle	Hospitalisations (N)	Reporting MSs (N hospitalisations)	Rank <sup>(c)</sup>	Hospitalisations (N/year) (range) <sup>(d)</sup>		2021 vs 2017–2020 <sup>(e)</sup>
1	Salmonella	Mixed foods <sup>(f)</sup>	82	Denmark (30), Slovakia (25), Spain (10), Germany (4), Finland (4), Poland (3), Italy (3), Belgium (2), Malta (1)	2	100.5 (11–198)	7.8	_
2	Salmonella	Eggs and egg products <sup>(g)</sup>	79	France (24), Spain (22), Poland (18), Denmark (12), Hungary (2), Lithuania (1)	1	251.5 (46–382)	10.0	<b>↓</b> ↓
3	Salmonella	Vegetables and juices and other products thereof <sup>(h)</sup>	75	Germany (30), Denmark (19), Belgium (17), Austria (7), Finland (2)	35	7.0 (2–12)	3.0	$\uparrow \uparrow$
4	Salmonella	Bakery products <sup>(i)</sup>	48	Poland (35), Slovakia (4), Romania (3), Italy (3), Czechia (2), Spain (1)	3	90.5 (21–148)	3.3	<b>\</b>
5	Salmonella	Pig meat and products thereof <sup>(j)</sup>	44	Germany (22), France (18), Italy (4)	6	54.5 (22–94)	7.8	_
6	E. coli other than STEC	Vegetables and juices and other products thereof <sup>(k)</sup>	35	Denmark (24), France (11)	_	_	_	NA
7	Salmonella	Buffet meals	33	Lithuania (23), Austria (10)	4	63.8 (3–168)	2.0	<b>\</b>
8	Norovirus (and other calicivirus)	Mixed foods <sup>(1)</sup>	32	Portugal (24), Italy (7), Belgium (1)	65	2.3 (0–6)	1.0	$\uparrow \uparrow$
8	Campylobacter	Broiler meat (Gallus gallus) and products thereof <sup>(m)</sup>	32	Denmark (29), Spain (3)	61	2.8 (1–8)	1.3	<b>↑</b> ↑
9	Campylobacter	Mixed foods <sup>(n)</sup>	25	Spain (17), Slovakia (4), Finland (2), Poland (2)	67	2.0 (0–3)	1.5	$\uparrow \uparrow$

NA: Not available.

<sup>(</sup>a): Data on FBOs from the United Kingdom (Northern Ireland) are taken into account for 2021. In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/Northern Ireland, the EU requirements on data sampling are also applicable to Northern Ireland.

<sup>(</sup>b): Data on FBOs from the United Kingdom are taken into account for 2017–2019, because the United Kingdom was an EU MS, but it became a third country on 1 February 2020.

<sup>(</sup>c): Ranking of the food vehicle based on the number of strong-evidence FBOs in which the combination (causative agent/food vehicle) was implicated (rank 1 is the highest rank meaning the most commonly implicated). Strong-evidence outbreaks with unknown causative agents are not included.

<sup>(</sup>d): Outbreaks reported by the United Kingdom are also included.

<sup>(</sup>e): A single arrow indicates variations of 25% and 50% in the number of outbreaks; double arrows indicate variations > 50%; a 'stable' value indicates variations of between -25% and +25%.

<sup>(</sup>f): 'Mixed food' includes 'Foodstuffs intended for special nutritional uses – dietary foods for special medical purposes', 'Mixed food', 'Other processed food products and prepared dishes – fish and seafood-based dishes', 'Other processed food products and prepared dishes – pasta', 'Other processed food products and prepared dishes – pasta', 'Sauce and dressings – mayonnaise'.

<sup>(</sup>g): 'Eggs and egg products' include 'Eggs', 'Eggs and egg products'.



- (h): 'Vegetables and juices and other products thereof' include 'Alfalfa sprouts', 'Melons (except watermelon)', 'Vegetables pre-cut', 'Vegetables and juices and other products thereof'.
- (i): 'Bakery products' include 'Bakery products', 'Bakery products cakes', 'Bakery products cakes containing heat-treated cream', 'Bakery products desserts containing raw cream', 'Bakery products desserts containing raw eggs'.
- (j): 'Pig meat and products thereof' includes 'Meat from pig meat products', 'Meat from pig meat products ready-to-eat', 'Meat from pig offal', 'Pig meat and products thereof'.
- (k): 'Vegetables and juices and other products thereof' include 'Spring onion', 'Vegetables pre-cut', 'Vegetables and juices and other products thereof'.
- (I): 'Mixed food' includes 'Mixed food', 'Other processed food products and prepared dishes', 'Other processed food products and prepared dishes sushi', 'Ready-to-eat salads'
- (m): 'Broiler meat (Gallus gallus) and products thereof' include 'Broiler meat (Gallus gallus) and products thereof'. 'Meat from broilers (Gallus gallus)'.
- (n): 'Mixed food' includes 'Foodstuffs intended for special nutritional uses other food for infants and children', 'Mixed food'.

**Table 67:** Top seven pathogen/food vehicle pairs causing the highest number of deaths in strong-evidence outbreaks in reporting EU MSs, 2021

		2021 <sup>(a)</sup>				b)	Evaluation	
Rank <sup>(c)</sup>	Causative agent	Food vehicle	Deaths (N)	Reporting MSs (N deaths)	Rank <sup>(c)</sup>	Deaths (N/year) (range) <sup>(d)</sup>	Reporting MSs (N/year) <sup>(d)</sup>	2021 vs. 2017–2020 <sup>(e)</sup>
1	Listeria monocytogenes	Fish and fish products <sup>(f)</sup>	4	Netherlands (4)	2	2.0 (0–8)	0.5	$\uparrow \uparrow$
2	Clostridium perfringens toxins	Pig meat and products thereof	3	France (3)	22	0.3 (0–1)	0.3	$\uparrow \uparrow$
3	Listeria monocytogenes	Meat and meat products, unspecified	2	Austria (2)	1	2.5 (0–9)	8.0	-
4	Cronobacter sakazakii	Dairy products (other than cheese)	1	Germany (1)	_	_	_	NA
4	Norovirus (and other calicivirus)	Eggs and egg products	1	France (1)	_	-	-	NA
4	Salmonella	Mixed foods	1	Spain (1)	12	0.5 (0–1)	0.5	$\uparrow \uparrow$
4	Bacterial toxins, unspecified	Mixed foods <sup>(g)</sup>	1	France (1)	_	_	_	NA

#### NA: Not available.

- (a): Data on FBOs from the United Kingdom (Northern Ireland) are taken into account for 2021. In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/Northern Ireland, the EU requirements on data sampling are also applicable to Northern Ireland.
- (b): Data on FBOs from the United Kingdom are taken into account for 2017–2019, because the United Kingdom was an EU MS, but it became a third country on 1 February 2020.
- (c): Ranking of the food vehicle based on the number of strong-evidence FBOs in which the combination (causative agent/food vehicle) was implicated (rank 1 is the highest rank meaning the most commonly implicated). Strong-evidence outbreaks with unknown causative agents are not included.
- (d): Outbreaks reported by the United Kingdom are also included.
- (e): A single arrow indicates variations of 25% and 50% in the number of outbreaks; double arrows indicate variations > 50%; a 'stable' value indicates variations of between -25% and +25%.
- (f): 'Fish and fish products' include 'Fish smoked'.
- (g): 'Mixed food' includes 'Other processed food products and prepared dishes'.



# 4.4. Overview of the places of exposure in strong-evidence outbreaks

Table 68 describes the characteristics of strong-evidence FBOs by place of exposure. More information regarding the places of exposures implicated in strong- and weak-evidence FBOs in 2021, with data per causative agent, are available on the dashboard (available here, see page 'Places of exposures').

In 2021, most reported FBOs occurred on 'domestic premises' similarly to what has been observed in recent years. In 2021, 103 FBOs taking place in a 'domestic setting' were described as household FBOs (85.1% of FBOs in domestic settings). However, since not all MSs communicate data on household FBOs, which frequently occur in a domestic setting, this value is likely to be underestimated. 10 MSs (Czechia, Finland, France, Germany, Italy, the Netherlands, Poland, Romania, Slovakia, Spain) reported strong-evidence FBOs on 'domestic premises'. In 2021, the proportion of strong-evidence FBOs occurring in this setting (34.1%) was the lowest since 2012. In 2020 it was 37.1% and over the period 2017–2019 this proportion stood at an average of 39.1%. Bacteria, in particular *Salmonella* were the causative agent most frequently reported in this context in 2021 (68 FBOs; 56.2%).

After domestic settings, '**restaurants**, **pubs**, **street vendors**, **takeaway**, **etc.**' were the most frequently mentioned places of exposure in FBOs, with almost one in four FBOs (23.1%) in the EU occurring in these settings. The proportion was similar in 2020 (25.0%). In the pre-pandemic years (2017–2019), FBOs in 'restaurants, pubs, street vendors, takeaway, etc.' accounted for 28.5% of total strong-evidence FBOs. The main causative agents included *Salmonella* (24 FBOs; 29.3%), bacterial toxins (*Staphylococcus aureus*, *Clostridium perfringens*, *Bacillus cereus*, bacterial toxins unspecified; 17 FBOs; 20.7%), norovirus (and other calicivirus) (14 FBOs; 17.1%) and histamine (10 FBOs; 12.2%).

Most human cases arising from FBOs in 2021 were exposed to contaminated food in 'canteen or catering at workplace, school, etc.'(N = 2,152), especially in 'school or kindergarten' (97.8% of FBOs in 'canteen or catering at workplace, school, etc.'). FBOs in 'school or kindergarten' saw a considerable relative increase, accounting for 11.8% of total strong-evidence FBOs in 2021, 4.8% in 2020 and 5.5% during the period 2017–2019. It is worth remembering that many catering activities in schools and workplaces were suspended in 2020, owing to COVID-19. *Salmonella* was the causative agent mostly reported in FBOs occurring in 'school or kindergarten' (10 FBOs, 23.8%).

A considerable increase was also observed in the proportion of FBOs in **'Health care and residential facilities**', which accounted for 5.1% of strong-evidence FBOs during the period 2017– 2019, 7.7% in 2020 and 9.0% in 2021. Outbreaks in 'Health care and residential facilities' caused the highest number of deaths (N = 7; 53.8% of all deaths), mainly as a result of FBOs in a **'residential institution (nursing home or prison or boarding school)'** (N = 6). *C. perfringens, Salmonella* and Norovirus (and other calicivirus) were the main causative agents.

'Restaurant or cafe or pub or bar or hotel or catering service' was the most frequent place of exposure described for 'general outbreaks' (76 outbreaks, 32.2% of strong-evidence general outbreaks).

FBOs in 'school or kindergarten' were the largest in terms of the mean number of cases per FBO (50.1 cases). The mean size of FBOs in 'restaurant or cafe or pub or bar or hotel or catering service' was much smaller (14.7 mean cases).

# 4.5. Contributing factors in strong-evidence foodborne outbreaks

Information on the factors contributing to food contamination or improper food preparation and storage were available for 126 FBOs reported by MSs in 2021 (35.5% of strong-evidence FBOs). 'Cross-contamination' was reported in 34 FBOs caused by Salmonella, Campylobacter, bacterial toxins, norovirus, Listeria monocytogenes, Aeromonas or non-toxigenic Vibrio cholera. 'Cross-contamination' was the most frequently reported factor in settings such as 'canteen or catering at workplace, school, etc.' and 'health care and residential facilities'. The use of an 'unprocessed contaminated ingredient' was reported in 39 FBOs, mainly involving Salmonella, and was the main contributory factor in domestic settings. An 'infected food handler' is a major source of virus transmission (Teunis et al., 2015) and was identified in 26 FBOs implicating mainly norovirus and Salmonella. This was the most important factor in 'restaurants, pubs, street vendors, takeaway, etc.' 'Time/temperature storage abuse' was reported in 14 FBOs caused by bacterial toxins, histamine, Salmonella and Cronobacter sakazakii. 'Inadequate heat treatment' and 'Inadequate chilling' contributed to 11 and 10 FBOs, respectively, involving bacterial toxins, Salmonella and histamine. 'Untreated drinking water' was linked to one Norovirus FBO.



Table 68: Frequency distribution of strong-evidence foodborne outbreaks by place of exposure, in reporting EU, MSs, 2021

			Stro	ng-evide	nce out	breaks			Outbreak Reporting Rate per 100,000		
Type of setting	Out	breaks	Ca	ses	Hosp	italisations		Deaths			
	N	% of total	N	% of total	N	% of total	N	% of total	2021 <sup>(a)</sup>	2020	2017–2019 (mean) <sup>(b)</sup>
Domestic setting	121	34.1	803	11.5	161	21.7	0	0	0.027	0.022	0.054
Restaurant, pub, street vendors, take away,	etc.										
Restaurant or Cafe or Pub or Bar or Hotel or Catering service	77	21.7	1,134	16.2	72	9.7	0	0	0.017	0.012	0.037
Take-away or fast-food outlet	4	1.1	26	0.37	0	0	0	0	0.001	0.002	0.001
Mobile retailer or market/street vendor	1	0.28	13	0.19	0	0	0	0	< 0.01	< 0.01	0.001
Canteen or catering at workplace, school, etc											
School or kindergarten	42	11.8	2,104	30.0	146	19.7	0	0	0.009	0.003	0.007
Canteen or workplace catering	3	0.85	48	0.69	0	0	0	0	0.001	0.002	0.005
Health care and residential facilities											
Residential institution (nursing home or prison or boarding school)	21	5.9	449	6.4	20	2.7	6	46.2	0.005	0.004	0.005
Hospital and medical care facility	11	3.1	228	3.3	14	1.9	1	7.7	0.002	< 0.01	0.002
Multiple places of exposure											
Multiple places of exposure in one country	20	5.6	715	10.2	111	15.0	6	46.2	0.004	0.003	0.005
Multiple places of exposure in more than one country	6	1.7	306	4.4	90	12.1	0	0	0.001	0.001	0.001
Other place of exposure											
Others	14	3.9	361	5.2	29	3.9	0	0	0.003	0.004	0.008
Camp or picnic	8	2.3	220	3.1	39	5.3	0	0	0.002	< 0.01	0.002
Farm	1	0.28	39	0.56	0	0	0	0	< 0.01	< 0.01	0.001
Temporary mass catering (fairs or festivals)	1	0.28	19	0.27	1	0.13	0	0	< 0.01	< 0.01	< 0.01
Primary production	1	0.28	3	0.04	0	0	0	0	< 0.01	_	_
Unknown	24	6.8	537	7.7	58	7.8	0	0	0.005	0.003	0.008
EU Total <sup>(b)</sup>	355	100	7,005	100	741	100	13	100	0.079	0.055	0.137

<sup>(</sup>a): Data on FBOs from the United Kingdom (Northern Ireland) are taken into account for 2021. In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/Northern Ireland, the EU requirements on data sampling are also applicable to Northern Ireland.

<sup>(</sup>b): Data on FBOs from the United Kingdom are taken into account for 2017–2019, because the United Kingdom was an EU MS, but it became a third country on 1 February 2020.



18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Islituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (three/onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Islituto Zooprofilation Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (three/onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Islituto Zooprofilation Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (three/onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Islituto Zooprofilation Sperimenta dell'Umbria e delle Marche (IZSUM). Wiley Online Library on [18/12/2023]. See the Terms and Conditions (three/onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Islituto Zooprofilation Sperimenta dell'Umbria e dell'Umbri


and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

# 4.6. Temporal trends by causative agent

# 4.6.1. Temporal trend at EU level

Figure 33 describes the number of FBOs reported in the EU over the period 2012–2021, by causative agent. Given that the collection of FBO data is not fully harmonised across the EU, and that changes in the surveillance of FBOs by MSs may have taken place over time, annual variations in the frequency distribution of the causative agents shown in Figure 33 may not necessarily reflect the true epidemiological pattern at EU level.

For a further interactive look into data, see the dashboard on FBOs (available here), where there is a dedicate page on time trends.



Note: Outbreaks reported by the United Kingdom are included for the years 2012–2019. However, data from United Kingdom (Northern Ireland) are taken into account for 2021. In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/Northern Ireland, the EU requirements on data sampling are also applicable to Northern Ireland.

2019

2020

2021

'Marine biotoxins' include ciguatoxin, muscle-paralysing toxin, okadaic acid.

Year

2015

2016

2017

2018

'Norovirus (and other calicivirus)' include norovirus (Norwalk-like virus), sapovirus (Sapporo-like virus) and calicivirus unspecified.

'Other bacterial agents' include Aeromonas, Arcobacter, Cronobacter sakazakii, E. coli other than STEC, Enterococcus, Francisella, Leptospira, Shigella, Streptococcus, non-toxigenic Vibrio cholerae, Vibrio parahaemolyticus and other unspecified bacterial agents.

Figure 33: Number of foodborne outbreaks by causative agent, reported to the EU by MSs, 2012–2021

Trichinella Brucella

0

2012

2013

2014



# 4.6.2. Country-specific temporal trends by causative agent over the period 2012–2021

Figures on statistically significant country-specific temporal trends by causative agent, for the period 2012–2021, are available in 'Addendum foodborne outbreaks, 2021' Section 6, published on the EFSA Knowledge Junction on Zenodo here.

A trend analysis of FBOs caused by **Salmonella** showed that three MSs (Germany, Latvia, Lithuania) reported a statistically significant decreasing number of FBOs over the period 2012–2021. For Germany and Lithuania, this trend was mainly driven by a significant decrease of **S. Enteritidis** FBOs over the same period. FBOs caused by **S. Enteritidis** also followed a statistically significant decreasing trend in three other MSs (Austria, Belgium and Hungary). A progressive fall in the number of FBOs caused by this serovar was also observed for Croatia, Czechia and Spain, even though it was not statistically significant. Findings from a study carried out in Europe as part of the DiSCoVeR project emphasised that outbreaks caused by **S.** Enteritidis significantly decreased in Northern and Southern Europe between 2015 and 2019, and significantly increased in Eastern Europe (Chanamé Pinedo et al., 2022). Austria reported a significant negative trend for **S. Typhimurium** and **S. Typhimurium monophasic.** For Germany and Slovakia a similar trend was also observed, although it was not statistically significant. These trends could point to the successful implementation of National Control Programmes (NCP) for *Salmonella* in poultry. Information on the results achieved with the NCPs in poultry by EU MSs is available in Section 2.4.4. (*Salmonella* in animals) of the *Salmonella* chapter.

In Austria, FBOs caused by *Campylobacter* fell significantly, while in France, figures followed an increasing trend over the same period, although the number of *Campylobacter* FBOs reported last year was lower than in 2020.

In France, FBOs caused by *Clostridium perfringens* toxins followed a statistically decreasing trend, even though annual reporting of FBOs caused by this agent has been relatively variable over the years.

In Germany, **norovirus (and other calicivirus)** saw a considerable decrease over the period 2012–2021.

Except in the case of *Salmonella*, hypotheses explaining these trends cannot be easily formulated based on the information reported to EFSA. Most of the FBOs were actually weak-evidence outbreaks characterised by a high degree of uncertainty concerning the implicated food vehicles and the environment in which the outbreak occurred.

In the Netherlands and Switzerland, the number of FBOs of **unknown** origin increased significantly during the study period, while Norway reported a substantial decrease. In 2021, the FBOs reported by the Netherlands (N=810) accounted for 20.2% of all FBOs reported in the EU, totalling more than 3.5 times the number of FBOs notified to EFSA in 2012 (N=216). The reasons underlying increasing or decreasing trends in FBOs caused by unknown agents may reflect changes in the sensitivity of outbreak surveillance, as well as changes in laboratory sample testing and subtyping. As a result, the diversion of both human and technical resources in recent years, to fight the COVID-19 pandemic, should be considered as a possible reason for the increase.

Information on the number of strong-evidence outbreaks, by causative agent, over the years 2016–2021 can be viewed on the dashboard (available here).

# 4.6.3. Temporal trends by implicated food vehicle

Figures on statistically significant country-specific temporal trends by food vehicle, for the period 2012–2020, are available in the 'Addendum foodborne outbreaks, 2021' Section 7, published on the EFSA Knowledge Junction on Zenodo here.

Few statistically significant increasing and decreasing trends were detected for food vehicles over the 2012–2021 period. Germany reported an increasing trend in FBOs associated with the consumption of 'cereal products including rice and seeds/pulses (nuts, almonds)'. The number of FBOs involving this foodstuff in annual reports was very low (< 10 FBOs/year), making it difficult to investigate possible reasons underlying this finding.

In France, FBOs caused by the consumption of 'fish and fish products' saw a statistically significant decrease over the period 2012–2021. This fall was mainly linked to a reduction in the number of FBOs caused by 'histamine and scombrotoxin' and 'marine biotoxins'.

Information on the number of strong-evidence outbreaks, by food vehicle, over the years 2016–2021 can be viewed on the dashboard (available here).



## 4.7. Waterborne outbreaks

In 2021, seven MSs (Austria, Finland, Greece, Ireland, Italy, Spain and Sweden) reported a total of 12 waterborne outbreaks caused by the intake of 'tap water, including well water' (11 outbreaks) and 'potable water' (one outbreak). Moreover, three outbreaks were notified by two non-MSs (Switzerland and Republic of North Macedonia) caused by 'tap water, including well water' (two outbreaks) and 'water, unspecified'.

At EU level, the number of waterborne outbreaks and cases reported in 2021 decreased by 55.6% and 28.1% compared with 2020 (27 outbreaks and 278 human cases in 2020), respectively. Compared with the pre-pandemic years, the number of outbreaks and human cases reported in 2021 were considerably lower than during the 2017–2019 period, with a relative decrease of 69.5% (39 mean outbreaks reported annually during 2017–2019) and 85.9% (1,422 cases on average, during 2017–2019) respectively. In 2021, the mean outbreak size in EU MSs was 16.7.

It is worth mentioning a large general outbreak reported in 2021 by the Republic of North Macedonia, which involved 93 human cases. No information was available on the causative agent and the outbreak was associated with 'water, unspecified'.

Four strong-evidence outbreaks were reported in 2021 by two MSs (Finland and Spain, one FBO each) and by Switzerland (N = 2), involving 146 human cases. All outbreaks were associated with the intake of 'tap water including well water' and were caused by 'norovirus' in MSs and by 'E. coli other than STEC' in Switzerland. Remarkably, the outbreak reported by Spain was described as a general outbreak and involved a total of 114 human cases.

Eleven weak-evidence outbreaks were reported by six MSs (Austria, Finland, Greece, Ireland, Italy and Sweden) and one non-MS (Republic of North Macedonia). In the EU, 'tap water, including well water' was the main source of waterborne outbreaks (N=9), followed by 'potable water' (one outbreak). The weak-evidence outbreak reported by the Republic of North Macedonia is described above. All together weak-evidence outbreaks reported in 2021 were caused by STEC (N=4) and Campylobacter (N=2). For five weak-evidence waterborne outbreaks, the agent was 'unknown'.

# 4.8. ECDC and EFSA rapid outbreak assessment (ROA) of multi-country foodborne outbreaks

The ECDC and EFSA jointly assessed the public health risk posed by various FBOs occurring in 2021, publishing their Joint Rapid Outbreak Assessments (ROA). The first ROA concerned an FBO attributable to *Salmonella* Braenderup sequence type 22, which was responsible for 348 cases, and 68 hospitalisations, in 12 European countries and the United Kingdom. The cases occurred between March and July 2021. Evidence from epidemiological, microbiological and trace-back investigations identified imported melons as the suspected source of the outbreak. In particular, following case interviews and an analytical epidemiological study, Galia melons were identified as the probable vehicle of infection. Although *S.* Braenderup matching the outbreak strain was isolated in melons, including Galia melons imported from Honduras, at the time of the publication of the ROA it was not possible to establish the exact point of contamination in the food production chain. Withdrawals and recalls of contaminated products were implemented as control measures (ECDC and EFSA, 2021a).

Another ROA concerned an FBO caused by multiple serovars of *Salmonella enterica*, namely *S.* Amsterdam, *S.* Havana, *S.* Kintambo, *S.* Mbandaka, *S.* Orion and *S.* Senftenberg, which has been ongoing since 2019. It involved 121 confirmed cases, and 22 hospitalisations from five EU/EEA countries (Denmark, Germany, the Netherlands, Norway and Sweden), between 2019 and 2021. Noticeably, almost half of the cases and most hospitalisations (12/22) involved children under 10 years of age. Epidemiological, microbiological and food tracing investigations identified sesame-based products imported from Syria as the probable vehicles of infection. Although control measures (withdrawals, recalls and destructions) have been implemented since 2020, these were not sufficient to prevent the occurrence of human cases in 2021. Moreover, sesame-based products have a long shelf life and could be stored in people's homes over a long period (ECDC and EFSA, 2021b), thereby highlighting the risk of a prolonged pattern of human exposure to *Salmonella* caused by these foodstuffs.

Another ROA concerned an FBO caused by *Salmonella* Enteritidis ST11 in 2021. In total, 272 confirmed human cases, 25 hospitalisations and two deaths were notified by five European countries and the United Kingdom. France reported the highest number of cases (N = 216), followed by Spain (N = 22), the Netherlands (N = 12), the United Kingdom (N = 12), Norway (N = 7) and Denmark (N = 3). Following epidemiological, microbiological and traceability investigations, eggs from farms in



Spain were identified as the potential source of the outbreak. Table eggs from the farms implicated in the outbreak were withdrawn and redirected for use in heat-treated egg products. Interestingly, this outbreak was microbiologically linked to a historical outbreak reported by the Netherlands in 2019. However, no epidemiological link between the two events was identified (ECDC and EFSA, 2022b).

## 5. Conclusions

## 5.1. Health impact, causative agents and trends

In 2021, an increase was observed in the reporting of FBOs by most MSs and non-MSs, compared with 2020. Across the EU and the United Kingdom (Northern Ireland) a total of 4,005 FBOs were reported, an increase of approximately one-third (29.8%) compared with 2020. However, this number was still well below the mean number of FBOs reported to EFSA in the pre-pandemic years (5,601 FBOs/year reported on average during the period 2017–2019). A similar pattern could be observed for other indicators relating to the impact of FBOs on public health (primarily cases and hospitalisations). To properly interpret these data, since FBO surveillance is not fully harmonised among MSs, it is important to remember that statistics at EU level reflect mostly the pattern of FBO occurrence of those MSs that contributed the most to data collection in 2021. The findings described above suggest that the consequences of the COVID-19 pandemic were still visible in 2021, in both the occurrence of FBOs in the EU and the activities of detection, investigation and reporting. This is also supported by the fact that the drop in FBO reporting observed in a few countries in 2020 following the arrival of the COVID-19 pandemic, was not reversed in 2021.

The year 2021 was the second year of the COVID-19 pandemic, with several new waves being reported across Europe. Also in this second year, a range of control measures were adopted by National Authorities in the EU to limit the spread of the virus, with content varying from one country to another. These measures included actions to reduce contact (e.g. stay-at-home orders, banning of private gatherings, etc.), promote hygiene and safety measures (e.g. the use of protective equipment, disinfection procedures, etc.) and other preventive measures.<sup>35</sup> The control measures implemented in 2021 were less stringent than in 2020, but could still explain the downturn in FBOs for the second year of the pandemic. These control measures could have had an impact in decreasing the foodborne exposure of the population to zoonotic agents (e.g. better hygiene, closure of restaurants, etc.), improving hygiene measures by consumers (e.g. gloves, hand sanitiser, cleaning of surfaces and equipment, etc.) or even in reducing travel-related FBOs, as observed in the literature (Ray et al., 2021; van Deursen et al., 2022). The diversion of technical, financial and human resources towards activities relating to the COVID-19 pandemic may also have had an important role. In many countries, the COVID-19 pandemic placed healthcare services under severe strain and negatively impacted doctor-patient interaction, leading to healthcare difficulties, with problems of under-diagnosis and under-reporting of non-COVID diseases (Kastritis et al., 2020; Verhoeven et al., 2020; Lim et al., 2021; WHO, 2022). Nonetheless, the increase in the number of FBOs compared with 2020 might indicate a gradual return to pre-pandemic stability in FBO surveillance for most MSs.

Overall, the causative agents associated with FBOs in 2021 were similar to those observed in previous years. In the EU, *Salmonella* remained the most frequently reported pathogen, for the number of FBOs (i.e. one in five), as well as for human cases and hospitalisations. *S.* Enteritidis and *S.* Typhimurium were the predominant serovars. At national level, *Salmonella* was the main cause of FBOs in most EU MSs (17 MSs) and the United Kingdom (Northern Ireland), as well as in six non-MSs. However, in a few countries, the number of FBOs caused by *Salmonella* was surpassed by other agents including either *Campylobacter*, STEC, *Staphylococcus aureus* toxins or norovirus. This finding highlights considerable differences in the epidemiology of FBOs among MSs, which is probably attributable to different risks of exposure to pathogens through food consumption, and differences in the structure of FBO surveillance.

One critical finding emerging from the 2021 data analysis concerns the number of listeriosis FBOs reported to EFSA, which was higher than in both 2020 and the pre-pandemic years. However, these outbreaks were apparently less severe compared with recent years, as fewer cases, hospitalisations and deaths were reported in 2021. Possible reasons for this increase are discussed in the section dedicated to *Listeria* (4.2.1 Bacteria). The rise in the reporting of FBOs caused by *Listeria monocytogenes* is a matter of concern since high rates of hospitalisation and death are associated with

<sup>&</sup>lt;sup>35</sup> More information is available at: https://covid-statistics.jrc.ec.europa.eu/RMeasures



this pathogen (Buchanan et al., 2017), especially in vulnerable and fragile populations, whose numbers are increasing in the EU. It is possible that FBO cases involving severe conditions, such as sepsis and meningitis associated with invasive listeriosis, are less likely to be under-diagnosed and under-reported.

In 2021, two strong-evidence FBOs caused by *Cronobacter sakazakii* and *Vibrio cholerae* (nontoxigenic) were reported for the first time to EFSA since the start of data collection. Both outbreaks occurred in 'Health care and residential facilities'. One new-born child infected by *C. sakazakii* died after the intake of a contaminated probiotic formula. These events draw attention to the high susceptibility of vulnerable population groups to FBOs.

# 5.2. Food vehicles and places of exposure

In 2021, the range of foodstuffs implicated in FBOs closely reflected the established epidemiology of the implicated causative agents. Overall, considering only single food items, 'mixed food' was the foodstuff implicated in most outbreaks, leading to the highest number of cases and hospitalisations. 'Mixed foods' are complex foods by definition since they include a variety of ingredients, making it difficult to identify the primary source of contamination. Many factors may contribute to the contamination of these foodstuffs, including unsafe food mixing, processing and handling of ingredients by infected food handlers, cross-contamination or even the development of favourable growing conditions for pathogens during food preparation. 'Eggs and egg products' were the second most frequently reported foodstuff in FBOs, followed by 'vegetables and juices and products thereof' and 'fish and fish products'. In 2021, FBOs caused by *Salmonella* coupled with a variety of food vehicles (i.e. 'eggs and egg products', 'mixed foods', 'vegetables and juices and other products thereof' and 'bakery products') led to some of the most severe health impacts in terms of total outbreaks, cases and hospitalisations in the EU.

Interestingly, the implication of 'vegetables and juices and other products thereof' in FBOs increased considerably compared with both 2020 and the pre-pandemic years. In addition, these foods caused the second highest number of human cases. For extensively handled vegetables and fruit, such as pre-cut or frozen items, the food chain leading to the shop or place of consumption may be complex and long, as indicated by the *Salmonella* Braenderup found in melons. Contamination may take place at various levels of the food chain, from the primary production level with the growers, up to the processing and retail chain (Olaimat and Holley, 2012; Alegbeleye et al., 2018). This is the reason why FBOs implicating this foodstuff may have a critical impact on public health and require strong public health and food safety preparedness, with intersectoral coordination for effective control.

As observed in the past, domestic settings remain the most frequently reported place of exposure to foodborne agents. FBOs occurring in this setting were mostly described as 'household' outbreaks. However, it is highly probable that these outbreaks have been under-reported, since household FBOs could have remained undetected or been poorly investigated, especially in the pandemic years. In 2021, these outbreaks were reported only by seven MSs across the EU. This finding highlights the importance of communicating with the public in order to improve awareness of the risk posed by handling food in domestic kitchens, and the need to adopt the correct practices for food preparation and storage.

One remarkable aspect highlighted in 2021 is the increase of outbreaks in 'canteen or catering at workplace, school, etc.', mainly due to the contribution of FBOs occurring in 'school or kindergarten'. Outbreaks occurring in 'school or kindergarten' are relatively easy to detect because they are often point-source outbreaks and/or involve a large number of people. In 2021, outbreaks in 'school or kindergarten' led to the highest number of cases and had a high mean outbreak size (50 cases per FBO). The highest number of deaths was observed in outbreaks occurring in 'health care and residential facilities', drawing attention to the high risks of foodborne outbreaks for vulnerable population groups. The causative agents frequently identified in these places of exposure were mostly unspecified bacterial toxins, *Salmonella* and norovirus. Interestingly, an infected food handler was the main factor contributing to FBOs. This finding underlines the importance of strengthening hygiene standards and HACCP protocols in the food production chain, including training food handlers in food safety principles and procedures.

In conclusion, it is important to remember that trends in the occurrence of FBOs over the years are highly influenced not only by the level of food contamination at consumer level, but also by changes in food consumption habits. Globalisation has revolutionised the food supply system, resulting in an extensive and complex food production chain that may also contribute to an increasing number of



1831/4732, 2022, 12, Downloaded from https://ds.aonlinelibrary.wiley.com/doi/10.2903j.ds.a.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms

foodborne outbreaks (Adinolfi et al., 2016). In addition, the EU is in the throes of demographic change. The ageing of the population will probably affect all EU countries, resulting in the considerably increased susceptibility of vulnerable population groups (Eurostat, 2020).



//efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico

Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common


# Zoonoses monitored in accordance with the epidemiological situation (Directive 2003/99 List B)

#### 1. Yersinia

Summary data substantiating this chapter, as well as additional information on related projects and internet sources are published for this report on the EFSA Knowledge Junction on Zenodo here. Summary statistics on yersiniosis foodborne outbreaks reported in the framework of Directive 2003/99/EC, with downloadable files, are retrievable using the EFSA foodborne outbreaks dashboard available here. Summary statistics on human surveillance data with downloadable files are retrievable using the ECDC Surveillance Atlas of Infectious Diseases here.

# Yersinia in the EU, 2021







# 1.1. Key facts

- Yersiniosis is the third most commonly reported zoonosis in humans in the EU.
- In 2021, the number of confirmed cases of human yersiniosis was 6,789, corresponding to an EU notification rate of 1.9 per 100,000 population. This was an increase of 11.8% compared with the rate in 2020 (1.7 per 100,000 population).
- In 2020, yersiniosis reporting recorded the lowest number of human cases since 2007, when *Yersinia* surveillance began. This low number was related to the impact of the COVID-19 pandemic and the withdrawal of the United Kingdom from the EU.
- Compared with the rate before the COVID-19 pandemic (2017–2019 annual mean), the 2021 EU notification rate increased by 11.3% and decreased 8.9% with and without data from the United Kingdom, respectively.
- The overall yersiniosis trend in 2017–2021 showed no statistically significant increase or decrease.
- In 2021, six MSs reported information on 355 'ready-to-eat' food sampling units tested for the presence of *Yersinia*. There were three positive units, and all were from the 'ready-to-eat' meat and meat products category, in particular 'meat and meat products from pigs' (2.1% positive samples).
- In 'non-ready-to-eat' food, six MSs provided results on 736 sampling units. There were 38 positive units and all originated from 'meat and meat products' with 5.4% positives in this category. In 'fresh meat', five MSs provided information from 418 sampling units, reporting 29 positive units (6.9%). In particular, *Yersinia* was found in 26 (7.1%) out of 366 samples tested in the 'fresh meat of pigs' food category.
- In animals, seven MSs and one non-MS reported results of sampling activities in 2021; the highest overall proportion of *Yersinia*-positive sampling units in farmed animals was observed in 'small ruminants' (N = 96, 2.7%).

# 1.2. Surveillance and monitoring of Yersinia in the EU

#### 1.2.1. Humans

In 2021, 26 MSs reported information on yersiniosis in humans. Surveillance of yersiniosis is mandatory in 22 MSs. In four MSs (Belgium, France, Greece and Italy) notification is based on a voluntary system. No yersiniosis surveillance system is in place in the Netherlands. The surveillance systems for yersiniosis have national coverage in all reporting countries, except for three: France, Italy and Spain. For these three countries, no information on population coverage was provided, and notification rates were not calculated. In 2020–2021, Spain did not received data from all regions that normally report, so the case number might not be complete. The EU case definition was used by 21 MSs, four MSs used a different case definition for reporting (Denmark, France, Germany and Italy), and Greece did not use any case definition. All countries reporting data on yersiniosis in 2021 provided case-based data, except Belgium, Bulgaria and Greece, which reported aggregated data.

#### 1.2.2. Food and animals

Although it is not mandatory to report the presence of *Yersinia* in food and animals, MSs can report monitoring data on *Yersinia* to EFSA, in accordance with the Zoonoses Directive (Directive 2003/99/EC). The directive specifies that, in addition to the zoonoses and zoonotic agents for which monitoring is mandatory, zoonoses such as yersiniosis and agents thereof must also be monitored if warranted by the epidemiological situation in an MS (Annex I, B. 2., in compliance with Article 4.1 of the same Directive). At present, no harmonised *Yersinia* monitoring plan is in place for food or animals in the EU. Therefore, data on *Yersinia* in food and animals submitted to EFSA by the MSs are not harmonised. Data allow for descriptive summary statistics at the EU level only, and do not support trend analyses and trend watching (Table 1). Harmonised monitoring and reporting criteria for *Y. enterocolitica* in slaughter pigs were recommended by EFSA in a scientific report (EFSA, 2009). The reported occurrence of *Yersinia* in major food categories for the year 2021 and for the four-year period 2017–2020 was summarised descriptively, making a distinction between 'ready-to-eat' (RTE) and non-RTE foods.

For the purpose of the 2021 data analysis, only results obtained from samples collected and tested for *Yersinia* under an 'objective sampling' strategy were considered, in order to limit selection bias.



Objective sampling means that MSs collected and tested the samples according to a planned strategy based on a random sampling design representative of the population under study.

#### 1.3. Results

## 1.3.1. Overview of key statistics, EU, 2017–2021

Tables 69 and 70 summarise EU-level statistics on human yersiniosis, and on the occurrence and prevalence of *Yersinia* in food and animals, for the period 2017–2021. Although yersiniosis was the third most frequently reported zoonosis in the EU in 2021, data on *Yersinia* in food and animals continue to be reported by few MSs, like in previous years. More detailed descriptions of these statistics are provided in the below subsections and in the chapter on foodborne outbreaks.

**Table 69:** Summary of *Yersinia* statistics related to humans, major food categories and main animal species, EU, 2017–2021

	2021 <sup>(a)</sup>	2020	2019 <sup>(b)</sup>	2018 <sup>(b)</sup>	2017 <sup>(b)</sup>	Data source
Humans						
Total number of confirmed cases	6,789	5,661	6,967	7,015	6,844	ECDC
Total number of confirmed cases/100,000 population (notification rate)	1.9	1.7	1.7	1.7	1.7	ECDC
Number of reporting MSs	26	26	27	27	27	ECDC
Infection acquired in the EU	3,478	2,686	3,468	3,446	3,410	ECDC
Infection acquired outside the EU	18	61	96	106	88	ECDC
Unknown travel status or unknown country of infection	3,293	2,914	3,403	3,463	3,346	ECDC
Number of foodborne outbreak-related cases	125	236	160	58	130	EFSA
Total number of foodborne outbreaks	21	16	23	12	11	EFSA
Food						
Meat and meat products						
Number of sampling units	1,789	1,597	2,304	1,470	1,211	EFSA
Number of reporting MSs	6	6	6	6	7	EFSA
Fruits and vegetable products						
Number of sampling units	142	256	17	7	116	EFSA
Number of reporting MSs	3	4	2	2	4	EFSA
Animals <sup>(c)</sup>						
Bovine animals						
Number of sampling units	19,218	14,796	16,885	13,101	17,404	EFSA
Number of reporting MSs	5	5	5	6	8	EFSA
Pigs						
Number of sampling units	2,164	2,368	2,591	2,347	2,781	EFSA
Number of reporting MSs	5	4	6	7	7	EFSA

ECDC: European Centre for Disease Prevention and Control; EFSA: European Food Safety Authority; MSs: Member States.

204

18314732, 2022, 12, Downloaded from https://efs.aonlinelibrary.wiley.com/doi/10.2903/jefs.a.2022.7666 by Istituto Zopprofilative Sperimenta dell'Umbria e delle Marche (ISSUM), Wiley Online library.wiley.com/rems-and-conditions) on Wiley Online library for rules of use. O. A articles are governed by the applicable Creative Common

<sup>(</sup>a): Data on food and animals from the United Kingdom (Northern Ireland) are taken into account for 2021. In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/ Northern Ireland, EU requirements on data sampling are also applicable to Northern Ireland.

<sup>(</sup>b): Data from the United Kingdom are taken into account for 2017–2019, because the United Kingdom was an EU MS, but it became a third country on 1 February 2020.

<sup>(</sup>c): Samples as single animals.



#### 1.3.2. Human yersiniosis

In 2021, 6,789 confirmed cases of yersiniosis were reported by 26 MSs. As in recent years, Germany accounted for the highest number of cases, followed by France (Table 70). Cases reported by these countries together accounted for 49.5% of all confirmed yersiniosis cases in the EU. In particular, the number of cases increased in France and Denmark from 2017 to 2021. The highest notification rates (per 100,000 population) were observed for Denmark (7.8), Finland (6.0), Lithuania (5.5) and Latvia (4.4).

The notification rate of confirmed yersiniosis cases in the EU was 1.9 cases per 100,000 population. This corresponds to an increase of 11.8% compared with the rate in 2020 (1.7 per 100,000 population), and an increase of 11.3% and decrease of 8.9% compared with the average annual notification rate from 2017 to 2019 (pre-pandemic period) with and without United Kingdom data, respectively.

The majority of yersiniosis cases, i.e. 3,478 (99.5%) reported with known origin of infection were from the EU (Table 70). Germany, Sweden, Denmark, Austria and Finland reported the highest number of travel-associated cases (33, 16, 15, 6 and 6 cases, respectively) out of a total of 84 cases. Overall, the proportion of travel-associated cases of yersiniosis was only 2.4%, including 60 travel-associated cases in the EU (1.7%), and 18 cases (0.5%) outside the EU reported by 18 MSs and 6 with origin unknown (0.2%).

**Table 70:** Reported confirmed human cases of yersiniosis and notification rates per 100,000 population in EU MSs and non-MS countries, by country and year, 2017–2021

		2021			20	2020		2019		18	201	L7
Country	National coverage <sup>(a)</sup>	Data format <sup>(a)</sup>	Confii cases rat	and	Confine cases	and	Confii cases rat	and	Confirmed cases and rates		Confirmed cases and rates	
			Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate
Austria	Y	С	138	1.5	128	1.4	112	1.3	136	1.5	95	1.1
Belgium	Υ	Α	418	3.6	260	2.3	406	3.5	392	3.4	317	2.8
Bulgaria	Υ	Α	5	0.07	4	0.06	11	0.16	9	0.13	17	0.24
Croatia	Υ	С	14	0.35	11	0.27	12	0.29	20	0.49	29	0.7
Cyprus	Υ	С	0	0	0	0	0	0	0	0	0	0
Czechia	Υ	С	456	4.3	437	4.1	618	5.8	622	5.9	611	5.8
Denmark	Υ	С	453	7.8	413	7.1	221	3.8	282	4.9	206	3.6
Estonia	Υ	С	45	3.4	44	3.3	42	3.2	63	4.8	43	3.3
Finland	Υ	С	331	6.0	386	7.0	406	7.4	529	9.6	423	7.7
France <sup>(b)</sup>	N	С	1,451	_	988	_	1,135	_	929	_	738	_
Germany	Υ	С	1,912	2.3	1,860	2.2	2,164	2.6	2,193	2.6	2,581	3.1
Greece	Υ	Α	7	0.07	3	0.0	13	0.12	21	0.20	19	0.18
Hungary	Y	С	50	0.51	25	0.26	38	0.39	36	0.37	30	0.31
Ireland	Υ	С	19	0.38	13	0.26	9	0.18	8	0.17	6	0.13
Italy <sup>(b)</sup>	N	С	35	_	21	_	12	_	14	_	8	_
Latvia	Υ	С	83	4.4	88	4.6	60	3.1	68	3.5	47	2.4
Lithuania	Y	С	153	5.5	123	4.4	181	6.5	139	4.9	174	6.1
Luxembourg	Υ	С	12	1.9	26	4.2	18	2.9	16	2.7	15	2.5
Malta	Y	С	0	0	0	0	0	0	0	0	0	0
Netherlands <sup>(c)</sup>	_	_	_	_	_	_	_	_	_	_	_	_
Poland	Υ	С	142	0.38	90	0.24	196	0.52	170	0.45	191	0.50
Portugal	Υ	С	34	0.33	25	0.24	29	0.28	30	0.29	35	0.34
Romania	Υ	С	15	0.08	6	0.03	36	0.19	22	0.11	36	0.2
Slovakia	Υ	С	213	3.9	168	3.1	255	4.7	259	4.8	242	4.5
Slovenia	Υ	С	49	2.3	26	1.2	28	1.3	32	1.5	18	0.87
Spain <sup>(b),(d)</sup>	N	С	444	_	296	_	409	_	549	_	585	_
Sweden	Υ	С	310	3.0	220	2.1	393	3.8	278	2.7	236	2.4



18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://doi.org/10.1001/10.100


and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons

		2021			20	2020		2019		18	201	.7
Country	National coverage <sup>(a)</sup>	Data format <sup>(a)</sup>	Confine cases	and	Confine cases	and	Confirmed cases and rates		Confirmed cases and rates		Confirmed cases and rates	
			Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate
EU Total 27	_	_	6,789	1.9	5,661	1.7	6,804	2.1	6,817	2.1	6,702	2.1
United Kingdom	_	_	_	_	_	_	163	0.24	198	0.30	142	0.22
EU Total <sup>(e)</sup>	_	_	6,789	1.9	5,661	1.7	6,967	1.7	7,015	1.7	6,844	1.7
Iceland	Υ	С	4	1.1	3	0.82	2	0.56	2	0.57	0	0
Norway	Υ	С	85	1.6	83	1.5	85	1.6	105	2.0	67	1.3
Liechtenstein	_	_	-	_	_	_	_	_	-	_	_	_
Switzerland	_	_	_	_								

<sup>-:</sup> Data not reported.

- (a): Y: yes; N: no; A: aggregated data; C: case-based data.
- (b): Sentinel surveillance; no information on estimated coverage. Notification rate not estimated.
- (c): No surveillance system.
- (d): Data not complete in 2020–2021, rate not estimated.
- (e): Cases reported by the United Kingdom for the period 2017–2019 were also considered for this estimation (EU-28). When the United Kingdom data were collected for the period 2017–2019, the United Kingdom was an EU MS, but it became a third country on 1 February 2020.

The trend of human yersiniosis cases in the EU for 2017–2021 showed no statistically significant increase or decrease (Figure 34). At the MS level, a statistically significant decrease (p < 0.05) was observed for Germany and Romania. For the same time period, there was a statistically significant increase (p < 0.05) for four MSs (Denmark, France, Italy and Latvia).



Source: Austria, Czechia, Denmark, Estonia, Finland, Germany, Hungary, Ireland, Italy, Latvia, Luxembourg, Malta, Poland, Romania, Slovakia, Slovenia and Sweden.

Figure 34: Trends in reported confirmed human cases of yersiniosis in the EU, by month, 2012–2021

Of the 1,564 (23.0%) yersiniosis cases with available hospitalisation information reported by 13 MSs, 508 cases (32.5%) were hospitalised. No deaths were reported among the 3,596 confirmed cases with available clinical outcome information. The outcome was reported for 53.0% of all confirmed cases of yersiniosis by 21 MSs.

Information on the *Yersinia* species detected in confirmed cases was reported by 21 MSs for 6,063 cases (95.3%) in 2021. *Y. enterocolitica* was the most commonly reported species in all countries, among 5,950 human cases (98.1% of all cases with information on species available). Serotype information on *Y. enterocolitica* was provided for 2,984 (50.2%) confirmed cases. The most commonly reported serotype was O3 (83.1% of cases with information on serotype available), followed by O9 (12.3%). Overall, the other serotypes (O1, O5, O5, 27, O8 and others) accounted for 4.6% of cases with known serotype. Biotype information was provided for 1,727 (29.0%) confirmed cases. The most commonly reported biotype of *Y. enterocolitica* was biotype 4 (82.3% of all cases with this information available), followed by biotype 2 (16.7%), biotype 1B (0.6%) and biotype 3 (0.4%). Information on *Y. enterocolitica* bioserotypes was provided for 1,682 (28.3%) confirmed cases. The most common bioserotypes were 4/O3 (83.2%) and 2/O9 (15.3%). Comparing these data with those for 2020, there was a slight decrease in the reporting of biotype 4/O3, from 87.7% in 2020 to 83.2% in 2021, while reporting of biotype 2/O9 increased from 10.9% in 2020 to 15.3% in 2021.

Eleven MSs reported a total of 113 Y. pseudotuberculosis cases in 2021.

#### 1.3.3. Yersinia in food

Statistics on the presence of *Yersinia* in the major food categories in 2021 and for 2017–2020 are summarised in Table 71 according to food type (i.e. RTE and non-RTE foods).

As in recent years, most RTE food samples analysed for the presence of *Yersinia* (N = 355) in 2021 belonged to 'meat and meat products' (232 samples; 65.3%) and were reported by four MSs. However, compared with 2020, the number of RTE samples decreased by 54%. A total of 3 (2.1%) sampling units of 'meat and meat products from pigs' were found to be positive for *Yersinia* in 2021. In the period 2017–2020, 121 (7.1%) sampling units positive for the presence of *Yersinia* were found in RTE food from 'meat and meat products', of which 111 (7.1%) sampling units in 'meat and mixed meat products from bovine animals and pigs' and 10 (10%) in 'mixed meat' samples. Compared with 2017–2020, no sampling was conducted in 2021 for the category 'mixed meat and meat products from bovine animals and pigs'.

Despite evidence supporting the need to sample RTE foods, few MSs reported monitoring this type of food for *Yersinia* to EFSA, even though they represent a direct risk to consumers.

**Table 71:** Occurrence of *Yersinia* in the main food categories, EU, 2021 and 2017–2020

		2021 <sup>(a)</sup>		2017–2020 <sup>(b)</sup>					
Food	N reporting MSs	N sampling units	N positive (%)	N reporting MSs	N sampling units	N positive (%)			
RTE food									
All	6	355	3 (0.85)	6	1,768	122 (6.9)			
Meat and meat products	4	232	3 (1.3)	4	1,707	121 (7.1)			
Meat and meat products from pigs	2	144	3 (2.1)	2	36	0 (0)			
Mixed meat and meat products from bovine animals and pigs	0	0	0 (0)	2	1,553	111 (7.1)			
Mixed meat	1	37	0 (0)	2	100	10 (10)			
Meat from other animals and raw cured (or seasoned) meat	2	51	0 (0)	3	18	0 (0)			
Milk and milk products	1	43	0 (0)	3	31	0 (0)			
Fruits and vegetables	1	55	0 (0)	4	10	0 (0)			
RTE salads	1	4	0 (0)	0	0	0 (0)			
Other processed food products and prepared dishes	1	21	0 (0)	2	4	1 (25.0)			
Other food	0	0	0 (0)	2	16	0 (0)			



//efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

Food		2021 <sup>(a)</sup>		2017–2020 <sup>(b)</sup>				
	N reporting MSs	N sampling units	N positive (%)	N reporting MSs	N sampling units	N positive (%)		
Non-RTE food								
All	6	736	38 (5.2)	8	4,092	283 (6.9)		
Meat and meat products	5	698	38 (5.4)	7	3,722	254 (6.8)		
Milk and milk products	1	23	0 (0)	3	238	29 (12.2)		
Other food	2	15	0 (0)	5	132	0 (0)		
Fresh meat <sup>(c)</sup>								
All fresh meat	5	418	29 (6.9)	7	4,026	310 (7.7)		
Fresh meat from pigs	5	366	26 (7.1)	5	2,400	148 (6.2)		
Fresh meat from bovine animals	1	10	0 (0)	3	34	1 (2.9)		
Fresh meat from bovine animals and pigs	0	0	0 (0)	2	1,401	129 (9.2)		
Other fresh meat	2	42	3 (7.1)	3	191	32 (16.8)		

MSs: Member States; RTE: 'ready-to-eat'. Fresh meat includes RTE and non-RTE food.

Other RTE food includes: Bakery products, Cereals and nuts - RTE, Chocolate, Other food - RTE

Other non-RTE food includes: Cereals, dried seeds – non-RTE, Egg and egg products – non-RTE, Fish and fishery products – non-RTE, Fruits, vegetables and juices – non-RTE, Foodstuffs intended for special nutritional uses – non-RTE, Other processed food products and prepared dishes – non-RTE and Seeds, sprouted – non-RTE.

- (a): Data from the United Kingdom (Northern Ireland) are taken into account for 2021. In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/Northern Ireland, the EU requirements on data sampling are also applicable to Northern Ireland.
- (b): Data from the United Kingdom are taken into account for 2017–2019, because the United Kingdom was an EU MS, but it became a third country on 1 February 2020.
- (c): Fresh meat sampling units are a subset of the two main categories above.

Regarding non-RTE foods, sampling and testing results were reported in 2021 by six MSs. As in recent years, most non-RTE food samples analysed for the presence of *Yersinia* (N = 736) belonged to 'meat and meat products' (698 samples; 94.8%). A total of 38 (5.4%) sampling units of 'meat and meat products' were positive for *Yersinia*. *Yersinia* was detected in 26 (7.1%) sampling units of 'fresh meat from pigs' (N = 366) belonging to the food category 'all fresh meat' (N = 418).

## 1.3.4. Yersinia in animals

Table 72 summarises the reported occurrence of *Yersinia* in animals for the year 2021 in MSs and non-MS countries. In 2021, seven MSs and one non-MS country reported monitoring data on the presence of *Yersinia* in animals. A total of 27,756 animals were analysed in EU for *Yersinia*. The MSs sampled different animal species, mainly cattle, small ruminants, pigs and pet animals with 223 (0.8%) positive animals. The non-MS country sampled mainly pet animals (N = 1,341) with only one positive sample (0.07%).

**Table 72:** Summary of *Yersinia enterocolitica* statistics related to major animal species, reported by EU MSs and non-MS countries, 2021

Animals			EU N	Non-MS countries				
	N reporting countries	N sampling units tested <sup>(b)</sup>	Positive sampling units		N reporting countries	N sampling units	Positive sampling units	
		testea	N	%		tested <sup>(b)</sup>	N	%
Cattle (bovine animals)	5	19,206	106	0.55	1	39	0	0
Small ruminants (sheep and goats)	4	3,531	96	2.7	1	18	0	0
Pigs	5	2,164	2	0.09	1	11	0	0

Animals		EU N	Non-MS countries					
	N reporting countries	N sampling units	Positive sampling units		N reporting countries	N sampling units tested <sup>(b)</sup>	Positive sampling units	
		tested <sup>(b)</sup>	N	%		testea	N	%
Other domestic/farmed animals	3	266	0	0	1	90	0	0
Pet animals	4	159	6	3.8	1	1,341	1	0.07
Wild animals (e.g. birds, roe and fallow deer, wild boar, foxes)	3	448	9	2.0	0	0	0	0
Zoo animals	5	80	4	5.0	0	0	0	0
Other (unspecified habitat)	4	1,902	0	0	1	6	0	0
Total	7	27,756	223	0.80	1	1,494	1	0.07

Other domestic\farmed animals: Alpacas – farmed, Poultry unspecified, Solipeds domestic – horses, Wild ducks – farmed, Camels – farmed, rabbits – farmed, Reindeers – farmed.

Pet animals: Cats – pet animals, Dogs – pet animals, Other animals – exotic pet animals, Rabbits – pet animals, Budgerigars – pet animals, Canaries – pet animals, Chinchillas – pet animals, Ferrets – pet animals, Guinea pigs – pet animals, Parrots – pet animals, Rats – pet animals, Reptiles – pet animals, Turtles – pet animals.

Wild: Badgers – wild, Birds – wild, Cantabrian chamois – wild, Deer – wild – fallow deer, Deer – wild – roe deer, Falcons – wild, Ferrets – wild, Foxes – wild, Martens – wild, Otters – wild, Rats – wild, Squirrels – wild, Steinbocks – wild, Turtles – wild, Wild boars – wild, Wolves – wild.

Zoo animals: Fish, Kangaroos – zoo animal, Monkeys – zoo animal, Penguins – zoo animals, Snakes – zoo animal, Wild cats (*Felis silvestris*) – zoo animals, Zoo animals, all.

Other (unspecified habitat): Fish, Cats, Deer, Hares, Rabbits, Geese, Kangaroos, Oscine birds and Pigeons.

- (a): Data from the United Kingdom (Northern Ireland) are taken into account for 2021. In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/Northern Ireland, EU requirements on data sampling are also applicable to Northern Ireland.
- (b): The summary statistics were obtained summing all sampling units (single and batch samples).

#### 1.4. Discussion

In 2021, yersiniosis was the third most commonly reported foodborne zoonotic disease in the EU. The EU notification rate for yersiniosis remained stable even during the COVID-19 pandemic when compared with other foodborne diseases and the overall trend for yersiniosis in 2017–2021 showed no statistically significant increase or decrease. The lowest number of yersiniosis cases were reported in 2020, which could be partially explained by the impact of the COVID-19 pandemic on national healthcare systems, and less importantly by the withdrawal of the United Kingdom from the EU. The EU notification rate in 2020, remained comparable with the previous years owing to the withdrawal of the United Kingdom from the EU. In 2021, the number of reported cases was comparable to those reported before the pandemic and the EU notification rate of yersiniosis increased compared to both 2020 and to the pre-pandemic period (2017–2019).

Among the two pathogenic species that are notified, *Y. enterocolitica* and *Y. pseudotuberculosis*, the first caused the majority (98.1%) of human infections. Information on biotypes would allow for better characterisation of the epidemiology of *Y. enterocolitica* infection in humans and a better investigation of relevant animal sources (Espenhain et al., 2019; Gruber et al., 2021). In 2021 as in previous years, the most common biotypes in humans were 4/O3 and 2/O9. In 2021, few MSs reported data on *Yersinia* sampling activities in food and animals. This is probably due to the lack of mandatory monitoring plans for non-human sources, resulting in significant differences between MSs in their approach to monitoring *Yersinia* in food and animals. The lack of homogeneity makes statistical analysis and monitoring of trends impossible. An EFSA scientific report suggested technical specifications for the harmonised monitoring and reporting of *Y. enterocolitica* in slaughter pigs in the EU (EFSA, 2009).

In 2021, only six MSs provided data on RTE food for *Yersinia* and the number of samples analysed was rather low (N = 355) compared with 2020 (N = 766), with a proportion of positive samples of 0.85% compared with 5.2% in 2020. Instead, it would be beneficial to sample more RTE foods that are consumed without any processing to reduce or eliminate *Yersinia* contamination.



18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/erms

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons

Yersinia enterocolitica was isolated in all foods that were found to be positive. Information on Y. enterocolitica biotypes was provided for a small fraction of data, although documenting trends and sources of Yersinia along the food chain, including reporting of information on Y. enterocolitica biotypes (EFSA BIOHAZ Panel, 2007), is essential to the overall goal of reducing yersiniosis at both the primary production level and at other critical points along the food production chain.

In this context, it would be useful to adopt real-time PCR-based methods that are able to rapidly provide information on the pathogenicity of *Y. enterocolitica* that may be present, even though they do not provide an indication of the specific biotype. Such methods can also easily be used for the detection of strains isolated from faeces, food, feed and environmental samples (Hallanvuo et al., 2019). For this purpose, the ISO method (ISO, 2015) describes horizontal methods based on real-time PCR for detection of the pathogenic bioserotypes of *Y. enterocolitica* and one for detection of *Y. pseudotuberculosis* in products for human consumption, animal feeding stuffs and environmental samples (Rivas et al., 2021).

## 2. Toxoplasma gondii

Summary data substantiating this chapter, as well as additional information on related projects and internet sources are published for this report on the EFSA Knowledge Junction on Zenodo here. Summary statistics on human surveillance data with downloadable files are retrievable using the ECDC Surveillance Atlas of Infectious Diseases here.

# 2.1. Key facts

- Only confirmed cases of congenital toxoplasmosis are reported to ECDC, with a two-year delay
  in human data analyses at the EU level as data from France are reported to TESSy with a twoyear delay.
- In 2020, the number of confirmed cases of congenital toxoplasmosis was 133, corresponding to an EU notification rate of 5.1 cases per 100,000 live births. Compared with the rates in 2019 with and without data from the United Kingdom (5.1 and 6.2 per 100,000 live births), this is unchanged or decreased by 17.7%, respectively.
- In 2020, France accounted for 82.7% of reported cases of congenital toxoplasmosis due to compulsory screening of pregnant women.
- Overall, the number of human cases of congenital toxoplasmosis in the EU has shown a
  constant decrease in the 2016–2020 period, mainly due to the reduction in cases reported by a
  single MS (France), but also due to the COVID-19 pandemic and withdrawal of the United
  Kingdom from the EU.
- In total, 14 MSs and three non-MSs reported 2021 monitoring data on *Toxoplasma gondii* infections in animals. Most animals tested were sheep and goats, which also showed the highest overall prevalence of *T. gondii* infections in animals (16.8%), as reported by 12 MSs. Most samples were obtained from clinical investigations. It is not possible to accurately estimate the prevalence of *T. gondii* infections in animals due to the use of different diagnostic methods, the different sampling schemes in the MSs and the lack of information on the animals' ages and rearing conditions.

# 2.2. Surveillance and monitoring of *Toxoplasma* in the EU

#### **2.2.1.** Humans

In 2020, surveillance of toxoplasmosis was mandatory in 18 MSs and in one (France) it was voluntary. Data from France are reported to TESSy with a 2-year delay. Eight MSs (Austria, Belgium, Denmark, Greece, Italy, the Netherlands, Portugal and Sweden) do not have a surveillance system for toxoplasmosis.

All reporting MSs except one (Spain) had a comprehensive surveillance system with full national coverage. Spain did not have national surveillance and could not provide any estimate for population coverage, so no notification rate was calculated. In 2020, Spain has not received data from all regions that normally report, so the case number might not be complete. Case-based data were reported by all MSs except Bulgaria, which reported aggregated data. The EU case definition was used by 17 MSs and two MSs (France and Germany) reported use of different case definitions.



Six MSs (Austria, Belgium, France, Greece, Slovakia and Slovenia) have active surveillance of congenital cases, with compulsory screening of pregnant women. Austria and Belgium, however, do not report to TESSy. In the case of Austria, the disease is not notifiable and official data are therefore lacking. In Belgium, there are no clear recommendations on the follow-up of seroconversion cases during pregnancy. In 2020, Greece did not report to TESSy. Four MSs (Bulgaria, Czechia, Germany and Hungary) have voluntary screening (ECDC, 2022).

On 1 February 2020, the United Kingdom became a third country, whereas before it was a MS. Human data were not collected from the United Kingdom for 2020 by ECDC.

#### 2.2.2. Animals and food

There are no EU regulations concerning the surveillance and monitoring of *T. gondii* in animals and food. Therefore, the available and reported information relies on national legislation and whether the countries have a mandatory reporting system following the detection of *T. gondii*. As a result, data allow only for descriptive summaries at the EU level (Table 73). The main animal species tested are those intended for human consumption (small ruminants (goats and sheep), cattle, pigs) as well as pet animals (cats and dogs), using samples from aborted animals (ruminants) or clinically suspect animals. Mainly blood samples, but also samples from tissues and organs, are analysed either by direct methods, such as PCR, histology and immunohistochemistry (IHC), or by indirect antibody detection methods, including enzyme-linked immunosorbent assay (ELISA), latex agglutination test (LAT), direct agglutination (DA) and immunofluorescence assay (IFA). For food testing, meat is the main matrix to be analysed either by PCR or by direct agglutination (DA).

#### 2.3. Results

## 2.3.1. Human congenital toxoplasmosis

In 2020, 19 MSs reported congenital toxoplasmosis data (Table 73) with a total of 133 confirmed cases. Sixteen MSs reported zero cases, whereas eight MSs (Austria, Belgium, Denmark, Greece, Italy, the Netherlands, Portugal and Sweden) did not report congenital toxoplasmosis at the EU level (Table 73). The notification rate was 5.1 per 100,000 live births in the EU, with the highest rate in France (15.5), followed by Poland (2.5) and Germany (1.8) (Table 73).

France accounted for 82.7% of all reported cases in the EU, followed by Germany and Poland. France regularly reports the highest number of congenital toxoplasmosis cases in the EU, whereas the other two MSs accounted for 17.3% of the cases reported in the EU.

Between 2016 and 2020, a gradual decrease in the number of reported cases was observed from France (43.6%) and Poland (55%, excluding data from 2018). A similar decrease was not observed for Germany.

As a result of fewer cases reported by France (and Poland), in 2020 the notification rate in the EU decreased by 17.7% compared with 2019, without data from the United Kingdom (Table 73), whereas no decrease was observed compared with 2019, including data from the United Kingdom.

In 2020, information on hospitalisation was provided by one MS (Poland) on 9 cases, representing 100% of cases reported by the MS. Cases with known fatal outcomes were 5 out of 109 (case fatality of 4.6%), with all fatal cases reported by France. Germany reported one case as imported, with Turkey being the probable country of infection.

18314732, 2022, 12, Downloaded from https://fs.aonlinelibrary.wiley.com/inchibrary.wiley.com/inchibrary.wiley.com/em-and-conditions) on Wiley Online Library on [18/12/2023], See the Terms and Conditions (http://oinineibrary.wiley.com/em-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons



**Table 73:** Reported confirmed human cases of congenital toxoplasmosis and notification rates per 100,000 live births in EU MSs and non-MS countries, by country and year, 2016–2020

		2019		20	18	20	17	2016				
Country	National Data coverage <sup>(a)</sup> format <sup>(a)</sup>		Confirmed cases and rates									
			Cases	Rate								
Austria	_	-	-	_	_	_	_	_	-	-	_	_
Belgium	_	-	_	_	_	_	_	_	_	_	_	_
Bulgaria	Υ	Α	0	0	0	0	0	0	2	3.1	0	0
Croatia	Υ	С	0	0	0	0	1	2.7	0	0	0	0
Cyprus	Y	С	0	0	0	0	0	0	0	0	0	0
Czechia	Υ	С	0	0	1	0.9	0	0	2	1.8	0	0
Denmark	_	_	_	_	_	_	_	_	_	_	_	_
Estonia	Υ	С	0	0	0	0	1	7.2	0	0	0	0
Finland	Υ	С	0	0	0	0	0	0	0	0	1	1.8
France	Υ	С	110	15.5	134	18.8	151	20.8	153	20.7	195	25.9
Germany	Υ	С	14	1.8	17	2.2	18	2.3	8	1.0	10	1.3
Greece	_	_	_	_	0	0	0	0	0	0	_	_
Hungary	Υ	С	0	0	1	1.1	0	0	0	0	0	0
Ireland	Y	С	0	0	0	0	0	0	0	0	0	0
Italy	_	_	_	_	_	_	_	_	_	_	_	_
Latvia	Y	С	0	0	1	5.2	0	0	0	0	0	0
Lithuania	Υ	С	0	0	0	0	0	0	0	0	0	0
Luxembourg	Y	С	0	0	0	0	0	0	0	0	0	0
Malta	Y	С	0	0	0	0	0	0	0	0	0	0
Netherlands	_	_	_	_	_	_	_	_	_	_	_	_
Poland	Y	С	9	2.5	14	3.7	25	6.3	18	4.8	20	5.5
Portugal	_	_	_	_	_	_	_	_	_	_	_	_
Romania	Y	С	0	0	0	0	1	0.48	0	0	0	0
Slovakia	Υ	С	0	0	0	0	0	0	0	0	2	3.6
Slovenia	Y	С	0	0	1	5.1	2	9.9	2	9.8	1	4.8
Spain <sup>(b),(c)</sup>	N	С	0	_	0	_	2	_	3	_	5	_
Sweden	_	_	_	_	_	_	_	_	_	_	_	_
EU Total 27	_	_	133	5.1	169	6.2	201	7.1	188	6.6	234	8.7
United Kingdom	_	_	_	_	7	1.0	7	0.9	7	0.91	8	1.0
EU Total <sup>(d)</sup>	_	_	133	5.1	176	5.1	208	5.8	195	5.4	242	6.9
Iceland	Υ	С	0	0	0	0	0	0	0	0	0	0
Norway	_	_	_	_	_	_	-	_	-	_	_	_
Liechtenstein	_	_	_	_	_	_	_	_	_	_	_	_
Switzerland	_	_	_	_								

<sup>-:</sup> Data not reported.

18314732, 2022, 12, Downloaded from https://efs.aonlinelibrary.wiley.com/doi/10.2903/j.efs.a.2022.7666 by Isituto Zooprofilation Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18122023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for release to exceed the applicable Creative Commonstation (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for release to exceed the applicable Creative Commonstation (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for release to exceed the applicable Creative Commonstation (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for release to exceed the applicable Creative Commonstation (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for release to exceed the applicable Creative Commonstation (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for release to exceed the applicable Creative Commonstation (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for release to exceed the applicable Creative Commonstation (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for release to exceed the applicable Creative Commonstation (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for release to exceed the applicable Creative Commonstation (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for release to exceed the applicable Creative Commonstation (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for release to exceed the applicable Creative Commonstation (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for release to exceed the applicable Creative Commonstation (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for release to exceed the applicable Creative Commonstatio

<sup>(</sup>a): Y: yes; N: no; A: aggregated data; C: case-based data.

<sup>(</sup>b): Notification rate was not calculated since information on estimated coverage was not available.

<sup>(</sup>c): Data not complete for 2020–2021, rate not estimated.

<sup>(</sup>d): Cases reported by the United Kingdom for the period 2016–2019 were also considered for this estimate (EU-28). When the United Kingdom data were collected for the period 2017–2019, the United Kingdom was an EU MS, but it became a third country on 1 February 2020.

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903j.efsa.2022.7666 by Istituo Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common



#### 2.3.2. Toxoplasma gondii in food and animals

#### Toxoplasma gondii in food

One MS (Poland) submitted monitoring results for *T. gondii* in non-RTE food in 2021. In total, 694 samples were reported from meat products from pigs (641) and bovine animals (53). Seventy-three (11.4%) and five (9.4%) samples were positive from pigs and from bovine animals, respectively.

#### Toxoplasma gondii in animals

Monitoring data on *T. gondii* in livestock, pet or zoo animals and wildlife were provided by 14 MSs, and by three non-MSs (North Macedonia, Norway and Switzerland).

In small ruminants (sheep and goats), 12 MSs and three non-MSs (North Macedonia, Norway and Switzerland) reported data. In total, 4,552 animals were tested and 762 were found to be positive (16.7%). In cattle, seven MSs (Austria, Germany, Ireland, Italy, Latvia, Slovakia and the United Kingdom (Northern Ireland)) and one non-MS (Switzerland) reported data on Toxoplasma. In total, 728 animals were tested and 18 were found to be positive (2.5%). In pigs, five MSs (Austria, Germany, Ireland, Italy and the United Kingdom (Northern Ireland)) reported monitoring data that are found in Table 74. For other livestock (solipeds, alpacas, fowl and water buffaloes), four MSs (Austria, Germany, Italy and the United Kingdom (Northern Ireland)) reported monitoring data: in total, 371 animals were tested and four (1.1%) were found to be positive. In pet animals (cats, dogs, birds and rodents), eight MSs (Austria, Finland, Germany, Italy, Latvia, the Netherlands, Slovakia and Slovenia) and one non-MS (Switzerland) tested in total 5,463 animals (3,640 cats, 1,776 dogs, 33 birds and 14 rodents). There were 778 (14.2%) positive samples, 428 (11.7%) samples from cats, 349 (19.6%) samples from dogs and 1 (3%) sample from birds obtained mainly from clinical investigations. As regards zoo animals, five MSs (Austria, Finland, Germany, Italy and Slovakia) and two non-MSs (North Macedonia and Switzerland) tested in total 810 animals and 14 (1.7%) were positive. Three MSs (Germany, Italy and Slovakia) and one non-MS (Switzerland) reported on testing for T. gondii in wildlife. In total, 1,299 animals (mainly from Italy and Slovakia) were tested and 150 were positive (11.5%).

**Table 74:** Summary of *Toxoplasma gondii* statistics in the main animal species, EU, 2017–2021

Animals <sup>(a)</sup>	2021 <sup>(a)</sup>	2020	2019 <sup>(b)</sup>	2018 <sup>(b)</sup>	2017 <sup>(b)</sup>	Data source
Small ruminants						
Number of tested animals	4,525	6,113	12,120	6,756	6,410	EFSA
% of positive animals	16.8	21.3	13.5	18.3	18.3	EFSA
Number of reporting MSs	12	10	11	12	13	EFSA
Bovine animals						
Number of tested animals	726	254	664	158	2,163	EFSA
% of positive animals	2.5	9.8	9.2	27.8	10.6	EFSA
Number of reporting MSs	7	4	6	6	7	EFSA
Pigs						
Number of tested animals	599	948	1,108	263	689	EFSA
% of positive animals	5.0	9.7	11.7	22.1	15.2	EFSA
Number of reporting MSs	5	3	4	4	4	EFSA
Cats						
Number of tested animals	3,275	1,880	1,525	1,382	690	EFSA
% of positive animals	9.8	6.5	5.2	4.7	7.5	EFSA
Number of reporting MSs	8	6	8	9	8	EFSA

Summary statistics referring to MSs were obtained by totalling all sampling units (single animals, slaughter animal batches, and herds or flocks).

For the summary statistics, indirect and direct diagnostic methods were taken together to calculate the proportion of positive units.

MS: Member State.

- (a): Data from the United Kingdom (Northern Ireland) are taken into account for 2021. In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/Northern Ireland, the EU requirements on data sampling are also applicable to Northern Ireland.
- (b): Data from the United Kingdom are taken into account for 2017–2019, because the United Kingdom was an EU MS, but it became a third country on 1 February 2020.



#### 2.4. Discussion

Cases of congenital toxoplasmosis in the EU are strongly biased by the high reporting rate of France, which has accounted for most reported cases since 2009 (ECDC Surveillance Atlas of Infectious Diseases), representing 83.3% to 75.1% of overall EU cases in 2016–2020 (excluding data from the United Kingdom). The high reporting rate for France reflects systematic screening for toxoplasmosis in pregnant women first established in 1978 and mandatory since 1992. Seronegative women are followed up during pregnancy to detect seroconversion early, and congenital toxoplasmosis cases are laboratory confirmed. The reported constant decrease in cases in the EU clearly mirrors the lower number of cases reported by France in the 2016-2020 period. In this 5-year period, the most remarkable decrease in reported cases occurred in 2016 and 2017 (20.7% and 21.5, respectively) and continued in subsequent years, with the lowest rate in 2020. Decreased seroprevalence in pregnant women in France (from 54% in 1995, to 31% in 2016) and a decreased number of seroconversions during pregnancy (from 5.4 per 1,000 at-risk pregnancies in 1995, to 2.1 in 2010, and expected to be 1.6 by 2020) have been reported, and may explain the decreasing number of reported cases of congenital toxoplasmosis in France (Robinson et al., 2021). Interestingly, the number of reported cases in France decreased by 11.2% and 17.9% between 2018 and 2019 and between 2019 and 2020, respectively, suggesting an additional effect of the COVID-19 pandemic in the case reporting decrease in 2020. An educational campaign for pregnant women and reduced exposure to contaminated raw/ undercooked meat (e.g. changes in food habits and improved hygiene practices in meat production) or other raw foods at risk of contamination (e.g. fresh produce, molluscs and raw milk) have likely contributed to the reduced incidence of *T. gondii* infection during pregnancy in France, reflecting the lower number of congenital toxoplasmosis cases (Opsteegh et al., 2015; Robinson et al., 2021). By contrast, surveillance for congenital toxoplasmosis among the other MSs is highly variable, with countries reporting zero cases or simply not reporting to ECDC, or countries lacking any surveillance. This may reflect discussions about the cost-effectiveness of screening pregnant women in preventing or reducing the impact of congenital toxoplasmosis.

Due to imbalanced data, it is not possible to estimate the prevalence of congenital toxoplasmosis in the EU, limiting assessment of the burden of this form of the disease. In addition to prenatal screening, sensitive and effective postnatal diagnostic screening should be implemented and improved in the EU in order to detect cases of congenital toxoplasmosis. This is particularly important as most prenatal infections are sub-clinical at birth or when maternal anti-toxoplasma treatment is performed, which may decrease diagnostic sensitivity, thus generating false negatives (Guegan et al., 2021). Primary infection in pregnancy, particularly if early in gestation, and even in the asymptomatic form, is of particular concern. Even though the risk of congenital infection increases if the mother becomes infected late in pregnancy (third trimester), outcomes of congenital disease are much more severe if the mother became infected in the first trimester. This can result in abortion, still-birth, perinatal death or congenital diseases with immediate or late (up to adolescence) manifestations, including ocular diseases, seizures and learning disabilities. All possible strategies for the prevention of congenital toxoplasmosis, including appropriate information to pregnant woman and active screening, should be reinforced.

The 2021 monitoring data from animals reported by MSs show that *T. gondii* is present in most livestock species across the EU, as well as in pet/zoo animals and wildlife. The limitations of these surveillance data preclude any trend watching or prevalence trend analysis in animals.

The current European surveillance system of *T. gondii* in animals is strongly affected by several important limitations: (i) the small number of tested animals, (ii) the use of different indirect and direct detection methods, which have not been validated by an independent body in most cases; (iii) unknown age of the tested animals; and (iv) no information on the type of husbandry system (housing). Furthermore, there is no relationship between the presence of anti-*T. gondii* antibodies and infecting parasites in cattle and horses (Aroussi et al., 2015; Blaga et al., 2019; Opsteegh et al., 2019). For pigs, poultry and small ruminants, serological methods could be useful for the detection of high-risk animals or herds, but not as an indicator of infection in individual animals, as the agreement between direct and indirect methods was estimated to be low to moderate.

The above-mentioned limitations associated with toxoplasmosis detection and diagnosis, and surveillance rules, do not allow for direct comparison of the reported data across MSs.



#### 3. **Rabies**

Summary data substantiating this chapter, as well as additional information on related projects and internet sources are published for this report on the EFSA Knowledge Junction on Zenodo here. Summary statistics on human surveillance data with downloadable files are retrievable using the ECDC Surveillance Atlas of Infectious Diseases here.

#### 3.1. **Key facts**

- In 2021, EU MSs and non-MS countries reported no human lyssavirus infections, as in 2020. However, travel-associated human rabies cases occurred regularly in Europe, as reported in recent years (N = 4 in 2019, N = 1 per year in the 2017–2018 period).
- In animals excluding bats, a total of 118 cases of rabies of autochthonous origin were reported by two MSs: 113 cases in Poland (96 foxes, 2 wild roe deer, 2 martens, 2 raccoon dogs, 7 cats, one stray cat, 2 dogs and one stray dog) and five cases in Romania (four cows and one fox). The total number of reported indigenous rabies cases in non-flying animals in the EU increased in 2021 compared with previous years (N = 12 in 2020, N = 5 in 2019; N = 8 in 2018; N = 6 in 2017). This is due to a rabies outbreak since 2021 in a Polish region that had been rabiesfree for more than 16 years.
- Surveillance data on lyssavirus in bats were reported by 16 EU MSs. Four MSs reported 29 positive results for lyssavirus, mainly European bat 1 lyssavirus (EBLV-1).
- A case of rabies was reported by Germany in an illegally imported dog infected with a rabies virus (RABV) strain.

#### 3.2. Surveillance and monitoring of rabies in the EU

#### 3.2.1. **Humans**

Reporting of rabies is mandatory in all 27 MSs. In 2021, all countries except Denmark reported case-based data. Twenty-three MSs used the EU case definition, three countries (Denmark, Germany and Italy) used another case definition and France did not specify the case definition used. Disease surveillance is comprehensive in all reporting countries.

#### 3.2.2. **Animals**

The objective of **passive rabies surveillance** is to detect the presence and the geographic distribution of the virus over time, to allow timely dissemination of information for immediate integrated control actions among different sectors, such as the public health and veterinary sectors. For rabies-free countries, surveillance aims to confirm the absence of the disease. In accordance with Regulation (EU) No 652/2014<sup>36</sup> and Commission Delegated Regulation (EU) No 2020/689<sup>37</sup>, multiannual programmes for eradication of rabies may be co-financed by the EU. In 2021, 12 MSs had approved elimination, control and surveillance programmes for rabies and oral rabies vaccination (ORV) campaigns were conducted in 8 MSs (Croatia, Estonia, Finland, Hungary, Latvia, Lithuania, Poland and Slovakia), as well as in some of the EU-bordering countries. Surveillance of rabies is carried out by sampling and testing 'indicator animals'; these are wild or domestic animals (foxes, raccoon dogs, jackals, badgers, dogs, cattle, cats, sheep, equines, goats, etc.) that are found dead (including road-killed) and/or suspect animals, i.e. animals showing neurological clinical signs or abnormal behaviour compatible with rabies like biting, licking a wound or scratching a human being in the absence of clear neurological signs.

<sup>37</sup> Commission Delegated Regulation (EU) 2020/689 of 17 December 2019 supplementing Regulation (EU) 2016/429 of the European Parliament and of the Council as regards rules for surveillance, eradication programmes, and disease-free status for

certain listed and emerging diseases.

<sup>&</sup>lt;sup>36</sup> Regulation (EU) No 652/2014 of the European Parliament and of the Council of 15 May 2014 laying down provisions for the management of expenditure relating to the food chain, animal health and animal welfare, and relating to plant health and plant reproductive material, amending Council Directives 98/56/EC, 2000/29/EC and 2008/90/EC, Regulations (EC) No 178/2002, (EC) No 882/2004 and (EC) No 396/2005 of the European Parliament and of the Council, Directive 2009/128/EC of the European Parliament and of the Council and Regulation (EC) No 1107/2009 of the European Parliament and of the Council and repealing Council Decisions 66/399/EEC, 76/894/EEC and 2009/470/EC. OJ L 189, 27.6.2014, pp. 1-32.



18314732, 2022, 12, Downloaded from https://disa.onlinelibrary.wiley.com/doi/10.2903/j.disa.2022.7666 by Isituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.2903/j.disa.2022.7666 by Isituto Zooprofilation Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.2903/j.disa.2022.7666 by Isituto Zooprofilation Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.2903/j.disa.2022.7666 by Isituto Zooprofilation Sperimenta dell'Umbria e delle Marche (IZSUM). Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.2903/j.disa.2022.7666 by Isituto Zooprofilation Sperimenta dell'Umbria e dell'Um

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons

To monitor the efficacy of the ORV campaigns, healthy animals of the wild species targeted by oral vaccination, which are foxes, raccoon dogs and golden jackals, are hunted. These animals' carcases are used to determine immunity and oral vaccine bait uptake. This specific **active rabies surveillance** is traditionally designated as 'ORV monitoring' or 'monitoring'. These hunted animals can also be tested for rabies and very few of them (below 5%) are usually found to be positive for the disease.

Imported or travel-related companion animals (mainly dogs and cats) from territories and non-EU countries not included in Annex II of Commission Implementing Regulation (EU) No 577/2013<sup>38</sup> are currently tested for rabies virus neutralising antibodies.

In accordance with CIR (EU) 2020/2002, EU MSs must notify outbreaks of infection with rabies virus in non-flying animals to the EU ADIS<sup>3</sup>; these are the animal species and groups of species Carnivora, Bovidae, Suidae, Equidae, Cervidae and Camelidae, but not Chiroptera (bats).

The data reported here include all animals tested for rabies, collected for disease surveillance and for ORV monitoring (active surveillance) purposes.

#### 3.3. Results

# 3.3.1. Overview of key statistics, EU, 2017–2021

A summary of EU-level rabies statistics in humans and in wild and domestic animals is shown in Table 75. For animals, the total number of samples analysed for passive surveillance from foxes, raccoon dogs, golden jackals, dogs, cats and bats, as well as the number of MSs from which these samples originated, are shown. An increase was observed in the number of tested samples of foxes, which are the main reservoir of the virus, compared with 2020. The number of tested raccoon dogs remained stable, while the number of golden jackals more than doubled in 2021 compared with 2020. In 2021, the number of tested bats remained stable as compared with 2020, and the substantial decrease observed in 2020 was confirmed for 2021 as compared with the 2017–2019 period. For cats, dogs and farmed animals, the number of tested samples remained stable in 2021 as compared with 2020.

Table 75 also displays the active surveillance (ORV monitoring) statistics for ORV target species (foxes, raccoon dogs and golden jackals). The number of tested foxes decreased considerably, while the number of tested raccoon dogs and golden jackals increased in 2021, as compared with the previous years.

**Table 75:** Summary of rabies lyssavirus statistics related to humans and the main animal reservoirs, EU, 2017–2021

,	2021 <sup>(a)</sup>	2020	2019 <sup>(b)</sup>	2018 <sup>(b)</sup>	2017 <sup>(b)</sup>	Data
						Source
Humans						
Total number of confirmed cases	0	0	4	1	1	ECDC
Total number of confirmed cases/100,000 population (notification rates)	0	0	0	0	0	ECDC
Number of reporting countries	26	26	28	28	28	ECDC
Infection acquired in the EU	0	0	1	0	0	ECDC
Infection acquired outside the EU	0	0	3	1	1	ECDC
Unknown travel status or unknown country of infection	0	0	0	0	0	ECDC
Animals under passive surveillance						
Dogs (Canis lupus familiaris)						
Number of tested animals	1,838	1,732	1,901	2,097	2,334	EFSA
Number of positive animals	4	4	_	1	1	EFSA
Number of reporting MSs	21	22	22	23	22	EFSA

<sup>&</sup>lt;sup>38</sup> Commission Implementing Regulation (EU) No 577/2013 of 28 June 2013 on the model identification documents for the non-commercial movement of dogs, cats and ferrets, the establishment of lists of territories and third countries and the format, layout and language requirements of the declarations attesting compliance with certain conditions provided for in Regulation (EU) No. 576/2013 of the European Parliament and of the Council. OJ L 178, 28.6.2013, pp. 109–148.



dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

	2021 <sup>(a)</sup>	2020	2019 <sup>(b)</sup>	2018 <sup>(b)</sup>	2017 <sup>(b)</sup>	Data Source
Cats (Felis catus)						
Number of tested animals	2,335	2,440	2,389	2,661	2,722	EFSA
Number of positive animals	8	2	_	_	1	EFSA
Number of reporting MSs	20	21	22	21	23	EFSA
Farmed mammals <sup>(c)</sup>						
Number of tested animals	406	392	394	570	796	EFSA
Number of positive animals	4	3	1	1	3	EFSA
Number of reporting MSs	17	17	15	17	17	EFSA
Wild animals – Red foxes ( <i>Vulpes</i>	vulpes)					
Number of tested animals	12,907	9,805	5,338	5,833	10,808	EFSA
Number of positive animals	97	6	2	6	2	EFSA
Number of reporting MSs	17	18	16	16	17	EFSA
Wild animals – Raccoon dogs ( <i>Ny</i> o	ctereutes procyo	noides)				
Number of tested animals	1,339	1,214	1,241	1,335	712	EFSA
Number of positive animals	2	_	_	_	_	EFSA
Number of reporting MSs	6	6	7	7	6	EFSA
Wild animals – Jackals ( <i>Canis aur</i> c	eus)					
Number of tested animals	230	102	42	44	1,000	EFSA
Number of positive animals	_	_	_	_	_	EFSA
Number of reporting MSs	5	6	5	4	5	EFSA
Wild animals – Bats (order Chirop	tera)					
Number of tested animals	1,316	1,308	2,069	2,278	2,079	EFSA
Number of positive animals	29	31	39	45	39	EFSA
Number of reporting MSs	16	15	19	18	19	EFSA
Animals under active surveillance	(ORV monitorin	g) <sup>(d)</sup>				
Red foxes (Vulpes vulpes)						
Number of tested animals	10,581	14,416	17,805	15,737	19,677	EFSA
Number of positive animals	_	_	1	_	_	EFSA
Number of reporting MSs	9	10	9	9	9	EFSA
Raccoon dogs ( <i>Nyctereutes procy</i>	onoides)					
Number of tested animals	369	324	301	23	280	EFSA
Number of positive animals	_	_	_	_	_	EFSA
Number of reporting MSs	4	4	3	2	3	EFSA
Jackals ( <i>Canis aureus</i> )						
Number of tested animals	1,499	1,319	1,045	1,304	870	EFSA
Number of positive animals		_	_	_	_	EFSA
Number of reporting MSs	3	3	2	3	2	EFSA

ECDC: European Centre for Disease Prevention and Control; EFSA: European Food Safety Authority; MSs: Member States. ORV oral rabies vaccination.

#### 3.3.2. Humans

EU MSs and non-MS countries reported no human lyssavirus infections for 2021, as was observed in 2020. In the 2017–2019 period, travel-associated cases were reported each year from EU MSs. In 2017, France reported one travel-associated case, with exposure occurring in Sri Lanka. In 2018, the

<sup>(</sup>a): Data on animal samples from the United Kingdom (Northern Ireland) are taken into account for 2021. In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/ Northern Ireland, the EU requirements on data sampling are also applicable to Northern Ireland.

<sup>(</sup>b): Data from the United Kingdom are taken into account for the 2017–2019 period because the United Kingdom was an EU MS, but it became a third country on 1 February 2020.

<sup>(</sup>c): Animals that are commonly raised or kept for breeding purposes or animal by-products.

<sup>(</sup>d): Samples collected in the framework of oral rabies vaccination monitoring (active surveillance).



onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico


Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

United Kingdom reported one human case following exposure in Morocco. In 2019, four human lyssavirus infections were reported by EU MSs. Three of these were travel-associated, reported by Italy, Latvia and Spain, and acquired in Tanzania, India and Morocco, respectively. Most of the human cases were linked to dog exposure. The fourth human infection in 2019 was reported in France and was due to European bat lyssavirus 1 (EBLV-1).

#### **3.3.3.** Animals

#### Rabies cases in wildlife

In 2021, 12,907 foxes (*Vulpes vulpes*) were tested using passive surveillance by 17 MSs. The majority of the tested samples (60.5%) were analysed by two MSs: Poland and Romania. In total, 97 cases of rabies in foxes were detected in the EU: 96 cases in Poland and one in Romania. The geographical distribution and number of cases in foxes, as well as a choropleth map of the total number of foxes sampled per MS for passive surveillance, are shown in Figure 35. Three non-EU countries (Norway, North Macedonia, and Switzerland) reported a total of 21 foxes tested under passive surveillance schemes and found none positive.



**Figure 35:** Choropleth map of the number of tested and positive foxes, and the geographical distribution of the rabies cases reported in foxes in EU MSs and non-EU countries, 2021

In 2021, 1,339 raccoon dogs (*Nyctereutes procyonoides*) were tested for rabies by six MSs (Estonia, Finland, Latvia, Lithuania, Poland and Slovakia). Most (93.4%) of these samples came from raccoon dogs originating from two MSs (Estonia and Latvia). Two tested samples (out of a total of 38)




onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico

dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

originating from Poland were positive for rabies. Regarding golden jackals, 89.5% of the samples tested came from two MSs (Bulgaria and Romania).

Seventeen MSs reported data of passive surveillance for 714 non-flying wild animals other than foxes, raccoon dogs or golden jackals. The other most widely tested wild animal species were deer and roe deer (152), badgers (148), martens (108) and raccoons (98). In Poland, two roe deer and two martens were tested positive for rabies. Other species tested included wolf, lynx, bear, ferret, hedgehog, mouse, mink, otter, polecat, rat, squirrel, wild boar, wolverine, coypus, moose, beaver, weasel and mole. All the animals tested negative for rabies.

In 2021, 16 MSs and two non-MSs reported surveillance data on bats. In total, 1,316 bats were investigated in the EU (Figure 36). Of these, 29 samples tested positive in four MSs: Germany (16 EBLV-1), France (5 EBLV-1), Poland (5 unspecified lyssavirus) and Spain (3 EBLV-1). Two non-MSs, Norway and Switzerland, analysed 4 and 18 bats, respectively and all samples tested negative.



For Germany and Spain, the geographical distribution of reported cases was not provided.

**Figure 36:** Choropleth map of the number of tested and positive bats, and the geographical distribution of the rabies cases reported in bats in EU MSs and non-EU countries, 2021

#### Rabies cases in domestic animals

In 2021, a total of 15 autochthonous domestic animals were tested positive for rabies. Romania reported four cases of rabies (unspecified lyssavirus) in cows. Poland recorded two cases in pet dogs, one case in a stray dog, seven cases in pet cats and one case in a stray cat. Except for the stray cat for which a rabies virus (RABV) strain was identified, all the other cases reported the strains as unspecified lyssavirus.




//efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico

dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

In Germany, one dog was reported positive for rabies. This unvaccinated 8-week-old puppy had been imported illegally from Turkey via Bulgaria. Genetic analysis revealed that it was an RABV strain showing a high sequence identity with a Turkish RABV fox isolate.

The geographical distribution and number of tested and reported cases in pets (dogs and cats) are shown in Figure 37




For Germany, the geographical distribution of the reported case was not provided.

**Figure 37:** Choropleth map of the number of tested and positive pets, and the geographical distribution of the rabies cases reported in pets in EU MSs and non-EU countries, 2021

In 2021, 21 MSs tested and reported 4,173 samples for dogs (1,838) and cats (2,335 from 20 MSs). The number of samples reported for both species remained similar to 2019 (1,732 dogs and 2,440 cats were reported in 2019). Three non-EU countries (Norway, North Macedonia and Switzerland) reported in total 72 tested dogs and 24 tested cats. None of them reported positive cases for rabies.

A total of 406 samples from farmed mammals (Figure 38) were tested by 17 MSs (reports included mainly cattle, small ruminants and domestic solipeds). The number of samples tested from domestic farmed mammals in 2021 was slightly higher than in 2020 (392 samples tested) but lower than in 2017 or and 2018.



**Figure 38:** Choropleth map of the number of tested and positive farmed animals, and the geographical distribution of the rabies cases reported in farmed animals in EU MSs and non-EU countries, 2021

## 3.4. Discussion

In Europe, human rabies is a rare disease, with the last autochthonous EU case of RABV human infection dating back to 2012 (Romania). Nowadays, the infection is mainly acquired abroad in countries where dog rabies is endemic and the development of the disease is due to the absence of pre-exposure prophylaxis or late/inappropriate/incomplete administration of post-exposure treatment. The illegal importation of pets also poses a constant risk of rabies introduction (Klevar et al., 2015). Another rare source of infection is through organ transplantation (Maier et al., 2010). The absence of human rabies deaths in 2021, and this for the second consecutive year, might be attributable to reduced travel due to the COVID-19 mitigation measures applied in many MSs.

Among infections caused by lyssaviruses other than RABV, five human deaths have been reported so far in Europe, more specifically in Ukraine (1977: species not characterised), Russia (1985: EBLV-1), Finland (1985: EBLV-2), the United Kingdom (2002: EBLV-2) and France (2019: EBLV-1) (Fooks et al., 2003; Regnault et al., 2022). All these infections were linked to direct exposure to infected bats; however, indirect exposure to lyssaviruses by contact with infected domestic animals (mainly free-roaming cats, occasionally infected by a bat lyssavirus) must not be overlooked. In this context, the absence of tools to prevent divergent lyssaviruses from circulating in European bats must be underlined (Echevarría et al., 2019).

The EU programmes for rabies eradication include disease surveillance, oral vaccination campaigns, the monitoring of ORV and awareness activities. Results show spread of the infection in 2021 in the

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/erms

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons

European wildlife reservoir, with a total of 118 autochthonous rabies cases reported in foxes and in domestic animals in Poland (113 cases) and Romania (5 cases), as compared with 12 cases reported in the same countries in 2020. The epidemiological data suggest stability in rabies incidence, with no more than five cases reported annually since 2017 in non-flying animals in most of the MSs reporting infections, except in Poland, which has been experiencing a rabies outbreak with domestic animal infections since 2021 in a Polish region that had been rabies-free for more than 16 years.

Data relating to rabies passive surveillance in wildlife, mainly in foxes and raccoon dogs, show relative stability in the number of samples tested, and even a slight increase in the fox sampling effort, due to the surveillance efforts maintained by two MSs for controlling the last foci. Data related to active surveillance (ORV monitoring) are reported for foxes, raccoon dogs and jackals. This surveillance is conducted by hunting apparently healthy animals and using the animals' carcases to assess the efficacy of vaccination campaigns in infected and rabies-free countries involved in eradication programmes. A sample size linked to the area covered by vaccination is recommended. The data show a substantial decrease over the years in the number of analysed foxes, reflecting the decrease in the size of areas vaccinated in the EU, hence the success of ORV programmes. The cases still being reported for several years in the few remaining MSs with infections, or a re-emergence of rabies, highlight the importance of a sustainable surveillance programme and awareness campaigns for the general public and professionals to ensure the early detection of any potential rabies cases.

Regarding rabies surveillance in bats, the number of tests has decreased since 2020. This is partly due to the withdrawal of the United Kingdom from the EU, since the UK analysed 347 bats in 2020 and 488 in 2019. The decrease observed is also related to a slightly lower number of tested bats in most reporting countries (for example, 660 and 308 samples analysed in France in 2019 and 2021, respectively and 275 and 194 bats tested in Poland in 2019 and 2021, respectively), as well as a slight decrease in the number of MSs reporting. Positive results obtained in the framework of bat surveillance (N = 29 cases) are in line with the findings of the previous years and confirm that European bats act as reservoirs for lyssaviruses other than rabies virus, reaffirming the public recommendation to handle bats with utmost caution, if at all. The public health hazard of bat lyssaviruses in Europe should not be underestimated.

In 2021, one case of imported rabies in a pet (a dog located in Lower Saxony in Germany) was reported. Such imported cases are recurrent in the EU (at a frequency of approximately one imported case per year (EFSA et al., 2022a)) and might pose a threat of rabies reintroduction into rabies-free areas.

This also underlines the need to improve public awareness, particularly among travellers, with regard to rabies risks and legislation involving pet movements. As rabies is still endemic in countries bordering the EU, in areas not far from the borders, several MSs are involved in collaborations with these countries for the implementation of vaccination and testing schemes in buffer zones. The Global Framework for the Progressive Control of Transboundary Animal Diseases (GF-TADs) created a new Standing Group of Experts on Rabies (SGE RAB) in 2019, and the third meeting was organised in 2021 with the goal of coordinating rabies control and improving surveillance activities, primarily in the Balkan sub-region, where a case was detected in a dog in 2020.

Maintaining appropriate surveillance is of paramount importance for all MSs, due to resurgence of the disease in Poland and the persistence of active foci in Romania. Apparent disappearance of the virus has been achieved in most EU territories, and maintaining appropriate surveillance efforts remains the most challenging issue to attaining rabies elimination in the EU.

## 4. Q fever

Summary data substantiating this chapter, as well as additional information on related projects and internet sources are published for this report on the EFSA Knowledge Junction on Zenodo here. Summary statistics on human surveillance data with downloadable files are retrievable using the ECDC Surveillance Atlas of Infectious Diseases here.

# 4.1. Key facts

• In 2021, the number of confirmed cases of human Q fever was 460 corresponding to an EU notification rate of 0.11 per 100,000 population. This is a decrease of 12.0% compared with the rate in 2020 (0.12 per 100,000 population).

18314732, 2022, 12, Downloaded from https://fs.aonlinelibrary.wiley.com/continelibrary.wiley.com

- Compared with the rate before the COVID-19 pandemic (2017–2019 annual mean), there was a decrease of 38.8% and 45.8% with or without data from the United Kingdom.
- Over the past 5 years (2017–2021), a significant decreasing trend (p < 0.05) in the number of O fever cases was observed.
- In 2021, Q fever cases occurred from April to September, in line with the spring/summer seasonal pattern. Cases were highest for the 50–55 years age group.
- In animals, cattle and small ruminants were mostly sampled during clinical investigations and passive monitoring of animals suspected to be infected with *Coxiella burnetii*. However, in the absence of harmonised reporting data in animals in the EU, the data reported to EFSA cannot be used to analyse spatial representativeness and trends over the years for Q fever at the EU level or to compare differences among reporting countries.
- In total, 17 MSs (15 in 2020) and five non-MSs (six in 2020) reported 2021 data for *C. burnetii*. The proportion of positive animals with direct tests was 5.9% in sheep (8.7% in 2020), 16.5% in goats (11.3% in 2020) and 5.2% in cattle (3.8% in 2020). The proportion of positive herds with direct tests was 4.1% in sheep (1.4% in 2020), 2.0% in goats (1.2% in 2020) and 4.8% in cattle (6.7% in 2020). The proportion of seropositive animals was 10.3% in sheep (11.4% in 2020), 24.6% in goats (25.0% in 2020) and 12.2% in cattle (9.6% in 2020). The proportion of seropositive herds was 18.9% in sheep (5.9% in 2020), 50.0% in goats (78.7% in 2020) and 15.1% in cattle (14.4% in 2020). Results from various other domestic and wild animal species were reported and only Italy reported positive results, mainly from dogs (73.2% out of 541) and water buffalos (4.7% out of 43).

# 4.2. Surveillance and monitoring of *Coxiella burnetii* in the EU

## 4.2.1. **Humans**

In 2021, 25 EU MSs provided data on Q fever in humans. No surveillance system is in place in Austria. Denmark did not report data for 2020 or 2021. Surveillance is mandatory in 24 MSs and voluntary in France. In 2020–2021, Spain did not receive data from all its regions; case numbers may therefore be incomplete. The EU case definition was used by 22 countries; three MSs (France, Germany and Italy) reported use of another case definition. All reporting countries had a comprehensive surveillance system. All countries reported case-based data except Belgium and Bulgaria, which provided aggregated data.

## **4.2.2.** Animals

Commission Implementing Regulation CIR (EU) 2018/1882<sup>39</sup> lists Q fever as a category E disease, meaning that, in accordance with CIR (EU) 2020/2002<sup>40</sup>, MSs shall report to the ADIS by 30 April of every year, covering the previous calendar year, on the detection of disease that has been confirmed on their territory in *Bison* ssp., *Bos* ssp., *Bubalus* ssp., *Ovis* ssp. and *Capra* ssp. MSs also submit data from annual surveillance and monitoring activities to EFSA in accordance with Directive 2003/99/EC. Q fever falls under Annex I, B. 4. 'Other zoonoses and zoonotic agents' as an agent to be monitored if warranted by the epidemiological situation in an MS, in compliance with Article 4.1 of the same Directive.

As there is no harmonised monitoring system in place for Q fever in animals in the EU, data can only be used for descriptive summaries and preclude analyses such as tracking or assessing EU-level temporal and spatial trends.

The main animal species tested are sheep, goats and cattle. Samples are mostly blood samples, samples from fetuses and stillborn animals, placentas, vaginal swabs from animals suspected of being infected with *C. burnetii*, as well as bulk milk samples for screening. Samples are tested either by serological methods (proving past or recent exposure to *C. burnetii*) or direct detection (showing carriage). In most MSs, reporting was based on clinical investigation and passive monitoring. A few countries (Belgium, Bulgaria, the Netherlands, Poland) implemented planned surveillance in cattle and

0

<sup>&</sup>lt;sup>39</sup> Commission Implementing Regulation (EU) 2018/1882 of 3 December 2018 on the application of certain disease prevention and control rules to categories of listed diseases and establishing a list of species and groups of species posing a considerable risk for the spread of those listed diseases.

<sup>40</sup> Commission Implementing Regulation (EU) 2020/2002 of 7 December 2020 laying down rules for the application of Regulation (EU) 2016/429 of the European Parliament and of the Council with regard to Union notification and Union reporting of listed diseases, to formats and procedures for submission and reporting of Union surveillance programmes and of eradication programmes and for application for recognition of disease-free status, and to the computerised information system.



Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

small ruminants, using mainly direct detection in herds in Poland (N = 4,940) and Belgium (N = 237), and mainly ELISA serological tests at the individual animal level. Belgium and the Netherlands carried out regular PCR tests on bulk tank milk from dairy sheep and goats. Some countries (Denmark, Finland, Slovenia and Spain) reported very low numbers of tests (< 100 units), corresponding to local surveys or selective tests. In addition, samples were taken from other domestic animals (dogs in particular for Italy in 2021) and wild animals on farms, in zoos or from natural habitats.

#### 4.3. Results

## 4.3.1. Overview of key statistics, EU, 2017–2021

Table 76 summarises EU-level statistics on Q fever in humans and in major animal species, respectively, during 2017–2021. More detailed descriptions of these statistics are provided in the below subsections.

**Table 76:** Summary of *Coxiella burnetii* statistics regarding humans and the main animal species, EU. 2017–2021

	2021	2020	2019 <sup>(a)</sup>	2018 <sup>(a)</sup>	2017 <sup>(a)</sup>	Data source
Humans						
Total number of confirmed cases	460	523	951	790	884	ECDC
Total number of confirmed cases/100,000 population (notification rates)	0.11	0.12	0.19	0.16	0.18	ECDC
Number of reporting EU MSs	25	25	27	27	27	ECDC
Infection acquired in the EU	359	347	810	629	720	ECDC
Infection acquired outside the EU	3	6	14	12	9	ECDC
Unknown travel status or unknown country of infection	98	170	127	149	155	ECDC
Animals						
Sheep <sup>(b)</sup>						
Animals						
Serology <sup>(c)</sup>						
Number of tested animals	750	911	666	2,986	867	EFSA
% positive animals	10.3	11.4	9.9	14.4	6.0	EFSA
Number of reporting MSs	8	8	9	10	7	EFSA
Direct detection <sup>(d)</sup>						
Number of tested animals	85	358	224	358	359	EFSA
% positive animals	5.9	8.7	18.3	12.8	8.9	EFSA
Number of reporting MSs	5	4	3	4	3	EFSA
Other methods/unknown						
Number of tested animals	521	1,555	2,244	802	1,166	EFSA
% positive animals	< 0.01	21.7	13.3	0.1	15.2	EFSA
Number of reporting MSs	1	2	2	3	2	EFSA
Herds						
Serology <sup>(c)</sup>						
Number of tested herds	2,238	17	25	197	3,654	EFSA
% positive herds	18.9	5.9	28.0	8.6	2.7	EFSA
Number of reporting MSs	1	2	1	3	4	EFSA
Direct detection <sup>(d)</sup>						
Number of tested herds	3,130	2,890	2,902	3,480	896	EFSA
% positive herds	4.1	1.4	1.2	0.9	9.6	EFSA
Number of reporting MSs	4	4	4	4	3	EFSA



	2021	2020	2019 <sup>(a)</sup>	2018 <sup>(a)</sup>	2017 <sup>(a)</sup>	Data source
Other methods/unknown						
Number of tested herds	0	39	53	1	342	EFSA
% positive herds	0	15.4	30.2	< 0.01	20.2	EFSA
Number of reporting MSs	0	1	1	1	1	EFSA
Goats						
Animals						
Serology <sup>(c)</sup>						
Number of tested animals	540	651	656	947	464	EFSA
% positive animals	24.6	25.0	18.0	20.6	3.7	EFSA
Number of reporting MSs	6	7	8	9	8	EFSA
Direct detection <sup>(d)</sup>						
Number of tested animals	200	248	189	217	148	EFSA
% positive animals	16.5	11.3	9.5	12.4	9.5	EFSA
Number of reporting MSs	4	3	4	5	2	EFSA
Other methods/unknown						
Number of tested animals	777	831	845	1,076	1,241	EFSA
% positive animals	0.13	0	0	0	8.0	EFSA
Number of reporting MSs	1	2	2	2	2	EFSA
Herds						
Serology <sup>(c)</sup>						
Number of tested herds	4	141	30	212	216	EFSA
% positive herds	50.0	78.7	73.3	25.5	25.9	EFSA
Number of reporting MS	1	1	1	3	3	EFSA
Direct detection(d)						
Number of tested herds	1,283	1,175	1,167	1,318	1,251	EFSA
% positive herds	2.0	1.2	2.8	3.0	2.6	EFSA
Number of reporting MSs	5	4	4	4	4	EFSA
Other methods/unknown						
Number of tested herds	0	12	207	11	0	EFSA
% positive herds	0	0.0	84.1	36.4	0	EFSA
Number of reporting MSs	0	1	1	1	0	EFSA
Cattle (bovine animals)						
Animals						
Serology <sup>(c)</sup>						
Number of tested animals	3,405	4,664	8,722	14,795	8,760	EFSA
% positive animals	12.2	9.6	10.5	9.8	12.4	EFSA
Number of reporting MSs	11	9	11	11	10	EFSA
Direct detection <sup>(d)</sup>						
Number of tested animals	458	842	739	4,703	4,963	EFSA
% positive animals	5.2	3.8	3.0	5.1	6.0	EFSA
Number of reporting MSs	458	842	739	4,703	4,963	EFSA
Other methods/unknown				,	,	
Number of tested animals	343	3,860	4,240	3,963	2,549	EFSA
% positive animals	0.29	0.08	0.66	2.8	0.43	EFSA
Number of reporting MSs	1	2	2	3	4	EFSA
Herds			_	-		
Serology <sup>(c)</sup>						
Number of tested herds	1,201	312	551	2,283	246	EFSA
% positive herds	15.1	14.4	30.7	10.2	21.1	EFSA

18314732, 2022, 12, Downloaded from https://drs.aonlinelibrary.wiey.com/doi/10.29033j.drs.2022.7666 by Istituto Zooprofilattico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiey.com/terms-and-conditions) on Wiley Online Library for rules of use; OA attacks are governed by the applicable Creative Commons



	2021	2020	2019 <sup>(a)</sup>	2018 <sup>(a)</sup>	2017 <sup>(a)</sup>	Data source
Number of reporting MSs	1	2	2	4	2	EFSA
Direct detection <sup>(d)</sup>						
Number of tested herds	4,311	3,571	3,673	1,262	1,244	EFSA
% positive herds	4.8	6.7	6.3	2.6	14.4	EFSA
Number of reporting MSs	8	5	5	5	4	EFSA
Other methods/unknown						
Number of tested herds	0	0	43	132	395	EFSA
% positive herds	0	0	95.3	3.8	3.8	EFSA
Number of reporting MSs	0	0	1	1	1	EFSA

ECDC: European Centre for Disease Prevention and Control; EFSA: European Food Safety Authority; MSs: Member States.

- (a): Data from the United Kingdom are taken into account for 2017–2019, because the United Kingdom was an EU MS, but it became a third country on 1 February 2020.
- (b): 'Sheep' includes the 'sheep and goats' category (NB, some countries report both livestock types together).
- (c): Serology includes the complement fixation test (CFT), ELISA, competitive ELISA (C-ELISA), indirect ELISA (I-ELISA), IgG ELISA and the immunofluorescence antibody test (IFAT).
- (d): Direct detection methods include fluorescent in situ hybridisation (FISH), immunohistochemistry (IHC), microbiological tests, multiplex PCR, PCR, real-time PCR, real-time PCR (CEN TC 275/WG 6).

#### **Animal categories**

Animal data of interest were classified into three major species categories (sheep, goats and cattle) and aggregated by year to obtain an annual overview of the volume of data submitted. The total number and the proportions of positive animals and herds were categorised according to the type of analytical testing method used (serological or direct detection).

In 2021, compared with the year 2020, the total number of samples from animals submitted by EU MSs decreased slightly for goats and more drastically for sheep and cattle. Concomitantly, the number of serological results from herds increased for sheep and for cattle (only reported this year by Italy), and slightly decreased for goats; however, the number of direct tests increased slightly for all three species.

### 4.3.2. Coxiella burnetii in humans

In 2021, 25 EU MSs reported a total of 460 confirmed cases of Q fever, with a notification rate of 0.11 cases per 100,000 population. This is a decrease of 12.0% compared with the rate in 2020 (0.12 per 100,000 population) and a decrease of 38.8% and 45.8% compared with the average annual notification rate from 2017 to 2019 (pre-pandemic period), with or without data from the United Kingdom. In 2021, Bulgaria had the highest notification rate with 0.45 cases per 100,000 population, followed by Hungary and Spain with 0.42 and 0.31 cases per 100,000 population, respectively (Table 77).

In 2021, 359 cases (78%) were acquired in the EU. Germany and Sweden reported travel-associated cases. Three cases were imported from Ethiopia, Iran and Sudan (0.7%), whereas for 98 cases (21.3%), there were no data on travel or the country of infection (Table 76).

In 2021, 11 MSs reported outcomes for 270 cases (58.7%) with four deaths (in Germany, Hungary, Spain and Portugal) recorded in patients over 50 years old, resulting in an EU case fatality of 1.5%.

18314732, 2022, 12, Downdoaded from https://sta.onlinelibrary.wiley.com/doi/10.2003.j.d/sa.2022.7666 by Istituto Cooperal lative Sperimental dell'Umbria e delle Marche (ISSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/emens-and-conditions) on Wiley Online Library for rules of use; Os Aurited sea geoverned by the applicable Creative Common



**Table 77:** Reported confirmed human cases of Q fever and notification rates per 100,000 population in EU MSs and non-MS countries, by country and year, 2017–2021

Country co	National overage <sup>(a)</sup>	2021 Data	Confi		20	20	20:	19	20:	18	201	L7	
cc	National overage <sup>(a)</sup>	Data	Confi										
A otuio (b)		format <sup>(a)</sup>	cases and rates		Confi cases rat	and	Confi cases rat	and	Confi cases rat	and	Confi cases rat	and	
A otuio(b)			Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate	
Austria	-	_	-	_	_	_	_	_	-	_	_	-	
Belgium	Υ	Α	6	0.05	4	0.03	10	0.09	6	0.05	7	0.06	
Bulgaria	Y	Α	31	0.45	103	1.48	36	0.51	45	0.64	28	0.39	
Croatia	Υ	С	0	0	2	0.05	8	0.20	11	0.27	23	0.55	
Cyprus	Υ	С	2	0.22	1	0.11	1	0.11	0	0	3	0.35	
Czechia	Υ	С	1	0.01	1	0.01	1	0.01	1	0.01	0	0	
Denmark	Υ	С	_	_	_	_	0	0	0	0	0	0	
Estonia	Υ	С	0	0	0	0	0	0	0	0	0	0	
Finland	Y	С	0	0	0	0	2	0.04	2	0.04	4	0.07	
France	Υ	С	92	0.14	96	0.14	156	0.23	172	0.26	194	0.29	
Germany	Υ	С	99	0.12	55	0.07	148	0.18	91	0.11	107	0.13	
Greece	Υ	С	4	0.04	4	0.04	14	0.13	13	0.12	4	0.04	
Hungary	Υ	С	41	0.42	34	0.35	47	0.48	28	0.29	29	0.30	
Ireland	Υ	С	0	0	2	0.04	2	0.04	0	0	2	0.04	
Italy	Υ	С	0	0	0	0	6	0.01	1	< 0.01	7	0.01	
Latvia	Υ	С	0	0	1	0.05	0	0	0	0	0	0	
Lithuania	Υ	С	0	0	0	0	0	0	0	0	0	0	
Luxembourg	Υ	С	0	0	2	0.32	0	0	0	0	0	0	
Malta	Υ	С	1	0.19	0	0	1	0.20	2	0.42	0	0	
Netherlands	Y	С	6	0.03	7	0.04	16	0.09	18	0.10	22	0.13	
Poland	Υ	С	0	0	0	0	4	0.01	0	0	0	0	
Portugal	Υ	С	18	0.17	22	0.21	32	0.31	36	0.35	48	0.47	
Romania	Υ	С	5	0.03	12	0.06	109	0.56	22	0.11	46	0.23	
Slovakia	Υ	С	2	0.04	5	0.09	1	0.02	2	0.04	0	0	
Slovenia	Υ	С	0	0	1	0.05	6	0.29	1	0.05	3	0.15	
Spain	Υ	С	149	0.31	170	0.36	332	0.71	313	0.67	333	0.72	
Sweden	Υ	С	3	0.03	1	0.01	10	0.10	7	0.07	3	0.03	
EU Total 27			460	0.11	523	0.12	942	0.22	771	0.18	863	0.20	
United Kingdom	-	_	-		-		9	0.01	19	0.03	21	0.03	
EU Total <sup>(c)</sup>			460	0.11	523	0.12	951	0.19	790	0.16	884	0.18	
Iceland	Υ	С	0	0	0	0	0	0	0	0	0	0	
Norway	Υ	С	3	0.06	5	0.09	8	0.15	5	0.09	4	0.08	
Liechtenstein	Υ	С	0	0	55	0.64	103	1.20	52	0.61	41	0.48	
Switzerland <sup>(d)</sup>	Υ	С	111	1.3									

<sup>-:</sup> Data not reported.

1831/4732, 2022, 12, Downloaded from https://eftsaonlinelibrary.wiley.com/doi/10.2903j.eftsa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/203]. See the Terms and Conditions (https://onlinelibrary.wiley.com/erms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

<sup>(</sup>a): Y: yes; N: no; A: aggregated data; C: case-based data.

<sup>(</sup>b): Not notifiable, no surveillance system exists.


<sup>(</sup>c): Cases reported by the United Kingdom for the period 2017–2019 were also considered in this estimation (EU-28). When the United Kingdom data were collected for the period 2017–2019, the United Kingdom was an EU MS, but it became a third country on 1 February 2020.

<sup>(</sup>d): Switzerland provided data directly to EFSA. The human data for Switzerland include data from Liechtenstein for years 2017–2020.



In 2021, cases occurred year-round. Most Q fever cases occurred from April to September, in line with the spring/summer seasonal pattern (Figure 39).

In 2021, the number of Q fever cases decreased compared with 2020 and the 2017–2019 period. In particular, a significant (p < 0.01) decreasing trend was noted for France, the Netherlands and Portugal. However, over the past 5 years, a significant decrease (p < 0.05) in Q fever cases has been observed in the EU. The rate of confirmed human Q fever cases was higher among men than women with a male-to-female ratio of 2.1:1. The highest notification rate occurred in the 50-55 years age group (12.5%).



Source: Data from Cyprus, Czechia, Estonia, Finland, France, Germany, Greece, Hungary, Ireland, Latvia, Lithuania, Malta, the Netherlands, Poland, Portugal, Romania, Slovakia, Slovenia and Sweden. Austria, Belgium, Bulgaria, Croatia, Denmark, Italy, Luxembourg and Spain did not report data at the level of detail required for the analysis.

Figure 39: Trend in reported confirmed human cases of Q fever in the EU by month, 2017–2021

## 4.3.3. Coxiella burnetii in animals

#### Sheep (including the 'sheep and goats' category)

Seven MSs and three non-MSs (North Macedonia, Serbia and Switzerland) provided data in 2021. In total, 5,368 herds and 1,356 animals were tested. At the animal level, serological tests were more often used (N = 750, 10.3% positive) than direct detection tests (N = 85, 5.9% positive). Most individual serological tests were reported by Greece (N = 288, 17.7% positive), Bulgaria (N = 185, 8.1% positive), Ireland (N = 123, 0.8% positive) and Sweden (N = 98, 0.0% positive). At the herd level, Italy accounted for all the reported serological tests (N = 2,238, 18.9% positive), and most direct tests were reported by Poland (N = 2,452, 0.1% positive), Italy (N = 508, 20.9% positive) and Belgium (N = 162, 8.6% positive).

## Goats

Eleven MSs and one non-MS (Switzerland) provided data in 2021. In total, 1,287 herds and 1,517 animals were tested. At the animal level, serological tests were more often used (N = 540, 24.6% positive) than direct detection tests (N = 200, 16.5% positive). Most serological tests were reported by

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/continelibrary.wiley.c

Italy (N = 215, 27.9% positive) and Bulgaria (N = 158, 36.7% positive). Italy and Switzerland reported the most direct detection tests (N = 162, 17.9% positive; N = 159, 11.3% positive, respectively). At the herd level, almost all tests were direct detection tests, mostly carried out in Poland (N = 926, 0.0% positive), Belgium (N = 241, 5.4% positive) and Spain (N = 81, 12.3% positive).

#### Cattle

Sixteen MSs and five non-MSs (Iceland, Norway, North Macedonia, Serbia and Switzerland) provided data in 2021. In total, 5,512 herds and 4,206 animals were tested. At the animal level, serological tests were more often employed (N = 3,405, 12.2% positive) than direct detection tests (N = 458, 5.2%). Slovakia (N = 1,137, 3.2% positive), Austria (N = 965, 13.7% positive) and Hungary (N = 451, 37.7% positive) provided 75% of the serological tests, whereas Switzerland tested 3,277 animals using direct detection tests, resulting in 1.7% positivity. For herds, most tests were direct detection methods and mainly reported from Belgium (N = 2,199, 7.6% positive) and Poland (N = 1,864, 1.0% positive), whereas Italy reported data mostly obtained using serological methods (N = 1,201, 15.1% positive).

#### Other animal species

Four MSs (Austria, Cyprus, Italy and Sweden) and two non-MSs (Norway and Switzerland) reported data in 2021 on animals other than sheep, goats and cattle. In total, 38 herds and 706 animals were tested from various domestic and wild animal species, such as other ruminant species (water buffalo, llama, etc.), pets (cats and dogs) or diverse other species. Positive results were only reported by Italy, mainly from dogs (N = 541, 73.2% positive) and water buffalos (N = 43, 4.7% positive).

#### 4.4. Discussion

*C. burnetii* is the aetiological agent of the Q fever disease and, if aerosolised, is also considered a potential biological weapon. Humans can acquire the infection mainly through environmental contamination due to bacterial shedding of infected animals, but also through tick-borne or foodborne transmissions. In Europe, the majority of clinical cases are sporadic. However, several outbreaks among humans have been reported. Up to 2016, France and Germany reported most of the confirmed cases. In 2017, two outbreaks were reported from Bulgaria in the Gabrovo and Blagoevgrad regions (Genova-Kalou et al., 2019). Since 2017, Spain has reported the highest number of cases annually. The increase in the number of human cases reported by Spain is most likely explained by a change in their reporting system from voluntary to mandatory.

In 2021, the number of human Q fever cases in the EU was the lowest recorded over the past 5 years. Cases decreased compared with the COVID-19 pandemic year 2020, but also compared with the pre-pandemic 2017–2019 period. Compared with 2020, a decrease was observed in Bulgaria and Spain. However, Spain accounted for about one third of the overall number of cases, whereas Germany reported an increase in Q fever disease in humans. Several Q fever cases recorded in Germany and Spain were linked to occupational (waste treatment plant or laboratory staff) or recreational activities, and the exposure was from animal and environmental contamination (ProMEDmail, 2021a,b,c). Overall, a significant decreasing trend was observed over the last 5 years (2017–2021). Year 2021 showed a reduction in case fatality (1.48%) compared with 2020 (2.13%). Moreover, no data were available for 2021 regarding the impact of the pandemic on Q fever surveillance and reporting.

The results obtained from animals – mainly from small ruminants and cattle – are insufficient for analysing trends for Q fever at the EU level. The start of annual Union reporting in 2021 (Q fever is a 'Category E' disease under the new EU Animal Health Law) has not drastically increased the total number of countries reporting to EFSA under Directive 2003/99/EC. The results submitted by different MSs and non-MSs are not directly comparable, mainly due to differences in sampling strategy (sample types, testing methods, coverage of the monitoring), data completeness and sensitivity of the surveillance method. Importantly, EU MSs should first agree on case definitions in domestic ruminants at the animal and herd levels and also improve data reporting harmonisation among countries. Case definitions could be based on EFSA harmonisation guidelines (Sidi-Boumedine et al., 2010) and the WOAH case definition draft (WOAH, 2022). Furthermore, monitoring could be extended to include species other than domestic ruminants, such as pets, which may be reservoirs. In 2021, Italy reported a high number (394) and proportion of positive dogs (76.8%), compared with published 2020 data indicating 8.2% positivity in Italian dogs (Ebani et al., 2020). Pets have already been suspected to be

18314732, 2022, 12, Downloaded from https://fs.aonlinelibrary.wiley.com/continelibrary.wiley.com

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/erms

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons

reservoirs for *C. burnetii* in Australia (Tozer et al., 2014; Orr et al., 2022), Asia (Lyoo et al., 2017) and Africa (Kelly et al., 1993; Boni et al., 1998). They may also be considered as sentinels for humans and farm animals. In coming years, it is of utmost importance to collect more data on the persistence of environmental contamination (animal waste, dust) to better assess risk factors (Carrié et al., 2019). The major challenge is to reduce human exposure to this zoonosis through a preventive 'One Health' approach.

#### 5. West Nile virus

Summary data substantiating this chapter, as well as additional information on related projects and internet sources are published for this report on the EFSA Knowledge Junction on Zenodo here. Summary statistics on human surveillance data with downloadable files are retrievable using the ECDC Surveillance Atlas of Infectious Diseases here.

# **5.1.** Key facts

- In 2021, the number of locally acquired probable and confirmed human cases of West Nile virus (WNV) infection was 152, corresponding to an EU notification rate of 0.03 per 100,000 population. This is a decrease of 57% compared with 2020 (0.07 per 100,000 population) and the lowest notification rate reported in the period 2017–2021.
- Most locally acquired infections were reported by Italy and Greece, accounting respectively for 43% and 39% of the total number of probable/confirmed human cases in the EU.
- Since the epidemic year of 2018, which was characterised by an unusually intense transmission season, the trend in the number of reported human cases of infection seems to be slowly decreasing. However, no significant decrease (p = 0.89) has been observed over the past 5 years (2017–2021). Nevertheless, over this same period, 2021 was the year in which countries reported the lowest number of cases in humans and the lowest number of outbreaks in animals.
- In Spain, a country that experienced intense transmission in 2020, a significant decrease in the number of human cases was documented over the same period.
- In 2021, 15 EU MSs and Switzerland submitted WNV monitoring and surveillance data from birds and/or equids to EFSA. Italy and Spain submitted respectively 62.3% and 13.7% of these data on birds, while Spain, Greece and Germany submitted most of the data for equids, at 42.8%, 17.6% and 15.1% respectively.
- Italy, Germany, Slovenia and Spain reported infections in birds (only 2 MSs reported outbreaks in ADIS) with Italy and Germany reporting the highest number of avian cases (73.3% and 23.3% respectively). Germany, Spain, Italy, Portugal, France, Hungary and Greece reported equid outbreaks to ADIS, with the highest number of cases being reported by Germany, Spain and Italy, accounting for 40.4%, 23.4% and 14.9% of the total number of cases respectively.
- The data reported to ECDC, ADIS and EFSA for 2021 indicated WNV circulation in central and eastern Europe (Germany, Hungary, Austria, Romania and Slovenia) as well as in southern and western Europe (Greece, Italy, France, Spain and Portugal). WNV infections of humans, equids and birds now regularly occur in these countries.

# 5.2. WNV ecology and epidemiology

West Nile fever, also known as West Nile virus disease, is an arboviral disease caused by an arbovirus called the West Nile virus (WNV). WNV genetic lineages one and two are associated with both human and animal diseases.

Mosquitoes serve as vectors and birds are the main amplifying hosts. This virus is transmitted to humans and equids through the bite of infected mosquitoes, mainly of the *Culex* genus (Hubálek and Halouzka, 1999). Occasionally, infection can occur through the transfusion/transplantation of substances of human origin (SoHO) (i.e. blood, organs or cells), percutaneous or conjunctival exposure in laboratories or transplacental passage from mother to fetus. The number of EU countries reporting locally acquired WNV infections in humans and cases of infection in birds/equids has increased in recent years.

WNV circulation is influenced by environmental conditions, and is helped by low precipitation in winter and high spring temperatures (Marini et al., 2021; Farooq et al., 2022). These conditions are

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/erms

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

beneficial to the *Culex* mosquito population, and enhance virus replication in the vector (Reisen et al., 2006; Fornasiero et al., 2020).

## 5.3. Surveillance and monitoring of West Nile virus in the EU

#### **5.3.1.** Humans

In 2021, 25 MSs reported information on WNV infections in humans. Surveillance is mandatory in 23 countries, in France it is voluntary, and Germany did not specify their surveillance system. There is no surveillance system in place in Denmark and the disease is not notifiable or reported at EU level. The EU case definition was used by 23 MSs, France reported another case definition and Germany did not specify which case definition was used. All countries conducting surveillance had a comprehensive surveillance system with full national coverage, except Germany, which did not specify the surveillance system but had full national coverage. All countries reported case-based data.

#### **5.3.2.** Animals

From a veterinary standpoint, WNV is the causative agent of West Nile Fever (WNF), a disease that develops into asymptomatic forms, benign forms (flu-like syndrome) and neuro-invasive forms (WOAH, 2018a). WNV surveillance in animals involves mostly passive surveillance, including surveillance based on the diagnosis of neuro-invasive cases in equids, but some countries implement active surveillance of equids and/or captive birds and/or wild birds. Alongside EU MSs, Switzerland submits reports to EFSA on animal surveillance and monitoring activities in animals. Two sources of information are used to complete this report. Firstly, data are submitted to EFSA by EU MSs and Switzerland from annual surveillance and monitoring activities in accordance with Directive 2003/99/ EC. WNV is listed in Annex I, Part B (viruses transmitted by arthropods) as a virus to be monitored, if warranted by the epidemiological situation in an MS, in compliance with Article 4.1 of the same Directive. Secondly, it is mandatory for MSs to notify outbreaks of equine and avian WNV to ADIS, <sup>41</sup> in accordance with CIR (EU) 2020/2002.

#### 5.4. Results

# 5.4.1. Overview of key statistics, EU, 2017–2021

In 2021, WNV cases were reported in humans, equids and birds. Over the past 5 years, there has been an increase in the number of countries reporting surveillance data in animals (Table 78). While the number of tested equids appears to correlate with WNV circulation, increasing in years when a greater number of cases are reported and decreasing when WNV circulation is observed to fall, there has been an overall increase in the number of tested birds over time, reflecting an intensification of the surveillance system in place.

The number of probable and confirmed human cases of West Nile virus (WNV) infection reported in 2021 was 158, corresponding to an EU notification rate of 0.04 per 100,000 population. This is a decrease of 50% compared with 2020 (0.08 per 100,000 population) and the lowest notification rate reported in the period 2017–2021.

More detailed descriptions of these statistics are provided in the below subsections.

\_

<sup>&</sup>lt;sup>41</sup> Animal Disease Information System.



Table 78: Summary of WNV infection statistics related to humans, birds and equids EU, 2017–2021

	2021 <sup>(a)</sup>	2020	2019 <sup>(b)</sup>	2018 <sup>(b)</sup>	2017 <sup>(b)</sup>	Data source
Humans						
Total number of confirmed and probable cases	158	333	443	1,615	208	ECDC
Total number of confirmed and probable cases/100,000 population (notification rates)	0.04	0.08	0.09	0.32	0.05	ECDC
Number of reporting MSs	26	26	27	27	26	ECDC
Infection acquired in the EU	153 <sup>(c)</sup>	331	435	1,573	205	ECDC
Infection acquired outside the EU	5	2	5	29	2	ECDC
Unknown travel status or unknown country of infection	0	0	3	13	1	ECDC
Animals <sup>(d)</sup>						
Birds						
Number of animals tested	19,596	11,141	14,932	13,970	11,173	EFSA
Number of positive animals by PCR <sup>(e)</sup> -based methods	146	165	104	425	93	EFSA
Number of MSs reporting surveillance/monitoring data to EFSA	13	11	13	11	7	EFSA
Number of outbreaks notified to ADIS	8	2	53	22	0	ADIS
Number of MSs notifying outbreaks to ADIS	2	1	2	6	0	ADIS
Equids						
Number of animals tested	5,985	6,749	5,563	13,785	11,668	EFSA
Number of positive animals by PCR <sup>(e)</sup> -based methods	1	1	4	7	1	EFSA
Number of animals positive for IgM by ELISA	47	209	74	393	110	EFSA
Number of MSs reporting surveillance/monitoring data to EFSA	14	14	14	12	12	EFSA
Number of outbreaks notified to ADIS	45	189	100	292	84	ADIS
Number of MSs notifying outbreaks to ADIS	7	9	8	10	7	ADIS

ADIS: Animal Disease Information System; ECDC: European Centre for Disease Prevention and Control; EFSA: European Food Safety Authority; ELISA: enzyme-linked immunosorbent assay; MSs: Member States; PCR: polymerase chain reaction.

Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.c

<sup>(</sup>a): Data on animals from the United Kingdom (Northern Ireland) are taken into account for 2021. In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/Northern Ireland, EU requirements on data sampling are also applicable to Northern Ireland.


<sup>(</sup>b): Data from the United Kingdom are taken into account for 2017–2019, because the United Kingdom was an EU MS, but it became a third country on 1 February 2020.

<sup>(</sup>c): One case reported from the EU was later found to have been infected in Mayotte.

<sup>(</sup>d): Samples tested using an unspecified analytical method are not included.

<sup>(</sup>e): PCR: polymerase chain reaction.





**Figure 40:** Geographical distribution of locally acquired West Nile virus infections among humans (NUTS 3 level) and outbreaks notified to ADIS among equids and birds (XY coordinates) across the EU, 2021 transmission season

### 5.4.2. West Nile virus infections in humans

In 2021, 158 confirmed and probable WNV infections in humans were reported. Of these, 153 were acquired in the EU. Of the 153 cases acquired in the EU, 152 were locally acquired (Table 79) and one was imported from another EU country. Of the 152 cases of probable/confirmed infection that were locally acquired in 2021, 93 concerned males, (61.2%), and over 80% of cases occurred in people aged 50 or older. Five cases of travel-associated WNV infection acquired outside the EU (Dominican Republic, Turkey and the United States) were also reported.



18314732, 2022, 12, Downloaded from https://efs.aonlinelibrary.wiley.com/doi/10.2903/jefs.a.2022.7666 by Istituto Zopprofilative Sperimenta dell'Umbria e delle Marche (ISSUM), Wiley Online library.wiley.com/rems-and-conditions) on Wiley Online library for rules of use. O. A articles are governed by the applicable Creative Common

**Table 79:** Locally acquired human WNV cases (confirmed and probable) and notification rates per 100,000 population in EU MSs and non-MS countries, by country and year, 2017–2021

			2021			20	20	20	19	20	18	20	17
Country	National	Data	Confirmed		cases* rates	Total o			ses* and tes		cases* rates		cases* rates
	coverage <sup>(a)</sup>	format <sup>(u)</sup>	cases	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate
Austria	Y	С	3	3	0.03	0	0	4	0.05	21	0.24	6	0.07
Belgium	Υ	С	0	0	0	0	0	0	0	0	0	0	0
Bulgaria	Υ	С	0	0	0	1	0.01	5	0.07	15	0.21	1	0.01
Croatia	Υ	С	0	0	0	0	0	0	0	58	1.4	5	0.12
Cyprus	Υ	С	0	0	0	0	0	23	2.6	1	0.12	0	0
Czechia	Υ	С	0	0	0	0	0	1	0.01	5	0.05	0	0
Denmark <sup>(b)</sup>	-	-	-	-	-	-	-	-	-	-	-	-	-
Estonia	Υ	С	0	0	0	0	0	0	0	0	0	0	0
Finland	Υ	С	0	0	0	0	0	0	0	0	0	0	0
France	Υ	С	1 <sup>(c)</sup>	1 <sup>(c)</sup>	< 0.01	0	0	2	< 0.01	27	0.04	2	< 0.01
Germany	Υ	С	4	4	< 0.01	22	0.03	5	0.01	1	< 0.01	-	_
Greece <sup>(d)</sup>	Υ	С	27	59	0.55	144	1.3	227	2.1	315	2.9	48	0.45
Hungary	Υ	С	6	7	0.07	3	0.03	36	0.37	215	2.2	20	0.20
Ireland	Υ	С	0	0	0	0	0	0	0	0	0	0	0
Italy	Υ	С	65	65	0.11	69	0.12	54	0.09	610	1.0	53	0.09
Latvia	Υ	С	0	0	0	0	0	0	0	0	0	0	0
Lithuania	Υ	С	0	0	0	0	0	0	0	0	0	0	0
Luxembourg	Υ	С	0	0	0	0	0	0	0	0	0	0	0
Malta	Υ	С	0	0	0	0	0	0	0	0	0	0	0
Netherlands	Υ	С	0	0	0	8	0.05	0	0	0	0	0	0
Poland	Υ	С	0	0	0	0	0	0	0	0	0	0	0
Portugal	Υ	С	0	0	0	0	0	0	0	0	0	0	0
Romania	Υ	С	6	7	0.04	6	0.03	67	0.35	277	1.4	66	0.34
Slovakia	Υ	С	0	0	0	0	0	1	0.02	0	0	0	0
Slovenia	Υ	С	0	0	0	0	0	0	0	4	0.19	0	0
Spain	Υ	С	6	6	0.01	77	0.16	0	0	0	0	0	0
Sweden	Υ	С	0	0	0	0	0	0	0	0	0	0	0
EU Total 27	-	_	118	152	0.03	330	0.07	425	0.10	1.55	0.35	201	0.06
United Kingdom	_	_	-	_	_	-	_	0	0	0	0	0	0
EU Total <sup>(e)</sup>	-	_	118	152	0.03	330	0.07	425	0.08	1.55	0.31	201	0.05
Iceland	Υ	С	0	0	0	0	0	0	0	0	0	0	0
Norway	Υ	С	0	0	0	0	0	0	0	0	0	0	0
Liechtenstein	Υ	С	0	0	0	1	0.01	1	0.01	0	0	0	0
Switzerland <sup>(f)</sup>	Υ	С	0	0	0								

<sup>-</sup>: Data not reported.

- (a): Y: yes; N: no; A: aggregated data; C: case-based data.
- (b): No surveillance system.
- (c): This case reported by France acquired the WNV infection in Mayotte.
- (d): Only locally acquired cases diagnosed in the country of exposure are included in the table.
- (e): Cases reported by the United Kingdom for the years 2017–2019 were also considered for this estimation (EU-28). When United Kingdom data were collected for the period 2017–2019, the United Kingdom was an EU MS, but it became a third country on 1 February 2020.
- (f): Switzerland provided data directly to EFSA. The human data for Switzerland include data from Liechtenstein for the years 2017–2020.

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Isituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (thtps://onlinelibrary.wiley.com/terms

**Table 80:** Locally acquired human WNND cases (confirmed and probable) and notification rates per 100,000 population in the EU MSs and non-MS countries, by country and year, 2017–2021

		2	021			20	020	2	019	20	18	20	17
Country	National coverage <sup>(a)</sup>	Data format <sup>(a)</sup>	Confirmed cases		cases* rates								
	Coverage	Tormac	Cases	Cases	Rate								
Austria	Υ	С	1	1	0.01	0	0	1	0.01	4	0.05	2	0.02
Belgium	Υ	С	0	0	0	0	0	0	0	0	0	0	0
Bulgaria	Υ	С	0	0	0	1	0.01	4	0.06	13	0.18	1	0.01
Croatia	Υ	С	0	0	0	0	0	0	0	47	1.1	5	0.12
Cyprus	Υ	С	0	0	0	0	0	20	2.3	1	0.12	0	0
Czechia	Υ	С	0	0	0	0	0	1	0.01	3	0.03	0	0
Denmark <sup>(b)</sup>	_	_	-	_	_	_	_	_	_	-	_	-	_
Estonia	Υ	С	0	0	0	0	0	0	0	0	0	0	0
Finland	Υ	С	0	0	0	0	0	0	0	0	0	0	0
France	Υ	С	0	0	0	0	0	1	< 0.01	7	0.01	0	0
Germany	Υ	С	2	2	< 0.01	7	0.01	3	< 0.01	0	0	-	_
Greece <sup>(c)</sup>	Υ	С	22	38	0.36	116	1.1	140	1.3	241	2.2	28	0.26
Hungary	Υ	С	6	7	0.07	1	0.01	23	0.24	152	1.6	17	0.17
Ireland	Υ	С	0	0	0	0	0	0	0	0	0	0	0
Italy	Υ	С	35	35	0.06	46	0.08	24	0.04	243	0.40	26	0.04
Latvia	Υ	С	0	0	0	0	0	0	0	0	0	0	0
Lithuania	Υ	С	0	0	0	0	0	0	0	0	0	0	0
Luxembourg	Υ	С	0	0	0	0	0	0	0	0	0	0	0
Malta	Υ	С	0	0	0	0	0	0	0	0	0	0	0
Netherlands	Υ	С	0	0	0	6	0.03	0	0	0	0	0	0
Poland	Υ	С	0	0	0	0	0	0	0	0	0	0	0
Portugal	Υ	С	0	0	0	0	0	0	0	0	0	0	0
Romania	Υ	С	6	7	0.04	6	0.03	65	0.33	277	1.4	66	0.34
Slovakia	Υ	С	0	0	0	0	0	0	0	0	0	0	0
Slovenia	Υ	С	0	0	0	0	0	0	0	4	0.19	0	0
Spain	Y	С	6	6	0.01	72	0.15	0	0	0	0	0	0
Sweden	Υ	С	0	0	0	0	0	0	0	0	0	0	0
EU Total 27	_	_	78	96	0.02	255	0.06	282	0.06	992	0.23	145	0.04
United Kingdom	_	-	_	_	_	_	_	0	0	0	0	0	0
EU Total <sup>(d)</sup>	_	-	78	96	0.02	255	0.06	282	0.06	992	0.20	145	0.03
Iceland	Υ	С	0	0	0	0	0	0	0	0	0	0	0
Norway	Υ	С	0	0	0	0	0	0	0	0	0	0	0
Liechtenstein	Υ	С	0	0	0	_	_	-	_	-	_	-	_
Switzerland <sup>(e)</sup>	Υ	С	_	_	_								

WNND: West Nile Neuro-invasive Disease.

Six EU MSs reporting locally acquired infections in humans provided data on the hospitalisation status of their cases. Of the cases with known hospitalisation status (83 cases, 55% of total infections) in 2021, 84% (N = 70) were hospitalised.

235

<sup>-:</sup> Data not reported.

<sup>(</sup>a): Y: yes; N: no; A: aggregated data; C: case-based data.

<sup>(</sup>b): No surveillance system.

<sup>(</sup>c): Only locally acquired cases diagnosed in the country of exposure are included in the table.

<sup>(</sup>d): Cases reported by the United Kingdom for the period 2017–2020 were also considered for this estimation (EU-28). When the United Kingdom data were collected for the period 2017–2020, the United Kingdom was an EU MS, but it became a third country on 1 February 2020.

<sup>(</sup>e): Switzerland provided data directly to EFSA. The human data for Switzerland include data from Liechtenstein for the years 2017–2020.



18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903j.gfsa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [1812/2021, See the Terms and Conditions (https://onlinelibrary.wiley.com/terms/

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

Clinical manifestations were reported for all of the 152 locally acquired cases. Among those, 63% (N = 96) of infections were neuro-invasive (Table 80) and 16% (N = 25) were asymptomatic. This compares with 79% (N = 252) neuro-invasive and 6% (N = 19) asymptomatic infections in 2020. In 2021, the remaining 31 cases (20%) were symptomatic cases with non-neurological symptoms. This compares with 46 cases (15%) in 2020.

Data on the outcome of infections were reported for all of the 152 locally acquired cases. In 2021, 11 deaths among cases with WNV infections were reported, compared with 39 in 2020. The case fatality rate in 2021 was 7% (12% in 2020) among all locally acquired WNV infections and 11% (15% in 2020) among locally acquired WNV infections with West Nile neuro-invasive disease (WNND).

Eight EU MSs (Austria, France, Germany, Greece, Hungary, Italy, Romania and Spain) reported at least one locally acquired human case of WNV infection in 2021. Most of the locally acquired infections were reported by Italy and Greece, accounting respectively for 43% and 39% of the total number of probable/confirmed cases in the EU.

## 5.4.3. West Nile virus infections in animals

In 2021, 15 EU MSs and Switzerland submitted WNV monitoring and surveillance data from birds and/or equids to EFSA. None of the analyses undertaken on other species (sheep and goats at farm, wild martens and one crocodile in a zoo) were positive. Summaries by country of reported cases, and of outbreaks, in birds and equids are provided in Table 81, which displays data from both data sources. In some cases, the statistics from both sources may show discrepancies, so the following points should be taken into consideration for interpreting the data:

- The reported number of animals tested based on surveillance activities submitted to EFSA includes the following registered methods: detection of IgG and IgM antibodies, seroneutralisation and the detection of flavivirus or WNV genomes using a PCR-based method. Samples tested using an unspecified analytical method are not included.
- The notification of outbreaks in equids and birds to ADIS is mandatory. However, some countries did not report confirmed outbreaks among birds to ADIS, while nevertheless including them in their national reports. An outbreak can refer to more than one infected animal if they constitute a single epidemiological unit and/or are from the same location.
- The number of positive animals includes cases for which the disease was confirmed clinically and/or in a laboratory setting either by the detection of IgM-specific antibodies indicating a recent infection by WNV (available for equids only) or by the detection of the WNV genome using a PCR-based method indicating an ongoing infection (for both birds and equids). The results of serum neutralisation testing and IgG ELISA, including observation of seroconversion in avian or equid sentinel animals by detecting specific neutralising antibodies, are not included, because it is not possible to determine when the infection occurred.
- Some countries did not submit data to either ADIS or EFSA.

The geographical distribution of outbreaks in animal is shown in Figure 40.

**Table 81:** Summary of WNV surveillance/monitoring results in animals reported to EFSA, and WNV outbreaks notified to ADIS, by EU MSs and non-MS countries, 2021

		Birds <sup>(a)</sup>			Equ	ıids <sup>(a)</sup>	
Country (EU	activities s	surveillance submitted to FSA		Data on s	N (0/ )		
MSs, non-EU country)	N (%) animals tested	N (%) animals positive using PCR- based methods <sup>(b)</sup>	N (%) outbreaks notified in ADIS	N (%) animals tested	N (%) animals positive using ELISA- IgM <sup>(c)</sup>	N (%) animals positive using PCR- based methods <sup>(b)</sup>	N (%) outbreaks notified in ADIS
Austria	170 (0.87)	0 (0)	NR	16 (0.27)	_	0 (0)	NR
Cyprus	498 (2.5)	_	NR	103 (1.7)	0 (0)	_	NR
Czechia	NR	_	NR	783 (13.1)	_	_	NR



18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilatico

Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [1812/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Common

		Birds <sup>(a)</sup>		Equids <sup>(a)</sup>							
Country (Ell	activities s	surveillance submitted to FSA			Data on surveillance activities submitted to EFSA						
Finland France Germany Greece Hungary Italy Portugal Romania Slovenia Spain Sweden EU Total (27 + XI)	N (%) animals tested	N (%) animals positive using PCR- based methods <sup>(b)</sup>	N (%) outbreaks notified in ADIS	N (%) animals tested	N (%) animals positive using ELISA- IgM <sup>(c)</sup>	N (%) animals positive using PCR- based methods <sup>(b)</sup>	N (%) outbreaks notified in ADIS				
Denmark	891 (4.5)	-	NR	NR	-	_	NR				
Finland	25 (0.13)	0 (0)	NR	144 (2.4)	0 (0)	_	NR				
France	85 (0.43)	0 (0)	NR	22 (0.37)	3 (6.4)	0 (0)	2 (4.4)				
Germany	1,909 (9.7)	34 (23.3)	NR	906 (15.1)	19 (40.4)	_	18 (40.0)				
Greece	38 (0.19)	0 (0)	NR	1,052 (17.6)	0 (0)	1 (100)	1 (2.2)				
Hungary	794 (4.1)	0 (0)	NR	109 (1.8)	3 (6.4)	0 (0)	3 (6.7)				
Italy	12,204 (62.3)	107 (73.3)	NR	195 (3.3)	7 (14.9)	_	6 (13.3)				
Portugal	NR	_	NR	6 (0.10)	4 (8.5)	_	4 (8.9)				
Romania	6 (0.03)	_	NR	84 (1.4)	0 (0)	_	NR				
Slovenia	38 (0.19)	1 (0.68)	1 (12.5)	2 (0.03)	_	0 (0)	NR				
Spain	2,684 (13.7)	4 (2.7)	7 (87.5)	2,562 (42.8)	11 (23.4)	_	11 (24.4)				
Sweden	254 (1.3)	0 (0)	NR	1 (0.02)	_	0 (0)	NR				
EU Total (27 + XI)	19,596 (100)	146 (100)	8 (100)	5,985 (100)	47 (100)	1 (100)	45 (100)				
Switzerland	0 (0)	_	NR	9 (0)	_	0 (0)	NR				

MSs: Member States; ADIS: Animal Disease Information System.

NR: Not reported to EFSA or to ADIS. These countries did not submit data for WNF surveillance activities to EFSA or did not notify outbreaks to ADIS.

- -: Analytical method not used.
- 0: Analytical method used with negative results.
- (a): Samples tested with an unspecified analytical method are not included.
- (b): PCR: polymerase chain reaction (for identification of the virus genome).
- (c): ELISA: enzyme-linked immunosorbent assay.

In birds, 13 MSs reported a total of 19,596 analysed samples as part of their surveillance and monitoring activities (Tables 78 and 81). Italy, Spain and Germany submitted 62.3%, 13.7% and 9.7% of these data respectively (Table 81). Of the total number, 16,247 samples from Hungary, Greece, France, Slovenia, Italy, Finland, Spain, Germany, Sweden, Austria were tested using a WNV-specific PCR method. A total of 146 infected birds were reported to EFSA by Italy, Germany, Slovenia and Spain (Table 81). WNV infection occurred in crows, doves, goshawks, jays, magpies, owls, Passeriformes, pheasants, pigeons and starlings. Only Slovenia and Spain reported avian outbreaks to ADIS.

In equids, 14 MSs reported a total of 5,985 analysed samples as part of their surveillance and monitoring activities (Tables 78 and 81). Spain, Greece and Germany submitted 42.8%, 17.6% and 15.1% of these data respectively. Seven MSs reported a total of 48 equid cases that were confirmed either through IgM by ELISA (47 animals) or PCR methods (1 animal) (Table 81). As a result, 45 outbreaks were notified to ADIS by France, Germany, Greece, Hungary, Italy, Portugal and Spain. Together, Germany, Spain and Italy reported a total of 77.7% of the outbreaks in equids (Table 81). One non-EU country, Switzerland, reported 9 analyses with no outbreaks.

### 5.4.4. Joint analysis of trends and seasonality


WNV is endemic in several EU countries and both animal and human cases of WNV infections are reported every year (Figure 41). Infections in Europe occur seasonally, with most cases being reported between June and October (Figures 42 and 43). However, since the epidemic year of 2018, which was



characterised by an unusually intense transmission season, no significant decrease (p=0.89) has been observed over the past 5 years (2017–2021). Notwithstanding, the trend in the number of reported human cases of infection seems to be slowly decreasing (Figure 41), and 2021 is the year, within that period, in which reports indicated the lowest number of cases among humans (total = 158; Table 78 and Figure 42) and the lowest number of outbreaks in animals (total = 53; Table 78 and Figure 43). In Spain, a country that experienced an intense transmission season in 2020, a significant decrease in the number of cases was documented over the same period (García San Miguel Rodríguez-Alarcón et al., 2021).

The transmission season of 2021 showed a similar pattern to pre-pandemic years, with an increase of human cases in the month of July and a peak in the month of August, albeit with lower intensity.

The season started early in 2021 with the first locally acquired human case reported in June (the earlier case reported in April was in the Island of Mayotte, France and does not really reflect the EU season) and the first animal outbreak also reported in June, with cases peaking, as in previous years, in August for humans (90 reported cases, Trend Statistical Analysis (TSA), Figure 42) and in September for animals (30 outbreaks, TSA, Figure 43).




Source: Austria, Belgium, Bulgaria, Cyprus, Czechia, Estonia, Greece, Spain, Finland, France, Hungary, Ireland, Italy, Lithuania, Luxembourg, Latvia, Malta, Netherlands, Poland, Portugal, Romania, Sweden, Slovenia, Slovakia.

Figure 41: Trend in locally acquired human WNV infections reported in EU MSs, by month, 2017–2022

18314732, 2022, 12, Downloaded from https://fs.aonlinelibrary.wiley.com/inchibrary.wiley.com/inchibrary.wiley.com/em-and-conditions) on Wiley Online Library on [18/12/2023], See the Terms and Conditions (http://oinineibrary.wiley.com/em-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903/j.efsa.2022.7666 by Istituto Zooprofilattico

Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [1812/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons



The data set includes only locally acquired WNF cases and only countries that consistently reported cases (or reported zero cases) over the whole reporting period (last 5 years) and to the level of detail required for trend analysis (not aggregated).

Figure 42: Reported human cases of West Nile virus infection in the EU MSs, by month, 2017–2021


#### 5.5. Discussion

Human cases of WNV infection are reported every year by some EU MSs while in others WNV is only diagnosed in people who have travelled in endemic areas.

The number of countries reporting local WNV transmission has increased in EU MSs in recent years (Hubálek and Halouzka, 1999; Young et al., 2021). In 2020, the Netherlands reported its first cases in birds and humans (Sikkema et al., 2020). Moreover, since detecting its first epizoonosis in 2018, Germany has reported locally-acquired human and animal cases every year, suggesting the establishment of local transmission (Ziegler et al., 2020). In 2021, all the countries that reported locally acquired cases of WNV in humans had also reported locally acquired cases in the past.

The intensity of WNV transmission varies from year to year depending on environmental conditions. In 2021, the number of reported human cases and reported animal outbreaks was lower than in previous years. The number of avian cases was higher in 2021 than in 2017 and 2019, but this could be the result of more intense surveillance. That year was characterised by higher-than-average precipitation over much of Europe in winter, followed by a dry, temperature-neutral spring and a warm, rainy summer. In contrast, past epidemic years, such as 2018, with case counts up to 10 times higher than expected, were marked by a dry winter and warm spring, favouring WNV replication and transmission by mosquitoes (ECDC, 2018; Haussig et al., 2018; Riccardo et al., 2018; Beck et al., 2020; Pervanidou et al., 2020; Riccardo et al., 2020; Young et al., 2021). Even though fewer human cases and animal outbreaks have been reported in Europe over the past 3 years, we can expect to see changes in the epidemiological pattern of WNV circulation in Europe during the coming years due to an overall increase in spring temperatures.

<sup>&</sup>lt;sup>42</sup> More information is available at: https://climate.copernicus.eu/



Data source: ADIS for animal outbreaks. Outbreaks in birds or equids that were not notified to ADIS are not included. The dataset includes only locally acquired WNF cases and only countries that consistently reported cases (or reported zero cases) over the whole reporting period (last 5 years) and to the level of detail required for trend analysis (not aggregated).

Figure 43: Outbreaks of West Nile infection in animals in the EU MSs, by month, 2017–2021

The two major lineages of WNV (lineage 1 and lineage 2) are considered endemic in Europe. They are derived from a limited number of independent introductions from Africa, followed by local spread and evolution. Lineage 2 has become the dominant lineage over the past few years and has been associated with significant human epidemics (Papa et al., 2011; Hernández-Triana et al., 2014; Beck et al., 2020; Pacenti et al., 2020). Yet lineage 1 is occasionally detected, indicating that it is still circulating. During the 2021 transmission season, and 8 years after the last human case of WNV lineage 1 in Italy, the co-circulation of both lineage 1 and lineage 2 was confirmed (Veneto region) (Barzon et al., 2022), revealing the introduction/re-introduction of WNV lineage 1.

The impact of WNV on human health in EU MSs is relevant in terms of hospitalisation and case fatality. As in prior years, neuro-invasive infections continued to be the most frequently reported clinical presentation in 2021. Notwithstanding, substantial under-detection/under-reporting of clinically asymptomatic and/or mildly symptomatic WNV infections exists and should be considered when reading the data presented. Similarly, in animals, most EU countries limit screening of WNV to suspected symptomatic cases.

Integrated One Health active surveillance provides a concrete advantage in the rapid detection and characterisation of WNV circulation across the human-animal interface and can effectively guide action to ensure the safety of substances of human origin (e.g. blood and organs) and enhance risk communication (Riccardo et al., 2020). Active surveillance, with planned sampling in sentinels or wild animals, and sometimes mosquito screening, allows earlier detection of cases and is useful to indicate when and where to start the screening of blood/organ donors.



#### 6. Tularaemia

Summary data substantiating this chapter, as well as additional information on related projects and internet sources are published for this report on the EFSA Knowledge Junction on Zenodo here. Summary statistics on human surveillance data with downloadable files are retrievable using the ECDC Surveillance Atlas of Infectious Diseases here.

## 6.1. Key facts

- In 2021, the number of confirmed cases of human tularaemia was 876, corresponding to an EU notification rate of 0.20 per 100,000 population. This is an increase of 33.3% compared with the rate in 2020 (0.15 per 100,000 population), and a decrease compared with the rate in 2019 (0.25 per 100,000 population).
- Compared with the rate before the COVID-19 pandemic (2017–2019 annual mean), there was an increase of 66.7% and 42.9%, with or without the data from the United Kingdom, respectively.
- Over the past 5 years (2017–2021), a significant increasing trend (p = 0.0005) in the number of tularaemia cases has been observed.
- In 2021, the seasonal pattern was similar to that in previous years with infections peaking in summer months. Cases increased with age and were highest in the 55- to 60-year age group.
- Tularaemia in animals is rarely reported in the EU as data are submitted to EFSA on a voluntary basis. In 2021, four MSs (Austria, Finland, the Netherlands and Sweden) reported data on the occurrence of *Francisella tularensis*, mainly in hares and dogs. One non-MS (Switzerland) reported samples from wild species, zoo animals and pets.
- Austria, Finland, the Netherlands and Sweden together reported that 75 out of 335 animals tested positive, 70 of which were hares. Among pets, four dogs tested serologically positive. In Switzerland, seven (41.2%) out of 17 tested hares were positive for *Francisella tularensis*.

# 6.2. Surveillance and monitoring of tularaemia in the EU

## **6.2.1.** Humans

In 2021, 26 EU MSs provided data on tularaemia in humans. Surveillance is mandatory and surveillance systems are comprehensive with full national coverage in all reporting countries. There is no surveillance system in place for tularaemia in Denmark and the disease is not notifiable or reported at the EU level. In 2020–2021, Spain did not receive data from all regions that normally report, so the case numbers may therefore not be complete. The EU case definition was used by 25 MSs, Germany and Italy reported using other case definitions. All countries reported case-based data except Belgium, which reported aggregated data.

#### **6.2.2.** Animals

Tularaemia in animals is an internationally reportable disease (with reporting to WOAH); therefore, at European level, each country receives communications from its veterinary services. In some countries, notification is mandatory under national law. Animal monitoring data on *F. tularensis* are submitted to EFSA on a voluntary basis by EU MSs and non-MS countries. Notably, for 2021, four EU MSs (Austria, Finland, the Netherlands and Sweden) and one non-MS (Switzerland) reported these data to EFSA. Surveillance was mostly passive.

# 6.3. Results

# 6.3.1. Overview of key statistics, EU, 2017–2021

Table 82 summarises EU-level statistics on tularaemia in humans and in major animal species, respectively, for 2017–2021. More detailed descriptions of these statistics are provided in the below subsections.

18314732, 2022, 12, Downloaded from https://fs.aonlinelibrary.wiley.com/doi/10.9293/j-fs.a.2022.7566 by Istituto Zoporofilatios Sperimenta dell'Umbria e delle Marche (ZSUM), Wiley Online Library on [1812023], See the Terms and Conditions (https://onlinelibrary.wiley.com/rerms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Cerawise Common.



**Table 82:** Summary of *Francisella tularensis* statistics related to humans and the main animal species, EU, 2017–2021

species, E0, 2017–2021	î				î	1
	2021 <sup>(a)</sup>	2020	2019 <sup>(b)</sup>	2018 <sup>(b)</sup>	2017 <sup>(b)</sup>	Data source
Humans						
Total number of confirmed cases	876	641	1,279	270	323	ECDC
Total number of confirmed cases/100,000 population (notification rates)	0.20	0.15	0.25	0.05	0.06	ECDC
Number of reporting MSs	26	26	27	27	27	ECDC
Infection acquired in the EU	715	466	1,118	233	236	ECDC
Infection acquired outside the EU	1	2	3	3	2	ECDC
Unknown travel status or unknown country of infection	160	173	158	34	85	ECDC
Number of foodborne outbreak-related cases	0	0	0	0	0	EFSA
Total number of foodborne outbreaks	0	0	0	0	0	EFSA
Animals						
Hares						
Number of sampled animals	317	222	211	112	39	EFSA
Number of positive animals	70	81	67	20	7	EFSA
% of positive animals	22.1	36.5	31.8	17.9	17.9	EFSA
Number of reporting MSs	3	3	2	2	1	EFSA
Animals other than hares						
Number of sampled animals	18	5	152	0	0	EFSA
Number of positive animals	5	1	8	0	0	EFSA
% of positive animals	27.8	20.0	5.3	-	-	EFSA
Number of reporting MSs	3	1	1	0	0	EFSA

ECDC: European Centre for Disease Prevention and Control; EFSA: European Food Safety Authority; MSs: Member States.

#### 6.3.2. Tularaemia in humans

In 2021, 26 EU MSs reported a total of 876 confirmed cases of tularaemia with a notification rate of 0.20 cases per 100,000 population. This was an increase of 33.3% compared with the rate in 2020 (0.15 per 100,000 population) and an increase of 66.7% and 42.9% compared with the average annual notification rate from 2017 to 2019 (pre-pandemic period), with and without the data from the United Kingdom, respectively. In 2021, Sweden had the highest notification rate with 2.8 per 100,000 population, followed by Slovenia and Finland, with 2.6 and 1.6 per 100,000 population, respectively (Table 83).

In 2021, 715 cases (81.6%) were acquired in the EU. Austria, France, Germany and the Netherlands reported travel-associated cases. Of those, only one case was imported from outside the EU, namely Nigeria, and for 160 cases (18.3%), there were no data on travel or the country of infection (Table 82).

Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms

<sup>(</sup>a): Data on animals from the United Kingdom (Northern Ireland) are taken into account for 2021. In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/Northern Ireland, the EU requirements on data sampling are also applicable to Northern Ireland.

<sup>(</sup>b): Data from the United Kingdom are taken into account for 2017–2019, because the United Kingdom was an EU MS, but it became a third country on 1 February 2020.



**Table 83:** Reported confirmed human cases of tularaemia and notification rates per 100,000 population in EU MSs and non-MS countries, by country and by year, 2017–2021

		2021			20	20	20	19	20	18	2017	
Country	National coverage <sup>(a)</sup>	Data format <sup>(a)</sup>	Confi cases rat	and	case	rmed s and tes	case	irmed s and tes	case	irmed s and tes	case	irmed s and tes
			Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate	Cases	Rate
Austria	Υ	С	58	0.65	33	0.37	20	0.23	7	0.08	13	0.15
Belgium	Y	Α	7	0.06	1	0.01	4	0.03	0	0	5	0.04
Bulgaria	Υ	Α	0	0	2	0.03	1	0.01	1	0.01	1	0.01
Croatia	Υ	С	0	0	0	0	1	0.02	0	0	3	0.07
Cyprus	Υ	С	0	0	0	0	0	0	0	0	0	0
Czechia	Υ	С	50	0.47	67	0.63	102	0.96	32	0.30	51	0.48
Denmark <sup>(b)</sup>	_	_	_	_	-	-	-	_	-	_	-	_
Estonia	Υ	С	1	0.08	1	0.08	2	0.15	1	0.08	0	0
Finland	Υ	С	86	1.6	143	2.6	48	0.87	7	0.13	32	0.58
France	Υ	С	143	0.21	45	0.07	45	0.07	11	0.02	19	0.03
Germany	Υ	С	113	0.14	59	0.07	70	0.08	52	0.06	52	0.06
Greece	Υ	С	0	0	0	0	0	0	0	0	0	0
Hungary	Υ	С	7	0.07	20	0.20	22	0.23	17	0.17	11	0.11
Ireland	Y	С	0	0	0	0	0	0	0	0	0	0
Italy	Υ	С	3	0.01	0	0	1	< 0.01	0	0	2	< 0.01
Latvia	Υ	С	0	0	0	0	2	0.10	0	0	0	0
Lithuania	Υ	С	7	0.25	2	0.07	4	0.14	5	0.18	5	0.18
Luxembourg	Υ	С	3	0.47	0	0	0	0	0	0	0	0
Malta	Υ	С	0	0	0	0	0	0	0	0	0	0
Netherlands	Υ	С	5	0.03	1	0.01	3	0.02	2	0.01	1	0.01
Poland	Υ	С	43	0.11	5	0.01	21	0.06	16	0.04	30	0.08
Portugal	Υ	С	0	0	1	0.01	1	0.01	2	0.02	0	0
Romania	Υ	С	0	0	0	0	0	0	0	0	0	0
Slovakia	Υ	С	0	0	12	0.22	20	0.37	6	0.11	2	0.04
Slovenia	Υ	С	54	2.6	1	0.05	7	0.34	4	0.19	1	0.05
Spain	Υ	С	4	0.01	1	< 0.01	88	0.19	4	0.01	11	0.02
Sweden	Υ	С	292	2.8	247	2.4	817	8.0	102	1.0	84	0.84
EU Total 27	_	-	876	0.20	641	0.15	1,279	0.29	269	0.06	323	0.07
United Kingdom	-	_	_	_	_	_	0	0	1	< 0.01	0	0
EU Total <sup>(c)</sup>	-	-	876	0.20	641	0.15	1,279	0.25	270	0.05	323	0.06
Iceland	Υ	С	0	0	0	0	0	0	0	0	0	0
Norway	Υ	С	95	1.8	99	1.8	183	3.4	58	1.1	92	1.7
Liechtenstein	Υ	С	1	2.6	133	1.5	154	1.8	117	1.4	152	1.8
Switzerland <sup>(d)</sup>	Υ	С	212	2.4								

<sup>-:</sup> Data not reported.

Tularaemia shows a seasonal pattern, with most cases occurring from July to November, but some cases are also observed in winter. In 2021, infections peaked in summer months, in line with the mean for the 2017–2020 period (Figure 44).

In 2021, the number of tularaemia cases increased compared with 2020 and vs. the 2017–2019 prepandemic period. Over the past 5 years, from 2017 to 2021, a significant increasing trend in the number


<sup>(</sup>a): Y: yes; N: no; A: aggregated data; C: case-based data.

<sup>(</sup>b): No surveillance system.

<sup>(</sup>c): Cases reported by the United Kingdom for the period 2017–2019 were also considered for this estimation (EU-28). When the United Kingdom data were collected for the period 2017–2019, the United Kingdom was an EU MS, but it became a third country on 1 February 2020.

<sup>(</sup>d): Switzerland provided data directly to EFSA. The human data for Switzerland include data from Liechtenstein for the years 2017–2020.

of tularaemia cases was observed in the EU (Figure 44). At the country level, a statistically significant (p < 0.01) increasing trend was observed in Austria, Czechia, France, Germany and Sweden.



Source: Austria, Cyprus, Czechia, Estonia, Finland, France, Germany, Greece, Hungary, Ireland, Latvia, Luxembourg, Malta, Poland, Romania, Slovakia, Slovenia, Spain and Sweden. Belgium, Bulgaria, Croatia, Denmark, Italy, Lithuania, the Netherlands and Portugal did not report data to the level of detail required for the analysis.

**Figure 44:** Trends in reported confirmed human cases of tularaemia in the EU, by month and year, 2017–2021

Data on hospitalisation status were provided by 10 MSs for 221 confirmed cases. A total of 112 hospitalisations were reported by eight MSs (Austria, Czechia, Hungary, Lithuania, the Netherlands, Poland, Slovenia and Spain), corresponding to 50.7% of confirmed cases with reported hospitalisation status; more than 35% were counted in the 45- to 60-year age group. The highest hospitalisation rates were reported in Lithuania (100%), Austria (64%) and Poland (55.8%). In 2021, 11 MSs reported outcomes for 341 cases (38.9%), with two deaths recorded (> 70 years) and an EU case fatality rate of 0.59%.

The proportion of male cases was higher, with a male-to-female ratio of 1.8:1. Children under 14 years of age accounted for 44 cases (5%). The number of cases increased with age up to 60 years and slowly decreased after 60 years. The highest notification rate was for the 55- to 60-year age group, followed by the 60- to 65-year age group.

#### 6.3.3. Tularaemia in animals

In 2021, overall, four EU MSs reported 70 hares, four dogs and one muskrat positive for tularaemia, while Switzerland (not an EU MS) reported 10 positive animals (Table 84).

The number of samples and number of animal species tested were greater than in 2020. Moreover, one additional MS country (the Netherlands) reported data.

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/continelibrary.wiley.c



**Table 84:** Francisella tularensis monitoring results for animals, by species in reporting EU MSs and non-MS countries, 2021

Country	N positive/N tested (% positive)										
	Domestic animals				Wild animals					Zoo animals	
	Cats	Dogs	<b>Guinea pigs</b>	Rabbits	Hares	Beavers	Foxes	Muskrats	Shrews	Hares	Other zoo animals
Austria	_	-	_	_	20/52 (38.5)	-	_	_	_	_	_
Finland	_	_	_	_	30/199 (15.1)	_	_	1/1 (100)	_	_	_
Netherlands	_	_	_	_	_	_	_	0/2 (0)	_	_	_
Sweden	_	4/11 (36.4)	0/1 (0)	0/2 (0)	20/66 (30.3)	_	_	_	0/1 (0)	_	_
EU Total (27 + XI)	_	4/11 (36.4)	0/1 (0)	0/2 (0)	70/317 (22.1)	_	_	1/3 (33.3)	0/1 (0)	-	_
Switzerland	1/5 (20.0)	0/1 (0)	_	0/3 (0)	7/17 (41.2)	0/2 (0)	0/1 (0)	-	_	1/3 (33.3)	1/7 (14.3)
Total EU (27+ XI) + non-EU countries	1/5 (20.0)	4/12 (33.3)	0/1 (0)	0/5 (0)	77/334 (23.1)	0/2 (0)	0/1 (0)	1/3 (33.3)	0/1 (0)	1/3 (33.3)	1/7 (14.3)

#### 6.4. Discussion

Francisella tularensis is the causative agent of tularaemia and is also considered a potential biological weapon. Humans can acquire the disease through several routes. The disease shows a seasonal pattern in humans (Hestvik et al., 2015), consistent with a higher likelihood of exposure in summer and autumn months due to recreational outdoor activities (notably hunting), exposure to contaminated water (Hennebique et al., 2019) and vector bites. Tularaemia is widely distributed throughout most of Europe, and in endemic regions within Scandinavian countries, it is typically transmitted by mosquito bites (Kenney et al., 2017). Notification rates for tularaemia vary among MSs and over time. Sweden has reported cases of tularaemia since 1931 and recorded two major outbreaks in 2015 and 2019 (Dryselius et al., 2019), whereas in 2016, Finland had the highest notification rate. In 2018, an outbreak occurred in France, whereas in 2019, Norway and Sweden reported highest number of cases. In 2020, the number of human cases in the EU was about half that observed in 2019.

In 2021, the number of human cases increased in the EU compared with the 2020 pandemic year but also vs. the 2017–2019 pre-pandemic period. More than 70% of cases were reported from Finland, France, Germany and Sweden. In particular, France recorded an increasing number of cases over a 5-year period, mainly due to the change of the case definition used in 2021 (EU case definition), being less restrictive for confirmed cases. However, from 2018, a significant increase in tularaemia cases was noted in France. In particular, the Brittany region reported more pulmonary forms than other regions, but the reasons for this deviating pattern are unknown (Figoni et al., 2022). Overall, from 2017 to 2021, there was a statistically significant increase in the trend for confirmed cases in the EU. Moreover, no data were available for 2021 regarding the impact of the COVID-19 pandemic on tularaemia surveillance and reporting.

Wildlife continues to play a role in maintaining *F. tularensis* in the ecological cycle, and also in the occurrence of human cases. *Francisella* spp. are present in wild animals (such as hares) and in vectors (e.g. ticks and mosquitos), which can be sources of infections for humans (WHO, 2007; Maurin and Gyuranecz, 2016).

Lately, the circulation of *F. tularensis* among wild animals has been reported in numerous countries in north-central Europe, where it is considered endemic (Hestvik et al., 2015; Faber et al., 2018; Seiwald et al., 2020), as well as in Spain (Mínguez-González et al., 2021).

In the last 5 years, among the reporting MSs, the number of hares tested has increased (from 39 to 317) and the rate of positivity has ranged from 17.9% to 36.5%.

The reporting of positive dogs (4/11) is interesting because dogs live in close contact with humans and may therefore share exposure to *F. tularensis* (Kwit et al., 2020).

As tularaemia surveys are very often passive, they do not reflect the status of the entire population, so it is difficult to paint an accurate picture of the spread of the disease among animals. For example, the positivity rate reported in dogs in 2021 is not indicative of the spread of the disease in this population.

It should be highlighted that risks of exposure and/or new outbreaks in humans are often preceded by the appearance of the disease in animals, which means that wildlife monitoring (hares are good indicators) is crucial.

Tularaemia is a disease with a multifaceted epidemiology; it is therefore difficult to control. All these aspects underline the importance of collaborative work between public health and veterinary units for the control of this zoonotic disease.

## 7. Other zoonoses and zoonotic agents

In 2021, data on *Bacillus, Chlamydia, Clostridium, Cronobacter sakazakii, Klebsiella*, non-pathogenic *Escherichia coli, Proteus, Shigella, Staphylococcus, Streptococcus, Vibrio*, caliciviruses, flaviviruses other than West Nile virus, hepatitis virus, *Cysticercus, Leishmania, Leptospira* and *Sarcocystis* were reported to EFSA.

## 7.1. Bacillus spp.

Luxembourg and Portugal submitted 2021 data on *Bacillus* spp. in various foods collected at hospitals or medical care facilities, retail establishments, collective catering establishments (restaurant or cafe or pub or bar, or hotel or catering service) and wholesale establishments. Out of 165 batches

1831472, 2022, 12, Downloaded from https://efs.aonlinelibray.wiley.com/doi/10/2033/jefs.a2022/7666 by Istituto Zooporficiation Sperimenta dell'Umbria e delle Marche (ZSUM), Wiley Online Library on [18/12/2023], See the Terms and Conditions (http://onlinelibray.wiley.com/rems-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Cerawise Commons of the Common of the Condition of the Common of the Com



and 1,130 single samples, 29 single samples (2.6%) tested positive. The positive food categories involved 'other processed food products and prepared dishes' or 'sauces and dressings'.

Greece reported one positive bovine and one positive sheep sample (15.4%) for *Bacillus anthracis* in 13 samples collected at the farm level during clinical investigations.

# 7.2. Caliciviruses (including norovirus)

Five MSs (Bulgaria, Croatia, France, Portugal and Romania) reported data on caliciviruses for a total of six batch samples and 937 single samples. Only one sample (0.1%) of non-pre-cut fruit collected at the retail level in France was positive.

# 7.3. Chlamydia spp.

Three MSs (Austria, Denmark and Greece) and one non-MS (North Macedonia) reported data on *Chlamydia* spp. in various animal species. Austria reported 106 (4.6%) positives out of 2,297 samples, Denmark reported 26 (24.5%) positives out of 106 samples and North Macedonia reported 12 (29.3%) positives out of 41 samples. Greece reported no positives from 12 tested samples.

# 7.4. Clostridium spp.

Greece and non-MS North Macedonia reported data on *Clostridium* spp. from various ruminant species for a total of 54 samples. Greece detected 30 (56.6%) positives out of 53 animal samples collected during clinical investigations and North Macedonia detected one (100%) positive sample from the one sample collected during monitoring.

Three MSs (Lithuania, Romania and Slovenia) and non-MS North Macedonia provided data on *Clostridium* spp. in foods for a total of 302 tested sampling units. In an investigation at the manufacturing level on five samples, four (80.0%) positives were found in Lithuania, and the identified species was *Clostridium perfringens*.

## 7.5. Hepatitis virus

Three MSs (Bulgaria, France and Romania) provided data on hepatitis virus in non-pre-cut fruit and leaf vegetables. None of the 153 tested samples were positive.

# 7.6. *Proteus* spp.

Greece tested for the presence of *Proteus* spp. in 179 samples of milk and organs or tissues collected from various animals during clinical investigations, resulting in 11 (6.1%) positives.

## 7.7. Staphylococcus spp. and staphylococcal enterotoxins

Four MSs (Bulgaria, Germany, Greece and Italy) provided data on Staphylococcus spp. (reported as Staphylococcus, Staphylococcus spp. unspecified or S. aureus) in various food matrices (N = 7,734) and animals (N = 2,491). Overall, 16.5% of food and 66.4% of animal sampling units were reported positive. 'Milk from other animal species or unspecified — pasteurised milk', 'cheeses, made from unspecified milk or other animal milk — unspecified', 'bakery products — pastry' and 'other processed food products and prepared dishes — pasta' were the food categories with the highest numbers of positive results.

Thirteen MSs reported data on staphylococcal enterotoxins collected in contexts other than those stipulated in Regulation (CE) No 2073/2005. None of the 118 tested batches were positive, whereas 12 out of 3,897 (0.3%) single samples were positive. 'Cheeses, made from unspecified milk or other animal milk' collected in Italy at the processing plant was the food category positive for staphylococcal enterotoxins.

## 7.8. *Cysticercus* spp.

Eight MSs (Belgium, Finland, Luxembourg, Malta, Slovakia, Slovenia, Spain and Sweden) submitted data on *Cysticercus* spp. in various animal species. Belgium collected 770,235 bovine carcases from slaughterhouses, revealing 857 positive samples (0.111%). None of the 2,200,672 carcases from cattle, pigs and wild boars collected in Finland were positive. Luxembourg found 74 positive bovine carcases out of the 27,326 collected samples (0.271%). None of the 65,334 cattle, pig, sheep or goat

18314732, 2022, 12, Downloaded from https://fs.aonlinelibrary.wiley.com/continelibrary.wiley.com

carcases collected in Malta were positive. Slovakia found three positive cattle carcases out of 34,771 (0.009%) and seven pig carcases out of 675,234 (0.001%) samples collected at the slaughterhouse. Slovenia provided results on 123,961 cattle and 242,584 pig carcases collected at the slaughterhouse, detecting 11 positive samples in cattle carcases (0.009%). Spain provided data on *Cysticercus* spp. in various animal species: 125 out of 2,332,666 cattle (0.005%), 17,332 out of 799,767 goats (2.18%), 2,902 out of 41,059,466 pigs (0.007%), 200,810 out of 7,077,050 sheep (2.84%), 110 out of 4,544 solipeds (2.42%) and 33 of 111,100 wild boars (0.03%) tested positive. No positives were detected upon testing 7,415 mouflons and 118,899 deer. Sweden detected one positive out of 411,650 cattle carcases and no positives from 2,651,110 pig carcases collected at the slaughterhouse.

#### 7.9. Leishmania

Greece and non-MS North Macedonia provided data on *Leishmania* collected from 6,006 (22.0%) dog blood samples, detecting 1,324 positive samples.

# 7.10. Sarcocystis spp.

Belgium reported data from 770,235 cattle carcases collected at the slaughterhouse, whereof 81 (0.01%) were positive for *Sarcocystis* spp.

#### **7.11.** Other

Twelve MSs reported data on *Cronobacter sakazakii* collected in contexts other those stipulated in Regulation (EC) No 2073/2005. 'Infant formulae', 'dried dietary foods for special medical purposes intended for infants below 6 months of age', 'dairy products intended for infants and young children', 'processed cereal-based food for infants and young children', 'milk and whey powder', 'ice cream', 'follow-on formulae' and 'ready-to-eat infant formulae' were collected as single samples and batches at the processing plant, retail, wholesale, hospital or medical care facility and farm and road transport levels. Three (0.25%) of the 1,205 single samples and three (0.74%) of the 404 batches were positive.

Two MSs (Greece and Latvia) provided data on *Escherichia coli* and non-pathogenic *E. coli* collected at the farm and slaughterhouse levels, detecting 293 (74.5%) positives out of 393 tested samples.

Slovenia tested 41 milk and milk products collected at the retail level to verify the presence of flaviviruses other than West Nile virus, obtaining no positive samples. Four MSs (Ireland, Italy, the Netherlands and Slovakia) and the non-MS Switzerland reported data collected from various animal categories. None of 12,474 collected samples tested positive.

Greece reported data on *Klebsiella* spp. in milk and organs or tissues collected from various ruminants (dairy cows, goats, sheep), obtaining four (2.2%) positives out of 179 tested samples.

Greece provided data on *Streptococcus* spp. in milk and organs or tissues collected from cattle and small ruminants, obtaining 37 (18.1%) positives out of 204 tested samples.

The Netherlands tested 750 single samples or batches of crustaceans, fishes, live bivalve molluscs and leaf vegetables to verify the presence of *Vibrio* spp. *Vibrio* cholerae was detected in 21 samples, whereas *Vibrio* parahaemolyticus was detected in 45 samples, for a total of 66 (8.8%) positives.

Slovenia provided data on *Leptospira* in various animal species, with no positives detected from 146 collected samples.

18314732, 2022, 12. Downloaded from https://eis.ao.nlinelibrary.wiley.com/doi/10.2903j.cfs.a.2022.7266 by Isitiuto Zoprofilatios Sperimenta dell'Umbria e delle Marche (ZSUM). Wiley Online Library on 11812023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/rems-and-conditions) on Wiley Online Library for rules of use O.A articles are governed by the applicable Creative Commons



# Microbiological contaminants subject to food safety criteria (Regulation (EC) No 2073/2005)

This chapter summarises the 2021 information and data provided by reporting countries on microbiological contaminants in food for which food safety criteria (FSC) have been set down in EU legislation (Regulation (EC) No 2073/2005), histamine, staphylococcal enterotoxins and *Cronobacter sakazakii*.

#### 1. Histamine

# 1.1. Histamine data in the context of Regulation (EC) No 2073/2005

Histamine is a thermostable biogenic amine occurring naturally in the human body. However, its ingestion at high concentrations through food is associated with the onset of health disorders such as scombroid poisoning.

Regulation (EC) No 2073/2005 on microbiological criteria for foodstuffs defines three categories concerning FSC for histamine in food at the retail level: 'fishery products from fish species associated with a high amount of histidine' (food category 1.26), 'fishery products which have undergone enzyme maturation treatment in brine, manufactured from fish species associated with a high amount of histidine' (food category 1.27) and 'fish sauce produced by fermentation of fishery products' (food category 1.27a). Information was also obtained at the manufacturing level, indicating the correct application of GMP (good manufacturing practices) and the proper maintenance of the cold chain, which is essential to avoid the development or increase of histamine in fish and fish products.

Data on histamine were reported by 15 MSs (Belgium, Cyprus, Czechia, Denmark, Estonia, France, Germany, Greece, Latvia, Luxembourg, Portugal, Romania, Slovakia, Slovenia and Spain) and two non-MSs (Iceland and Serbia).

In official control samples (n = 3,154) for histamine in food category **1.26** at the distribution level (wholesale establishments, retail establishments, border control posts and restaurants), 0.89% had a histamine content higher than 200 mg/kg, 0.98% a histamine content of between 100 and 20 mg/kg and 35.76% a histamine content above the limit of detection, but less than or equal to 100 mg/kg. Of the total number of samples, 13.9% were of EU origin (Latvia, Romania, European Union), 21.7% were of non-EU origin (Cape Verde, Colombia, Ecuador, El Salvador, Indonesia, Mauritius, Morocco, Papua New Guinea, Philippines, Senegal, Seychelles, Taiwan, Thailand, Vietnam, non-EU origin), while for 64.2% no information was available. Of the total number of official control samples, 15.6% came from canned fish and 3.9% from raw fish, while for 80% no information was provided.

At the manufacturing level (processing plants, packaging centres), 2,178 official control sampling units were collected and the results were as follows: 0.14% had a histamine content higher than 200 mg/kg, 0.18% a histamine content of between 100 and 200 mg/kg and 60.2% a histamine content higher than the limit of detection, but less than or equal to 100 mg/kg. Of the total number of samples, 30% were of EU origin (Estonia, European Union, Latvia, Portugal, Spain), while for 67.2% no information was reported. Of the total number of control sample units, 24.8% came from canned fish and 17.9% from raw fish, while for 57.1% no information was reported.

For food category **1.27**, 1,223 and 162 official control sample units were collected at the distribution and manufacturing levels, respectively. At the distribution level, 63.2% of the samples had a histamine concentration less than or equal to 200 mg/kg and 0.33% a histamine content between 200 and 400 mg/kg. At the manufacturing level, the percentage of samples with a histamine concentration less than or equal to 200 mg/kg was 44.4%. At the distribution level, 7.35% of samples were of EU origin, 8.1% were of non-EU origin, while for 84% no information was reported.

For food category **1.27a** at the distribution level, 6 official control samples were reported: 67% of samples had a histamine content of less than 400 mg/kg but above the limit of detection; at the manufacturing level, 1 sample was collected and had a histamine content below 400 mg/kg.

# 1.2. Other monitoring or surveillance data for histamine

MSs also collected and analysed fishery products in contexts other than Regulation (EC) 2073/2005 on microbiological criteria for foodstuffs, such as Surveillance and Monitoring.

18314732, 2022, 12, Downloaded from https://fs.aonlinelibrary.wiley.com/doi/10.9293/j-fs.a.2022.7566 by Istituto Zoporofilatios Sperimenta dell'Umbria e delle Marche (ZSUM), Wiley Online Library on [1812023], See the Terms and Conditions (https://onlinelibrary.wiley.com/rerms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Cerawise Common.



A total of 2,162 samples and 20 batches were collected as part of Surveillance and Monitoring activities, respectively.

#### Monitoring

A total 20 batches were collected at the distribution level. Of this number, 60% of batches were of EU origin and 40% of non-EU origin. All the batches fell into category 1.26 and had a histamine content of less than 100 mg/kg.

#### Surveillance

A total of 1,803 and 359 sampling units were collected at the distribution and manufacturing levels, respectively.

At the distribution level, 99% of samples were taken at border control posts; 5% and 95% of the collected samples were classified in categories 1.26 and 1.27, respectively. At the manufacturing level, all samples were taken at processing plants; 87% and 13% of the samples were classified in categories 1.26 and 1.27, respectively.

All samples taken (n = 2,162) were negative.

# 2. Staphylococcal enterotoxins

Data on staphylococcal enterotoxins were reported by seven MSs (Bulgaria, Croatia, Cyprus, Estonia, Romania, Slovakia and Spain). No positives were reported out of 1,391 collected official control samples.

#### 3. Cronobacter sakazakii

Data on *Cronobacter sakazakii* in 'infant formulae' and 'foodstuffs intended for special nutritional uses – dried dietary foods for special medical purposes intended for infants under 6 months of age' collected at the distribution level (retail and wholesale) were reported by four MSs (Croatia, Hungary, Slovenia, Slovakia). No positives were detected out of the 364 collected official control samples.

Estonia and Spain collected samples at the processing plant, detecting one (3.1%) positive out of 32 tested official control samples.

# Note on outbreak data of SARS-CoV-2 in minks, mustelids and raccoon dogs, in EU

In accordance with Commission Implementing Decision (CID) (EU) 2021/788<sup>43</sup> Member States monitor and report infections with SARS-CoV-2 in minks, mustelids and raccoon dogs to the European Commission. The European Commission forwards these reports to EFSA, who, in accordance with Article 9(2) of Directive 2003/99/EC is required to summarise these outbreak data in the EU One Health Zoonoses report. EFSA and the European Commission agreed that the summary of these data will be made available in the EFSA scientific opinion 'SARS-CoV-2 in animals: susceptibility of animal species, monitoring, prevention and control' to be released by EFSA in January 2023.

#### References

Adamse P, Van Egmond H, Noordam M, Mulder P and De Nijs M, 2014. Tropane alkaloids in food: poisoning incidents. Quality Assurance and Safety of Crops & Foods, 6, 15–24. https://doi.org/10.3920/QAS2013.0314
Adinolfi F, Di Pasquale J and Capitanio F, 2016. Economic issues on food safety. Ital J Food Saf, 5, 5580. https://doi.org/10.4081/ijfs.2016.5580

Alban L, Pozio E, Boes J, Boireau P, Boué F, Claes M, Cook AJ, Dorny P, Enemark HL, van der Giessen J, Hunt KR, Howell M, Kirjusina M, Nöckler K, Rossi P, Smith GC, Snow L, Taylor MA, Theodoropoulos G, Vallée I, Viera-Pinto MM and Zimmer IA, 2011. Towards a standardised surveillance for *Trichinella* in the European Union. Prev Vet Med, 99, 148–160. https://doi.org/10.1016/j.prevetmed.2011.02.008

Alegbeleye OO, Singleton I and Sant'Ana AS, 2018. Sources and contamination routes of microbial pathogens to fresh produce during field cultivation: a review. Food Microbiol, 73, 177–208. https://doi.org/10.1016/j.fm. 2018.01.003

18314732, 2022, 12, Downloaded from https://fs.aonlinelibrary.wiley.com/doi/10.9293/j-fs.a.2022.7566 by Istituto Zoporofilatios Sperimenta dell'Umbria e delle Marche (ZSUM), Wiley Online Library on [1812023], See the Terms and Conditions (https://onlinelibrary.wiley.com/rerms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Cerawise Common.

<sup>&</sup>lt;sup>43</sup> Commission Implementing Decision (EU) 2021/788 laying down rules for the monitoring and reporting of infections with SARS-CoV-2 in certain animal species. OJ L 173, 17.5.2021, pp. 6–14.



- Aroussi A, Vignoles P, Dalmay F, Wimel L, Dardé ML, Mercier A and Ajzenberg D, 2015. Detection of Toxoplasma gondii DNA in horse meat from supermarkets in France and performance evaluation of two serological tests. Parasite, 22, 14. https://doi.org/10.1051/parasite/2015014
- Balog T, Nagy G, Halász T, Csányi E, Zomborszky Z and Csivincsik Á, 2021. The occurrence of *Echinococcus* spp. in golden jackal (Canis aureus) in southwestern Hungary: should we need to rethink its expansion? Parasitology International, 80, 102214. https://doi.org/10.1016/j.parint.2020.102214
- Barzon L, Montarsi F, Quaranta E, Monne I, Pacenti M, Michelutti A, Toniolo F, Danesi P, Marchetti G, Gobbo F, Sinigaglia A, Riccetti S, Dal Molin E, Favero L, Russo F and Capelli G, 2022. Early start of seasonal transmission and co-circulation of West Nile virus lineage 2 and a newly introduced lineage 1 strain, northern Italy, June 2022. Euro Surveill, 27. https://doi.org/10.2807/1560-7917.Es.2022.27.29.2200548
- Beck C, Leparc Goffart I, Franke F, Gonzalez G, Dumarest M, Lowenski S, Blanchard Y, Lucas P, Lamballerie X, Grard G, Durand GA, Zientara S, Tapprest J, L'Ambert G, Durand B, Desvaux S and Lecollinet S, 2020. Contrasted epidemiological patterns of west nile virus lineages 1 and 2 infections in France from 2015 to 2019. Pathogens, 9, 908. https://doi.org/10.3390/pathogens9110908
- Bergis H, Bonanno L, Asséré A and Lombard B, 2021. EURL Lm technical guidance document on challenge tests and durability studies for assessing shelf-life of ready-to-eat foods related to Listeria monocytogenes; c2021 [cited 4 Aug 2021]. Available online: https://food.ec.europa.eu/system/files/2021-07/biosafety\_fh\_mc\_techguide-doc\_listeria-in-rte-foods\_en\_0.pdf
- Blaga R, Aubert D, Thébault A, Perret C, Geers R, Thomas M, Alliot A, Djokic V, Ortis N, Halos L, Durand B, Mercier A, Villena I and Boireau P, 2019. Toxoplasma gondii in beef consumed in France: regional variation in seroprevalence and parasite isolation. Parasite, 26, 77. https://doi.org/10.1051/parasite/2019076
- Bless PJ, Schmutz C and Mäusezahl D, 2017. The recurrent campylobacteriosis epidemic over Christmas and New Year in European countries, 2006-2014. BMC Res Notes, 10, 266. https://doi.org/10.1186/s13104-017-2587-8
- Boelaert F, Amore G, Van der Stede Y and Hugas M, 2016. EU-wide monitoring of biological hazards along the foodchain: achievements, challenges and EFSA vision for the future. Current Opinion in Food Science, 12, 52–62. https://doi.org/10.1016/j.cofs.2016.08.004
- Bonardi S, Blagojevic B, Belluco S, Roasto M, Gomes-Neves E and Vågsholm I, 2021. Food chain information in the European pork industry: where are we? Trends in Food Science & Technology, 118, 833–839. https://doi.org/10.1016/j.tifs.2021.10.030
- Boni M, Davoust B, Tissot-Dupont H and Raoult D, 1998. Survey of seroprevalence of Q fever in dogs in the southeast of France, French Guyana, Martinique, Senegal and the Ivory Coast. Vet Microbiol, 64, 1–5. https://doi.org/10.1016/s0378-1135(98)00247-8
- Buchanan RL, Gorris LG, Hayman MM, Jackson TC and Whiting RC, 2017. A review of Listeria monocytogenes: an update on outbreaks, virulence, dose-response, ecology, and risk assessments. Food control, 75, 1–13. https://doi.org/10.1016/j.foodcont.2016.12.016
- Byrne AW, Barrett D, Breslin P, O'Keeffe J, Murphy KJ, Conteddu K, Morera-Pujol V, Ryan E and Ciuti S, 2022. Disturbance ecology meets bovine tuberculosis (bTB) Epidemiology: a before-and-after study on the association between forest clearfelling and bTB herd risk in cattle herds. Pathogens, 11, 807. https://doi.org/10.3390/pathogens11070807
- Canals A, Martínez CV, Diogène J, Gago-Martínez A, Cebadera-Miranda L, de Vasconcelos FM, Gómez IL, Sánchez EVM, Alférez RC and Núñez D, 2021. Risk characterisation of ciguatera poisoning in Europe. EFSA Supporting Publications, 18, 6647E. https://doi.org/10.2903/sp.efsa.2021.EN-6647
- Carrié P, Barry S, Rousset E, de Crémoux R, Sala C, Calavas D, Perrin JB, Bronner A, Gasqui P, Gilot-Fromont E, Becker CAM, Gache K and Jourdain E, 2019. Swab cloths as a tool for revealing environmental contamination by Q fever in ruminant farms. Transbound Emerg Dis, 66, 1202–1209. https://doi.org/10.1111/tbed.13137
- Casulli A, 2020. Recognising the substantial burden of neglected pandemics cystic and alveolar echinococcosis. The Lancet Global Health, 8, e470–e471. https://doi.org/10.1016/S2214-109X(20)30066-8
- Casulli A, Massolo A, Saarma U, Umhang G, Santolamazza F and Santoro A, 2022. Species and genotypes belonging to *Echinococcus* granulosus sensu lato complex causing human cystic echinococcosis in Europe (2000-2021): a systematic review. Parasit Vectors, 15, 109. https://doi.org/10.1186/s13071-022-05197-8
- Cegar S, Kuruca L, Vidovic B, Antic D, Hauge SJ, Alvseike O and Blagojevic B, 2022. Risk categorisation of poultry abattoirs on the basis of the current process hygiene criteria and indicator microorganisms. Food control, 132, 108530. https://doi.org/10.1016/j.foodcont.2021.108530
- Chanamé Pinedo L, Mughini-Gras L, Franz E, Hald T and Pires SM, 2022. Sources and trends of human salmonellosis in Europe, 2015–2019: an analysis of outbreak data. International Journal of Food Microbiology, 379, 109850. https://doi.org/10.1016/j.ijfoodmicro.2022.109850
- Ciaravino G, Laranjo-González M, Casal J, Sáez-Llorente JL and Allepuz A, 2021. Most likely causes of infection and risk factors for tuberculosis in Spanish cattle herds. Vet Rec, 189, e140. https://doi.org/10.1002/vetr.140
- Clayton D and Hills M, 2013. Statistical models in epidemiology. OUP Oxford, Chapter 11.2 pp.
- Conraths FJ, Probst C, Possenti A, Boufana B, Saulle R, La Torre G, Busani L and Casulli A, 2017. Potential risk factors associated with human alveolar echinococcosis: systematic review and meta-analysis. PLoS Neglected Tropical Diseases, 11, e0005801. https://doi.org/10.1371/journal.pntd.0005801



- Crowe SJ, Bottichio L, Shade LN, Whitney BM, Corral N, Melius B, Arends KD, Donovan D, Stone J, Allen K, Rosner J, Beal J, Whitlock L, Blackstock A, Wetherington J, Newberry LA, Schroeder MN, Wagner D, Trees E, Viazis S, Wise ME and Neil KP, 2017. Shiga toxin-producing *E. coli* infections associated with flour. N Engl J Med, 377, 2036–2043. https://doi.org/10.1056/NEJMoa1615910
- Cuperlovic K, Djordjevic M and Pavlovic S, 2005. Re-emergence of trichinellosis in southeastern Europe due to political and economic changes. Vet Parasitol, 132, 159–166. https://doi.org/10.1016/j.vetpar.2005.05.047
- De Massis F, Sacchini F, Averaimo D, Garofolo G, Lecchini P, Ruocco L, Lomolino R, Santucci U, Sgariglia E, Crotti S, Petrini A, Migliorati G, D'Alterio N, Gavaudan S and Tittarelli M, 2021. First isolation of Brucella canis from a breeding kennel in Italy. Vet Ital, 57, 3. https://doi.org/10.12834/VetIt.2497.15848.1
- Deplazes P, Hegglin D, Gloor S and Romig T, 2004. Wilderness in the city: the urbanization of *Echinococcus multilocularis*. Trends Parasitol, 20, 77–84. https://doi.org/10.1016/j.pt.2003.11.011
- van Deursen B, Hagenaars M, Meima A, van Asten L, Richardus JH, Fanoy E and Voeten H, 2022. A sharp decrease in reported non-COVID-19 notifiable infectious diseases during the first wave of the COVID-19 epidemic in the Rotterdam region, the Netherlands: a descriptive study. BMC Infect Dis, 22, 208. https://doi.org/10.1186/s12879-022-07209-5
- van Dijk MAM, Engelsma MY, Visser VXN, Keur I, Holtslag ME, Willems N, Meij BP, Willemsen PTJ, Wagenaar JA, Roest HIJ and Broens EM, 2021. Transboundary spread of Brucella canis through import of infected dogs, the Netherlands, November 2016–December 2018. Emerg Infect Dis, 27, 1783–1788. https://doi.org/10.3201/eid2707.201238
- DIN (Deutsches Institut Fur Normung E.V. German National Standard-), 2004a. DIN 10118:2004-06. Analysis of foodstuffs detection of verotoxine-forming Escherichia (E.) coli-strains (VTEC) in food derived from animals.
- DIN (Deutsches Institut Fur Normung E.V. German National Standard-), 2004b. DIN 10167:2004 DE. Nachweis von Escherichia coli O157 in Fleisch und Fleischerzeugnissen. Detection of Escherichia coli O157 in meat and meat products.
- Djordjevic M, Bacic M, Petricevic M, Cuperlovic K, Malakauskas A, Kapel CM and Murrell KD, 2003. Social, political, and economic factors responsible for the reemergence of trichinellosis in Serbia: a case study. J Parasitol, 89, 226–231. https://doi.org/10.1645/0022-3395(2003)089[0226:Spaefr]2.0.Co;2
- Dryselius R, Hjertqvist M, Mäkitalo S, Lindblom A, Lilja T, Eklöf D and Lindström A, 2019. Large outbreak of tularaemia, central Sweden, July to September 2019. Euro Surveill, 24. https://doi.org/10.2807/1560-7917.Es. 2019.24.42.1900603
- Dušek D, Vince A, Kurelac I, Papić N, Višković K, Deplazes P and Beck R, 2020. Human alveolar Echinococcosis, Croatia. Emerging Infectious Diseases, 26, 364–366. https://doi.org/10.3201/eid2602.181826
- Ebani VV, Bertelloni F, Najar B, Nardoni S, Pistelli L and Mancianti F, 2020. Antimicrobial activity of essential oils against staphylococcus and malassezia strains isolated from canine dermatitis. Microorganisms, 8, 252. https://doi.org/10.3390/microorganisms8020252
- ECDC (European Centre for Disease Prevention and Control), 2018. Early large increase in West Nile virus infections in the EU/EEA and EU neighbouring countries. Available online: https://www.ecdc.europa.eu/en/publications-data/rapid-risk-assessment-early-large-increase-west-nile-virus-infections-reported
- ECDC (European Centre for Disease Prevention and Control), 2019. ECDC strategic framework for the integration of molecular and genomic typing into European surveillance and multi-country outbreak investigations 2019–2021. https://doi.org/10.2900/805317 Available online: https://www.ecdc.europa.eu/en/publications-data/ecdc-strategic-framework-integration-molecular-and-genomic-typing-european
- ECDC and EFSA (European Centre for Disease Prevention and Control and European Food Safety Authority), 2018a. Multi-country outbreak of Listeria monocytogenes serogroup IVb, multi-locus sequence type 6, infections probably linked to frozen corn. EFSA Supporting Publications. 2397-8325. https://doi.org/10.2903/sp.efsa.2018.EN-1402
- ECDC and EFSA (European Centre for Disease Prevention and Control and European Food Safety Authority), 2018b. Multi-country outbreak of Listeria monocytogenes sequence type 8 infections linked to consumption of salmon products. EFSA Supporting Publications . 2397-8325. https://doi.org/10.2903/sp.efsa.2018.EN-1496
- ECDC (European Centre for Disease Prevention and Control) and EFSA (European Food Safety Authority), 2019. Multi-country outbreak of Listeria monocytogenes sequence type 6 infections linked to ready-to-eat meat products. https://doi.org/10.2903/sp.efsa.2019.EN-1745
- ECDC (European Centre for Disease Prevention and Control) and EFSA (European Food Safety Authority), 2021a. Multi-country outbreak of *Salmonella* Braenderup ST22, presumed to be linked to imported melons. EFSA Supporting Publications. Available online: https://www.efsa.europa.eu/en/efsajournal/pub/en-6807
- ECDC (European Centre for Disease Prevention and Control) and EFSA (European Food Safety Authority), 2021b. Multi-country outbreak of multiple *Salmonella* enterica serotypes linked to imported sesame-based products. EFSA Supporting Publications, 18, 6922E. https://doi.org/10.2903/sp.efsa.2021.EN-6922
- ECDC (European Centre for Disease Prevention and Control) and EFSA (European Food Safety Authority), 2022a. Multi-country outbreak of monophasic *Salmonella* Typhimurium sequence type 34 linked to chocolate products—first update—18 May 2022a. EFSA Supporting Publications, 19, 7352E. https://doi.org/10.2903/sp.efsa.2022.EN-7352



- ECDC (European Centre for Disease Prevention and Control) and EFSA (European Food Safety Authority), 2022b. Multi-country outbreak of *Salmonella* Enteritidis sequence type (ST) 11 infections linked to eggs and egg products–8 February 2022b. EFSA Supporting Publications, 19, 7180E. https://doi.org/10.2903/sp.efsa.2022. FN-7180
- ECDC (European Centre for Disease Prevention and Control) and WHO (WHO Regional Office for Europe), 2022. Tuberculosis surveillance and monitoring in Europe 2022 2020 data. Copenhagen: WHO Regional Office for Europe and Stockholm, Licence: CC BY 3.0 IGO. Available online: https://www.ecdc.europa.eu/en/publications-data/tuberculosis-surveillance-and-monitoring-europe-2022-2020-data
- ECDC (European Centre for Disease Prevention and Control), EFSA (European Food Safety Authority) and ANSES (French Agency for Food, Environmental and Occupational Health and Safety), 2021. European Listeria typing exercise (ELiTE). https://doi.org/10.2900/314391 Available online: https://www.ecdc.europa.eu/en/publications-data/joint-ecdc-efsa-and-eurl-Im-report-european-listeria-typing-exercise-elite
- ECDC (European Centre for Disease Prevention and Control), 2022. Congenital toxoplasmosis Annual epidemiological report for 2019. Available online: https://www.ecdc.europa.eu/en/publications-data/congenital-toxoplasmosis-annual-epidemiological-report-2019
- Echevarría JE, Banyard AC, McElhinney LM and Fooks AR, 2019. Current rabies vaccines do not confer protective immunity against divergent Lyssaviruses circulating in Europe. Viruses, 11, 892. https://doi.org/10.3390/v11100892
- EFSA (European Food Safety Authority), 2009. Technical specifications for harmonised national surveys on *Yersinia enterocolitica* in slaughter pigs. EFSA Journal 2009;7(11):1374, 23 pp. https://doi.org/10.2903/j.efsa.2009. 1374. Available online: www.efsa.europa.eu/efsajournal
- EFSA (European Food Safety Authority), 2009a. Statistical analysis of temporal and spatial trends of zoonotic agents in animals and food Part I: Critical review of the statistical analysis carried out on the Community Summary Report 2006 data. EFSA Journal 2009;7(5):253r, 77 pp. https://doi.org/10.2903/j.efsa.2009.253r
- EFSA (European Food Safety Authority), 2009b. Analysis of the baseline survey on the prevalence of *Salmonella* in holdings with breeding pigs in the EU, 2008-Part A: *Salmonella* prevalence estimates. EFSA Journal 2009;7 (12):1377, 93 pp. https://doi.org/10.2903/j.efsa.2009.1377
- EFSA (European Food Safety Authority), 2009c. Technical specifications for the monitoring and reporting of verotoxigenic Escherichia coli (VTEC) on animals and food (VTEC surveys on animals and food). EFSA Journal 2009;7(11):1366, 43 pp. https://doi.org/10.2903/j.efsa.2009.1366
- EFSA (European Food Safety Authority), 2011. Statistical analysis of temporal and spatial trends of zoonotic agents in animals and food Part II: Applications of spatial analysis and further developments of temporal analysis. EFSA Journal 2011;9(8):2331, 72 pp. https://doi.org/10.2903/j.efsa.2011.2331
- EFSA (European Food Safety Authority), 2014. Update of the technical specifications for harmonised reporting of foodborne outbreaks through the European Union reporting system in accordance with Directive 2003/99/EC. EFSA Journal 2014;12(3):3598, 25 pp. https://doi.org/10.2903/j.efsa.2014.3598
- EFSA (European Food Safety Authority), 2022. Catalogues for 2021 zoonoses data reporting. Zenodo. https://doi.org/10.5281/zenodo.6395940
- EFSA (European Food Safety Authority) and ECDC (European Centre for Disease Prevention and Control), 2009. The community summary report on trends and sources of zoonoses and zoonotic agents in the European Union in 2007. EFSA Journal 2009;7(1):223r, 350 pp. https://doi.org/10.2903/j.efsa.2009.223r
- EFSA (European Food Safety Authority) and ECDC (European Centre for Disease Prevention and Control), 2010. The community summary report on trends and sources of zoonoses, zoonotic agents and foodborne outbreaks in the European Union in 2008. EFSA Journal 2010;8:1496, 410 pp. https://doi.org/10.2903/j.efsa.2010.1496
- EFSA (European Food Safety Authority) and ECDC (European Centre for Disease Prevention and Control), 2011. The European Union summary report on trends and sources of zoonoses, zoonotic agents and foodborne outbreaks in 2009. EFSA Journal 2011;9(3):2090, 378 pp. https://doi.org/10.2903/j.efsa.2011.2090
- EFSA (European Food Safety Authority) and ECDC (European Centre for Disease Prevention and Control), 2012. The European Union summary report on trends and sources of zoonoses, zoonotic agents and foodborne outbreaks in 2010. EFSA Journal 2012;10(3):2597, 442 pp. https://doi.org/10.2903/j.efsa.2012.2597
- EFSA (European Food Safety Authority) and ECDC (European Centre for Disease Prevention and Control), 2013. The European Union summary report on trends and sources of zoonoses, zoonotic agents and foodborne outbreaks in 2011. EFSA Journal 2013;11(4):3129, 250 pp. https://doi.org/10.2903/j.efsa.2013.3129
- EFSA (European Food Safety Authority) and ECDC (European Centre for Disease Prevention and Control), 2014. The European Union summary report on trends and sources of zoonoses, zoonotic agents and foodborne outbreaks in 2012. EFSA Journal 2014;12(2):3547, 312 pp. https://doi.org/10.2903/j.efsa.2014.3547
- EFSA (European Food Safety Authority) and ECDC (European Centre for Disease Prevention and Control), 2015a. The European Union summary report on trends and sources of zoonoses, zoonotic agents and foodborne outbreaks in 2014. EFSA Journal 2015;13(12):4329, 190 pp. https://doi.org/10.2903/j.efsa.2015.4329
- EFSA (European Food Safety Authority) and ECDC (European Centre for Disease Prevention and Control), 2015b. The European Union summary report on trends and sources of zoonoses, zoonotic agents and foodborne outbreaks in 2013. EFSA Journal 2015;13(1):3991, 165 pp. https://doi.org/10.2903/j.efsa.2015.3991



- EFSA (European Food Safety Authority) and ECDC (European Centre for Disease Prevention and Control), 2016. The European Union summary report on trends and sources of zoonoses, zoonotic agents and foodborne outbreaks in 2015. EFSA Journal 2016;14(12):4634, 231 pp. https://doi.org/10.2903/j.efsa.2016.4634
- EFSA (European Food Safety Authority) and ECDC (European Centre for Disease Prevention and Control), 2017. The European Union summary report on trends and sources of zoonoses, zoonotic agents and foodborne outbreaks in 2016. EFSA Journal 2017;15(12):5077, 228 pp. https://doi.org/10.2903/j.efsa.2017.5077
- EFSA (European Food Safety Authority) and ECDC (European Centre for Disease Prevention and Control), 2021. The European Union one health 2020 zoonoses report. EFSA Journal 2021;19(12):6971, 324 pp. https://doi.org/10.2903/j.efsa.2021.6971
- EFSA (European Food Safety Authority) and Zancanaro G, 2021. Annual assessment of *Echinococcus multilocularis* surveillance reports submitted in 2020 in the context of Commission Delegated Regulation (EU) 2018/772. EFSA Journal 2021;19(1):6382, 63 pp. https://doi.org/10.2903/j.efsa.2021.6382
- EFSA (European Food Safety Authority), Amore G, Beloeil P-A, Bocca V, Boelaert F, Gibin D, Papanikolaou A, Rizz V and Stoicescu A-V, 2021a. Zoonoses, antimicrobial resistance and foodborne outbreaks guidance for reporting 2020 data. EFSA Supporting Publication 2021:EN-6438, 112 pp. https://doi.org/10.2903/sp.efsa.2021. Available online: www.efsa.europa.eu/efsajournal
- EFSA (European Food Safety Authority), Amore G, Boelaert F, Papanikolaou A, Rizzi V and Stoicescu A-V, 2021b. Manual for reporting on zoonoses and zoonotic agents, within the framework of Directive 2003/99/EC, and on some other pathogenic microbiological agents for information derived from the year 2020. EFSA Supporting Publication 2021:EN-6440, 67 pp. https://doi.org/10.2903/sp.efsa.2021.EN-6440 Available online: www.efsa.europa.eu/efsajournal
- EFSA (European Food Safety Authority), Amore G, Boelaert F, Gibin D, Papanikolaou A, Rizzi V and Stoicescu A-V, 2022. Zoonoses and foodborne outbreaks guidance for reporting 2021 data. EFSA Supporting Publications. EN-7131. https://doi.org/10.2903/sp.efsa.2022.EN-7131
- EFSA (European Food Safety Authority), Alvarez J, Nielsen SS, Robardet E, Stegeman A, Van Gucht S, Vuta V, Antoniou SE, Aznar I and Papanikolaou A, 2022a. Risks related to a possible reduction of the waiting period for dogs after rabies antibody titration to 30 days compared with 90 days of the current EU legislative regime. EFSA Journal 2022;20(6):7350, 78 pp. https://doi.org/10.2903/j.efsa.2022.7350
- EFSA (European Food Safety Authority), Amore G, Boelaert F, Papanikolaou A, Rizzi V and Stoicescu A-V, 2022b. Manual for reporting on zoonoses and zoonotic agents, within the framework of Directive 2003/99/EC, and on some other pathogenic microbiological agents for information derived from the year 2021. EFSA Supporting Publication 2022:EN-7130, 78 pp. https://doi.org/10.2903/sp.efsa.2022.EN-7130
- EFSA (European Food Safety Authority), Boelaert F, Gibin D, Papanikolaou A, Rizzi V and Stoicescu A-V, 2022c. Prevalence sample-based guidance for reporting 2021 data. EFSA Supporting Publications 2022:EN-7129. https://doi.org/10.2903/sp.efsa.2022.EN-7129
- EFSA (European Food Safety Authority), Costa G, Di Piazza G, Koevoets P, Iacono G, Liebana E, Pasinato L, Rizzi V and Rossi M, 2022d. Guidelines for reporting Whole Genome Sequencing-based typing data through the EFSA One Health WGS System. EFSA Supporting Publications 2022:EN-7413. https://doi.org/10.2903/sp.efsa.2022. EN-7413
- EFSA AHAW Panel (EFSA Panel on Animal Health and Welfare), 2014. Statement on a conceptual framework for bovine tuberculosis. EFSA Journal 2014;12(5):3711, 59 pp. https://doi.org/10.2903/j.efsa.2014.3711
- EFSA BIOHAZ Panel (EFSA Panel on Biological Hazards,), EFSA CONTAM Panel (EFSA Panel on Contaminants in the Food Chain) and EFSA AHAW Panel (EFSA Panel on Animal Health and Welfare), 2012. Scientific Opinion on the public health hazards to be covered by inspection of meat (poultry). EFSA Journal 2012;10(6):2741, 179 pp. https://doi.org/10.2903/j.efsa.2012.2741
- EFSA BIOHAZ Panel (EFSA Panel on Biological Hazards), 2007. Scientific Opinion of the Panel on BIOHAZ on a request from EFSA on monitoring and identification of human enteropathogenic Yersinia spp. EFSA Journal 2007;595:1–30, 30 pp. https://doi.org/10.2903/j.efsa.2007.595
- EFSA BIOHAZ Panel (EFSA Panel on Biological Hazards), 2011. Scientific opinion on risk based control of biogenic amine formation in fermented foods. EFSA Journal 2011;9(10):2393, 93 pp. https://doi.org/10.2903/j.efsa. 2011.2393
- EFSA BIOHAZ Panel (EFSA Panel on Biological Hazards), 2013a. Scientific Opinion on the public health hazards to be covered by inspection of meat from farmed game. EFSA Journal 2013;11(6):3264, 181 pp. https://doi.org/10.2903/j.efsa.2013.3264
- EFSA BIOHAZ Panel (EFSA Panel on Biological Hazards), 2013b. Scientific Opinion on the public health hazards to be covered by inspection of meat (solipeds). EFSA Journal 2013;11(6):3263, 161 pp. https://doi.org/10.2903/j.efsa.2013.3263
- EFSA BIOHAZ Panel (EFSA Panel on Biological Hazards), 2013c. Scientific Opinion on VTEC-seropathotype and scientific criteria regarding pathogenicity assessment. EFSA Journal 2013;11(4):3138, 106 pp. https://doi.org/10.2903/j.efsa.2013.3138



- EFSA BIOHAZ Panel (EFSA Panel on Biological Hazards), EFSA CONTAM Panel (EFSA Panel on Contaminants in the Food Chain) and EFSA AHAW Panel (EFSA Panel on Animal Health and Welfare), 2011. Scientific opinion on the public health hazards to be covered by inspection of meat (swine). EFSA Journal 2011;9(10):2351, 198 pp. https://doi.org/10.2903/j.efsa.2011.2351
- EFSA BIOHAZ Panel (EFSA Panel on Biological Hazards), Ricci A, Allende A, Bolton D, Chemaly M, Davies R, Fernández Escámez PS, Girones R, Herman L, Koutsoumanis K, Nørrung B, Robertson L, Ru G, Sanaa M, Simmons M, Skandamis P, Snary E, Speybroeck N, Ter Kuile B, Threlfall J, Wahlström H, Takkinen J, Wagner M, Arcella D, Da Silva Felicio MT, Georgiadis M, Messens W and Lindqvist R, 2018. Scientific Opinion on the Listeria monocytogenes contamination of ready-to-eat foods and the risk for human health in the EU. EFSA Journal 2018;16(1):5134, 173 pp. https://doi.org/10.2903/j.efsa.2018.5134
- EFSA BIOHAZ Panel (EFSA Panel on Biological Hazards), Koutsoumanis K, Allende A, Alvarez-Ordóñez A, Bolton D, Bover-Cid S, Chemaly M, Davies R, De Cesare A, Hilbert F, Lindqvist R, Nauta M, Peixe L, Ru G, Simmons M, Skandamis P, Suffredini E, Jenkins C, Malorny B, Ribeiro Duarte AS, Torpdahl M, da Silva Felício MT, Guerra B, Rossi M and Herman L, 2019. Whole genome sequencing and metagenomics for outbreak investigation, source attribution and risk assessment of foodborne microorganisms. EFSA Journal 2019;17(12):5898, 78 pp. https://doi.org/10.2903/j.efsa.2019.5898
- EFSA BIOHAZ Panel (EFSA Panel on Biological Hazards), Koutsoumanis K, Allende A, Alvarez-Ordóñez A, Bolton D, Bover-Cid S, Davies R, De Cesare A, Herman L, Hilbert F, Lindqvist R, Nauta M, Peixe L, Ru G, Simmons M, Skandamis P, Suffredini E, Alter T, Crotta M, Ellis-Iversen J, Hempen M, Messens W and Chemaly M, 2020a. Update and review of control options for *Campylobacter* in broilers at primary production. EFSA Journal 2020;18(4):6090, 89 pp. https://doi.org/10.2903/j.efsa.2020.6090
- EFSA BIOHAZ Panel (EFSA Panel on Biological Hazards), Koutsoumanis K, Allende A, Alvarez-Ordóñez A, Bover-Cid S, Chemaly M, Davies R, De Cesare A, Herman L, Hilbert F, Lindqvist R, Nauta M, Peixe L, Ru G, Simmons M, Skandamis P, Suffredini E, Jenkins C, Monteiro Pires S, Morabito S, Niskanen T, Scheutz F, da Silva Felício MT, Messens W and Bolton D, 2020b. Pathogenicity assessment of Shiga toxin-producing Escherichia coli (STEC) and the public health risk posed by contamination of food with STEC. EFSA Journal 2020;18(1):5967, 105 pp. https://doi.org/10.2903/j.efsa.2020.5967
- EFSA BIOHAZ Panel (EFSA Panel on Biological Hazards), Koutsoumanis K, Alvarez-Ordóñez A, Bolton D, Bover-Cid S, Chemaly M, Davies R, De Cesare A, Herman L, Hilbert F, Lindqvist R, Nauta M, Peixe L, Ru G, Simmons M, Skandamis P, Suffredini E, Jordan K, Sampers I, Wagner M, Da Silva Felicio MT, Georgiadis M, Messens W, Mosbach-Schulz O and Allende A, 2020c. The public health risk posed by Listeria monocytogenes in frozen fruit and vegetables including herbs, blanched during processing. EFSA Journal 2020;18(4):6092, 102 pp. https://doi.org/10.2903/j.efsa.2020.6092
- Espenhain L, Riess M, Müller L, Colombe S, Ethelberg S, Litrup E, Jernberg C, Kühlmann-Berenzon S, Lindblad M, Hove NK, Torpdahl M and Mörk MJ, 2019. Cross-border outbreak of Yersinia enterocolitica O3 associated with imported fresh spinach, Sweden and Denmark, March 2019. Euro Surveill, 24, pii=1900368. https://doi.org/10.2807/1560-7917.Es.2019.24.24.1900368
- European Parliament, 2020. The EU pig meat sector. Available online: https://www.europarl.europa.eu/RegData/etudes/BRIE/2020/652044/EPRS\_BRI(2020)652044\_EN.pdf
- Eurostat (European Commission), 2020. Ageing Europe looking at the lives of older people in the EU. Statistical Books. https://doi.org/10.2785/628105
- Eurostat (European Commission) 2022. Population structure and ageing. Available online: https://ec.europa.eu/eurostat/statistics-explained/index.php?title=Population\_structure\_and\_ageing [Accessed: 03/11/2022].
- Faber M, Heuner K, Jacob D and Grunow R, 2018. Tularemia in Germany-A Re-emerging Zoonosis. Front Cell Infect Microbiol, 8, 40. https://doi.org/10.3389/fcimb.2018.00040
- FAO (Food and Agriculture Organization of the United Nations) and WHO (World Health Organization), 2018. Shiga Toxin-producing Escherichia Coli (STEC) and Food: Attribution, Characterization and Monitoring. World Health Organization Available online: https://www.fao.org/documents/card/en/c/CA0032EN
- FAO (Food and Agriculture Organization of the United Nations) and WHO (World Health Organization), 2022. Listeria monocytogenes in ready-to-eat (RTE) foods: attribution, characterization and monitoring Meeting report. Microbiological Risk Assessment Series No. 38. Rome. https://doi.org/10.4060/cc2400en
- Farooq Z, Rocklöv J, Wallin J, Abiri N, Sewe MO, Sjödin H and Semenza JC, 2022. Artificial intelligence to predict West Nile virus outbreaks with eco-climatic drivers. Lancet Reg Health Eur, 17, 100370. https://doi.org/10.1016/j.lanepe.2022.100370
- Ferrari RG, Rosario DKA, Cunha-Neto A, Mano SB, Figueiredo EES and Conte-Junior CA, 2019. Worldwide epidemiology of *Salmonella* serovars in animal-based foods: a meta-analysis. Appl Environ Microbiol, 85. https://doi.org/10.1128/aem.00591-19
- Figoni J, Gagnière B, Pouey J, Mailles A, Decors A, Madani N, Revest M, Boisset S and Caspar Y, 2022. Augmentation de l'incidence de la tularémie en France: données de la déclaration obligatoire 2012–2021. Médecine et Maladies Infectieuses Formation, 1, S18–S19. https://doi.org/10.1016/j.mmifmc.2022.03.063
- Fooks AR, Johnson N, Brookes SM, Parsons G and McElhinney LM, 2003. Risk factors associated with travel to rabies endemic countries. J Appl Microbiol, 94(Suppl), 31s–36s. https://doi.org/10.1046/j.1365-2672.94.s1.4.x



- Fornasiero D, Mazzucato M, Barbujani M, Montarsi F, Capelli G and Mulatti P, 2020. Inter-annual variability of the effects of intrinsic and extrinsic drivers affecting West Nile virus vector Culex pipiens population dynamics in northeastern Italy. Parasit Vectors, 13, 271. https://doi.org/10.1186/s13071-020-04143-w
- Forsythe SJ, 2015. Chapter 13 New Insights into the Emergent Bacterial Pathogen Cronobacter. In: SC Ricke, JR Donaldson and CA Phillips (eds). Food Safety. Academic Press, San Diego. pp. 265–308 Available online: https://www.sciencedirect.com/science/article/pii/B9780128002452000137
- García San Miguel Rodríguez-Alarcón L, Fernández-Martínez B, Sierra Moros MJ, Vázquez A, Julián Pachés P, García Villacieros E, Gómez Martín MB, Figuerola Borras J, Lorusso N, Ramos Aceitero JM, Moro E, de Celis A, Oyonarte S, Mahillo B, Romero González LJ, Sánchez-Seco MP, Suárez Rodríguez B, Ameyugo Catalán U, Ruiz Contreras S, Pérez-Olmeda M and Simón Soria F, 2021. Unprecedented increase of West Nile virus neuroinvasive disease, Spain, summer 2020. Euro Surveill, 26. https://doi.org/10.2807/1560-7917.Es.2021.26. 19.2002010
- Genova-Kalou P, Vladimirova N, Stoitsova S, Krumova S, Kurchatova A and Kantardjiev T, 2019. Q fever in Bulgaria: laboratory and epidemiological findings on human cases and outbreaks, 2011 to 2017. Euro Surveill, 24. https://doi.org/10.2807/1560-7917.Es.2019.24.37.1900119
- Gill A, Carrillo C, Hadley M, Kenwell R and Chui L, 2019. Bacteriological analysis of wheat flour associated with an outbreak of Shiga toxin-producing Escherichia coli O121. Food Microbiology, 82, 474–481. https://doi.org/10.1016/j.fm.2019.03.023
- Gormley E and Corner LA, 2013. Control strategies for wildlife tuberculosis in Ireland. Transbound Emerg Dis, 60 (Suppl 1), 128–135. https://doi.org/10.1111/tbed.12095
- Gruber JF, Morris S, Warren KA, Kline KE, Schroeder B, Dettinger L, Husband B, Pollard K, Davis C, Miller J, Weltman A, Mattioli M, Ray L, Tarr C and Longenberger AH, 2021. Yersinia enterocolitica outbreak associated with pasteurized milk. Foodborne Pathog Dis, 18, 448–454. https://doi.org/10.1089/fpd.2020.2924
- Guegan H, Stajner T, Bobic B, Press C, Olariu RT, Olson K, Srbljanovic J, Montoya JG, Djurković-Djaković O and Robert-Gangneux F, 2021. Maternal anti-toxoplasma treatment during pregnancy is associated with reduced sensitivity of diagnostic tests for congenital infection in the neonate. J Clin Microbiol, 59. https://doi.org/10.1128/jcm.01368-20
- Hallanvuo S, Herranen M, Jaakkonen A, Nummela M, Ranta J, Botteldoorn N, De Zutter L, Fredriksson-Ahomaa M, Hertwig S, Johannessen GS, Ludewig M, Messelhäußer U, Sigvart-Mattila P, Thisted-Lambertz S, Thure T and Vatunen E, 2019. Validation of EN ISO method 10273 detection of pathogenic Yersinia enterocolitica in foods. Int J Food Microbiol, 288, 66–74. https://doi.org/10.1016/j.ijfoodmicro.2018.01.009
- Haussig JM, Young JJ, Gossner CM, Mezei E, Bella A, Sirbu A, Pervanidou D, Drakulovic MB and Sudre B, 2018. Early start of the West Nile fever transmission season 2018 in Europe. Euro Surveill, 23, 1800428. https://doi.org/10.2807/1560-7917.Es.2018.23.32.1800428
- Hennebique A, Boisset S and Maurin M, 2019. Tularemia as a waterborne disease: a review. Emerg Microbes Infect, 8, 1027–1042. https://doi.org/10.1080/22221751.2019.1638734
- Hernández-Triana LM, Jeffries CL, Mansfield KL, Carnell G, Fooks AR and Johnson N, 2014. Emergence of west Nile virus lineage 2 in Europe: a review on the introduction and spread of a mosquito-borne disease. Front Public Health, 2, 271. https://doi.org/10.3389/fpubh.2014.00271
- Hestvik G, Warns-Petit E, Smith LA, Fox NJ, Uhlhorn H, Artois M, Hannant D, Hutchings MR, Mattsson R, Yon L and Gavier-Widen D, 2015. The status of tularemia in Europe in a one-health context: a review. Epidemiol Infect, 143, 2137–2160. https://doi.org/10.1017/s0950268814002398
- Hubálek Z and Halouzka J, 1999. West Nile fever--a reemerging mosquito-borne viral disease in Europe. Emerg Infect Dis, 5, 643–650. https://doi.org/10.3201/eid0505.990505
- ISO (International Organization for Standardization), 2001. ISO 16654:2001. Microbiology of food and animal feeding stuffs Horizontal method for the detection of Escherichia coli O157.
- ISO (International Organization for Standardization), 2012. ISO TS 13136:2012. . Microbiology of food and animal feed Real-time polymerase chain reaction (PCR)-based method for the detection of foodborne pathogens Horizontal method for the detection of Shiga toxin-producing Escherichia coli (STEC) and the determination of O157, O111, O26, O103 and O145 serogroups.
- ISO (International Organization for Standardization), 2015. ISO/TS 18867. Microbiology of the Food Chain-Polymerase Chain Reaction (PCR) for the Detection of Food-Borne Pathogens- Detection of Pathogenic *Yersinia enterocolitica* and *Yersinia pseudotubculosis*.
- Kapusta-Duch J, Leszczyńska T, Florkiewicz A and Filipiak-Florkiewicz A, 2011. Comparison of calcium and magnesium contents in cruciferous vegetables grown in areas around steelworks, on organic farms, and those available in retail. Ecol Food Nutr, 50, 155–167. https://doi.org/10.1080/03670244.2011.552373
- Kärssin A, Remes N, Korge K, Viigipuu M, Stensvold CR, Gómez-Morales MA, Ludovisi A, Jokelainen P and Lassen B, 2021. Herbivores as accidental hosts for *Trichinella*: search for evidence of trichinella infection and exposure in free-ranging moose (alces alces) in a highly endemic setting. J Wildl Dis, 57, 116–124. https://doi.org/10.7589/jwd-d-19-00011



- Kastritis E, Tsitsimpis K, Anninos E, Stamatelopoulos K, Kanakakis I, Lampropoulos C, Chatzidou S, Michopoulos S, Papamichail C, Kostis E, Manios E, Kontogiannis S, Paraskevaidis I, Terpos E, Mitrakou A and Dimopoulos MA, 2020. Significant reduction in the visits to the emergency room department during the COVID-19 pandemic in a tertiary hospital in Greece: Indirect victims of the pandemic? Medicine (Baltimore), 99, e23845. https://doi.org/10.1097/md.0000000000023845
- Kefaloudi C, Mellou K, Dougas G, Vorou R, Mitrou K and Kontopidou F, 2022. Human Brucellosis in Greece, 2005–2020: a persistent public health problem. Vector Borne Zoonotic Dis, 22, 163–169. https://doi.org/10.1089/vbz.2021.0050
- Kelly PJ, Matthewman LA, Mason PR and Raoult D, 1993. Q fever in Zimbabwe. A review of the disease and the results of a serosurvey of humans, cattle, goats and dogs. S Afr Med J, 83, 21–25.
- Kenney A, Cusick A, Payne J, Gaughenbaugh A, Renshaw A, Wright J, Seeber R, Barnes R, Florjanczyk A and Horzempa J, 2017. The potential for flower nectar to allow mosquito to mosquito transmission of Francisella tularensis. PLoS One, 12, e0175157. https://doi.org/10.1371/journal.pone.0175157
- Kern P, Bardonnet K, Renner E, Auer H, Pawlowski Z, Ammann RW, Vuitton DA and Kern P, 2003. European echinococcosis registry: human alveolar echinococcosis, Europe, 1982–2000. Emerg Infect Dis, 9, 343–349. https://doi.org/10.3201/eid0903.020341
- Klevar S, Høgåsen HR, Davidson RK, Hamnes IS, Treiberg Berndtsson L and Lund A, 2015. Cross-border transport of rescue dogs may spread rabies in Europe. Vet Rec, 176, 672. https://doi.org/10.1136/vr.102909
- Kuhn KG, Nygård KM, Guzman-Herrador B, Sunde LS, Rimhanen-Finne R, Trönnberg L, Jepsen MR, Ruuhela R, Wong WK and Ethelberg S, 2020. *Campylobacter* infections expected to increase due to climate change in Northern Europe. Scientific Reports, 10, 13874. https://doi.org/10.1038/s41598-020-70593-y
- Kwit NA, Middaugh NA, VinHatton ES, Melman SD, Onischuk L, Aragon AS, Nelson CA, Mead PS and Ettestad PJ, 2020. Francisella tularensis infection in dogs: 88 cases (2014-2016). J Am Vet Med Assoc, 256, 220–225. https://doi.org/10.2460/javma.256.2.220
- Lake IR, Colón-González FJ, Takkinen J, Rossi M, Sudre B, Dias JG, Tavoschi L, Joshi A, Semenza JC and Nichols G, 2019. Exploring *Campylobacter* seasonality across Europe using the European Surveillance System (TESSy), 2008 to 2016. Euro Surveill, 24. https://doi.org/10.2807/1560-7917.Es.2019.24.13.180028
- Lim J, Broughan J, Crowley D, O'Kelly B, Fawsitt R, Burke MC, McCombe G, Lambert JS and Cullen W, 2021. COVID-19's impact on primary care and related mitigation strategies: A scoping review. Eur J Gen Pract, 27, 166–175. https://doi.org/10.1080/13814788.2021.1946681
- Lyoo KS, Kim D, Jang HG, Lee SJ, Park MY and Hahn TW, 2017. Prevalence of antibodies against *Coxiella burnetii* in Korean native cattle, dairy cattle, and dogs in South Korea. Vector Borne Zoonotic Dis, 17, 213–216. https://doi.org/10.1089/vbz.2016.1977
- Maier T, Schwarting A, Mauer D, Ross RS, Martens A, Kliem V, Wahl J, Panning M, Baumgarte S, Müller T, Pfefferle S, Ebel H, Schmidt J, Tenner-Racz K, Racz P, Schmid M, Strüber M, Wolters B, Gotthardt D, Bitz F, Frisch L, Pfeiffer N, Fickenscher H, Sauer P, Rupprecht CE, Roggendorf M, Haverich A, Galle P, Hoyer J and Drosten C, 2010. Management and outcomes after multiple corneal and solid organ transplantations from a donor infected with rabies virus. Clin Infect Dis, 50, 1112–1119. https://doi.org/10.1086/651267
- Marini G, Manica M, Delucchi L, Pugliese A and Rosà R, 2021. Spring temperature shapes West Nile virus transmission in Europe. Acta Trop, 215, 105796. https://doi.org/10.1016/j.actatropica.2020.105796
- Marquer P, Rabade T and Forti R, 2014. Pig farming in the European Union: considerable variations from one Member State to another. Statistics in focus 15/2014.
- Maurin M and Gyuranecz M, 2016. Tularaemia: clinical aspects in Europe. Lancet Infect Dis, 16, 113–124. https://doi.org/10.1016/s1473-3099(15)00355-2
- Migliori GB, Thong PM, Alffenaar JW, Denholm J, Tadolini M, Alyaquobi F, Blanc FX, Buonsenso D, Cho JG, Codecasa LR, Danila E, Duarte R, García-García JM, Gualano G, Rendon A, Silva DR, Souleymane MB, Tham SM, Thomas TA, Tiberi S, Udwadia ZF, Goletti D, Centis R, D'Ambrosio L, Sotgiu G and Ong CWM, 2021. Gauging the impact of the COVID-19 pandemic on tuberculosis services: a global study. Eur Respir J, 58, 2101786. https://doi.org/10.1183/13993003.01786-2021
- Mínguez-González O, Gutiérrez-Martín CB, Martínez-Nistal MDC, Esquivel-García MDR, Gómez-Campillo JI, Collazos-Martínez J, Fernández-Calle LM, Ruiz-Sopeña C, Tamames-Gómez S, Martínez-Martínez S, Caminero-Saldaña C, Hernández M, Rodríguez-Lázaro D and Rodríguez-Ferri EF, 2021. Tularemia outbreaks in Spain from 2007 to 2020 in humans and domestic and wild animals. Pathogens, 10, 892. https://doi.org/10.3390/pathogens10070892
- Mughini-Gras L, van Hoek A, Cuperus T, Dam-Deisz C, van Overbeek W, van den Beld M, Wit B, Rapallini M, Wullings B, Franz E, van der Giessen J, Dierikx C and Opsteegh M, 2021. Prevalence, risk factors and genetic traits of *Salmonella* Infantis in Dutch broiler flocks. Vet Microbiol, 258, 109120. https://doi.org/10.1016/j.vetmic.2021.109120
- NACMCF (National Advisory Committee on Microbiological Criteria for Foods), 2019. Response to Questions Posed by the Food and Drug Administration Regarding Virulence Factors and Attributes that Define Foodborne Shiga Toxin-Producing Escherichia coli (STEC) as Severe Human Pathogens. J Food Prot, 82, 724–767. https://doi.org/10.4315/0362-028x.Jfp-18-479

18314732, 2022, 12, Downloaded from https://efs.aonlinelibrary.wiley.com/doi/10.2903/jefs.a.2022.7666 by Istituto Zopprofilative Sperimenta dell'Umbria e delle Marche (ISSUM), Wiley Online library.wiley.com/rems-and-conditions) on Wiley Online library for rules of use. O. A articles are governed by the applicable Creative Common

- Newell DG, Elvers KT, Dopfer D, Hansson I, Jones P, James S, Gittins J, Stern NJ, Davies R, Connerton I, Pearson D, Salvat G and Allen VM, 2011. Biosecurity-based interventions and strategies to reduce *Campylobacter* spp. on poultry farms. Appl Environ Microbiol, 77, 8605–8614. https://doi.org/10.1128/aem.01090-10
- Newton K, Withenshaw SM, Cawthraw SA and Davies R, 2021. In-depth farm investigations and an exploratory risk factor analysis for the presence of *Salmonella* on broiler farms in Great Britain. Prev Vet Med, 197, 105498. https://doi.org/10.1016/j.prevetmed.2021.105498
- Nikolayevskyy V, Holicka Y, van Soolingen D, van der Werf MJ, Ködmön C, Surkova E, Hillemann D, Groenheit R and Cirillo D, 2021. Impact of the COVID-19 pandemic on tuberculosis laboratory services in Europe. Eur Respir J, 57, 2003890. https://doi.org/10.1183/13993003.03890-2020
- NMKL (Nordisk Metodikkomité for Næringsmidler Nordic Committee on Food Analysis), 2005. Escherichia coli O157. Detection in food and feeding stuffs. NMKL No. 164, 2. Available online: http://www.nmkl.org/index.php?option=com\_zoo&task=item&item\_id=337&Itemid=319&lang=en
- Office fédérel de la santé publique, 2022. Bulletin semaine 4/2022; magazine d'information pour professionnels de la santé et pour les médias. Available online: https://www.bag.admin.ch/dam/bag/fr/dokumente/cc/Kampagnen/Bulletin/2022/bu-4-22.pdf.download.pdf/BU 4 22 FR.pdf
- Oksanen A, Siles-Lucas M, Karamon J, Possenti A, Conraths FJ, Romig T, Wysocki P, Mannocci A, Mipatrini D, La Torre G, Boufana B and Casulli A, 2016. The geographical distribution and prevalence of *Echinococcus multilocularis* in animals in the European Union and adjacent countries: a systematic review and meta-analysis. Parasit Vectors, 9, 519. https://doi.org/10.1186/s13071-016-1746-4
- Olaimat AN and Holley RA, 2012. Factors influencing the microbial safety of fresh produce: a review. Food Microbiol, 32, 1–19. https://doi.org/10.1016/j.fm.2012.04.016
- Olea-Popelka F, Muwonge A, Perera A, Dean AS, Mumford E, Erlacher-Vindel E, Forcella S, Silk BJ, Ditiu L and El Idrissi A, 2017. Zoonotic tuberculosis in human beings caused by *Mycobacterium bovis* a call for action. The Lancet Infectious Diseases, 17, e21–e25. https://doi.org/10.1016/S1473-3099(16)30139-6
- Opsteegh M, Kortbeek TM, Havelaar AH and van der Giessen JW, 2015. Intervention strategies to reduce human Toxoplasma gondii disease burden. Clin Infect Dis, 60, 101–107. https://doi.org/10.1093/cid/ciu721
- Opsteegh M, Spano F, Aubert D, Balea A, Burrells A, Cherchi S, Cornelissen JBWJ, Dam-Deisz C, Guitian J, Györke A, Innes EA, Katzer F, Limon G, Possenti A, Pozio E, Schares G, Villena I, Wisselink HJ and van der Giessen JWB, 2019. The relationship between the presence of antibodies and direct detection of Toxoplasma gondii in slaughtered calves and cattle in four European countries. International Journal for Parasitology, 49, 515–522. https://doi.org/10.1016/j.ijpara.2019.01.005
- Orr B, Malik R, Westman ME and Norris JM, 2022. Seroprevalence of *Coxiella burnetii* in pig-hunting dogs from north Queensland, Australia. Aust Vet J, 100, 230–235. https://doi.org/10.1111/avj.13151
- Ottaiano M, Brunetti R, Calistri P, Di Lorenzo A, Di Sabatino D, Cito F, Chiodini P, Baldi L and De Carlo E, 2022. Risk factors for the persistence and spread of Brucellosis in buffaloes in the province of Caserta (2015–2020). Available online: https://brucellosis2022.izs.it/wp-content/uploads/2022/09/Conference\_Proceedings\_Brucellosis2022.pdf
- Pacenti M, Sinigaglia A, Franchin E, Pagni S, Lavezzo E, Montarsi F, Capelli G and Barzon L, 2020. Human West Nile Virus Lineage 2 infection: epidemiological, clinical, and virological findings. Viruses, 12. https://doi.org/10.3390/v12040458
- Papa A, Xanthopoulou K, Gewehr S and Mourelatos S, 2011. Detection of West Nile virus lineage 2 in mosquitoes during a human outbreak in Greece. Clin Microbiol Infect, 17, 1176–1180. https://doi.org/10.1111/j.1469-0691. 2010.03438.x
- Persad AK and LeJeune JT, 2014. Animal reservoirs of Shiga toxin-producing Escherichia coli. Microbiol Spectr, 2, Ehec-0027-2014. https://doi.org/10.1128/microbiolspec.EHEC-0027-2014
- Pervanidou D, Vakali A, Georgakopoulou T, Panagiotopoulos T, Patsoula E, Koliopoulos G, Politis C, Stamoulis K, Gavana E, Pappa S, Mavrouli M, Emmanouil M, Sourvinos G, Mentis A, Tsakris A, Hadjichristodoulou C, Tsiodras S and Papa A, 2020. West Nile virus in humans, Greece, 2018: the largest seasonal number of cases, 9 years after its emergence in the country. Euro Surveill, 25. https://doi.org/10.2807/1560-7917.Es.2020.25.32.1900543
- Possenti A, Manzano-Román R, Sánchez-Ovejero C, Boufana B, La Torre G, Siles-Lucas M and Casulli A, 2016. Potential risk factors associated with human cystic echinococcosis: systematic review and meta-analysis. PLoS Neglected Tropical Diseases, 10, e0005114. https://doi.org/10.1371/journal.pntd.0005114
- Pozio E, 2014. Searching for *Trichinella*: not all pigs are created equal. Trends Parasitol, 30, 4–11. https://doi.org/10.1016/j.pt.2013.11.001
- Pozio E, 2016. *Trichinella* pseudospiralis an elusive nematode. Veterinary Parasitology, 231, 97–101. https://doi.org/10.1016/j.vetpar.2016.03.021
- Pozio E, Rinaldi L, Marucci G, Musella V, Galati F, Cringoli G, Boireau P and La Rosa G, 2009. Hosts and habitats of *Trichinella* spiralis and *Trichinella* britovi in Europe. Int J Parasitol, 39, 71–79. https://doi.org/10.1016/j.ijpara. 2008.06.006
- ProMED-mail, 2021a. Q fever-Spain (02): (PV) bouldering, Baltzola caves, RFI. 04 May: 20210504.8343274. Available online: https://www.promedmail.org
- ProMED-mail, 2021b. Q fever-Spain: (PV) recurrent, waste treatment facility, RFI. 27 March: 20210327.8273195. Available online: https://www.promedmail.org



- ProMED-mail, 2021c. Q fever-Germany: (Berlin) sheep, human, laboratory staff, RFI. 26 September: 20210926.8698730 Available online: https://www.promedmail.org
- Quereda JJ, Morón-García A, Palacios-Gorba C, Dessaux C, García-Del Portillo F, Pucciarelli MG and Ortega AD, 2021. Pathogenicity and virulence of Listeria monocytogenes: A trip from environmental to medical microbiology. Virulence, 12, 2509–2545. https://doi.org/10.1080/21505594.2021.1975526
- Radoshevich L and Cossart P, 2018. Listeria monocytogenes: towards a complete picture of its physiology and pathogenesis. Nat Rev Microbiol, 16, 32–46. https://doi.org/10.1038/nrmicro.2017.126
- Ray LC, Collins JP, Griffin PM, Shah HJ, Boyle MM, Cieslak PR, Dunn J, Lathrop S, McGuire S, Rissman T, Scallan Walter EJ, Smith K, Tobin-D'Angelo M, Wymore K, Kufel JZ, Wolpert BJ, Tauxe R and Payne DC, 2021. Decreased incidence of infections caused by pathogens transmitted commonly through food during the COVID-19 pandemic foodborne diseases active Surveillance Network, 10 U.S. Sites, 2017-2020. MMWR Morb Mortal Wkly Rep, 70, 1332–1336. https://doi.org/10.15585/mmwr.mm7038a4
- Regnault B, Evrard B, Plu I, Dacheux L, Troadec E, Cozette P, Chrétien D, Duchesne M, Vallat JM, Jamet A, Leruez M, Pérot P, Bourhy H, Eloit M and Seilhean D, 2022. First case of lethal encephalitis in Western Europe due to European Bat Lyssavirus Type 1. Clin Infect Dis, 74, 461–466. https://doi.org/10.1093/cid/ciab443
- Reisen WK, Fang Y and Martinez VM, 2006. Effects of temperature on the transmission of west Nile virus by Culex tarsalis (Diptera: Culicidae). J Med Entomol, 43, 309–317. https://doi.org/10.1603/0022-2585(2006)043[0309: Eotott]2.0.Co;2
- Riccardo F, Monaco F, Bella A, Savini G, Russo F, Cagarelli R, Dottori M, Rizzo C, Venturi G, Di Luca M, Pupella S, Lombardini L, Pezzotti P, Parodi P, Maraglino F, Costa AN, Liumbruno GM and Rezza G, 2018. An early start of West Nile virus seasonal transmission: the added value of One Heath surveillance in detecting early circulation and triggering timely response in Italy, June to July 2018. Euro Surveill, 23. https://doi.org/10.2807/1560-7917.Es.2018.23.32.1800427
- Riccardo F, Bolici F, Fafangel M, Jovanovic V, Socan M, Klepac P, Plavsa D, Vasic M, Bella A, Diana G, Rosi L, Pezzotti P, Andrianou XD, Di Luca M, Venturi G, Maraglino F, Pervanidou D, Cenciarelli O, Baka A, Young J, Bakonyi T, Rezza G and Suk JE, 2020. West Nile virus in Europe: after action reviews of preparedness and response to the 2018 transmission season in Italy, Slovenia. Serbia and Greece. Global Health, 16, 47. https://doi.org/10.1186/s12992-020-00568-1
- Richter J, Esmann L, Lindner AK, Trebesch I, Equihua-Martinez G, Niebank M, Georgi S, Schürmann D, Pfäfflin F, Pasić M and Gertler M, 2019. Cystic echinococcosis in unaccompanied minor refugees from Afghanistan and the Middle East to Germany, July 2016 through June 2017. Eur J Epidemiol, 34, 611–612. https://doi.org/10.1007/s10654-019-00492-8
- Rivas L, Strydom H, Paine S, Wang J and Wright J, 2021. Yersiniosis in New Zealand. Pathogens, 10. https://doi.org/10.3390/pathogens10020191
- Robinson E, de Valk H, Villena I, Le Strat Y and Tourdjman M, 2021. National perinatal survey demonstrates a decreasing seroprevalence of Toxoplasma gondii infection among pregnant women in France, 1995 to 2016: impact for screening policy. Euro Surveill, 26. https://doi.org/10.2807/1560-7917.Es.2021.26.5.1900710
- Rosner BM, Gassowski M, Albrecht S and Stark K, 2021. Investigating the *Campylobacter* enteritis winter peak in Germany, 2018/2019. Scientific Reports, 11, 22902. https://doi.org/10.1038/s41598-021-02423-8
- Rossi P, Tamarozzi F, Galati F, Akhan O, Cretu CM, Vutova K, Siles-Lucas M, Brunetti E and Casulli A, 2020. The European Register of Cystic Echinococcosis, ERCE: state-of-the-art five years after its launch. Parasit Vectors, 13, 236. https://doi.org/10.1186/s13071-020-04101-6
- Rouzic N, Desmier L, Cariou ME, Gay E, Foster JT, Williamson CHD, Schmitt F, Le Henaff M, Le Coz A, Lorléac'h A, Lavigne JP, O'Callaghan D and Keriel A, 2021. First case of Brucellosis caused by an amphibian-type *Brucella*. Clin Infect Dis, 72, e404–e407. https://doi.org/10.1093/cid/ciaa1082
- Sack DA, Sack RB, Nair GB and Siddique AK, 2004. Cholera. Lancet, 363, 223–233. https://doi.org/10.1016/s0140-6736(03)15328-7
- Šamec D, Urlić B and Salopek-Sondi B, 2019. Kale (Brassica oleracea var. acephala) as a superfood: review of the scientific evidence behind the statement. Critical Reviews in Food Science and Nutrition, 59, 2411–2422. https://doi.org/10.1080/10408398.2018.1454400
- Sapountzis P, Segura A, Desvaux M and Forano E, 2020. An overview of the elusive passenger in the gastrointestinal tract of cattle: the Shiga toxin producing Escherichia coli. Microorganisms, 8, 877. https://doi.org/10.3390/microorganisms8060877
- Seiwald S, Simeon A, Hofer E, Weiss G and Bellmann-Weiler R, 2020. Tularemia goes west: epidemiology of an emerging infection in Austria. Microorganisms, 8, 1597. https://doi.org/10.3390/microorganisms8101597
- Sidi-Boumedine K, Rousset E, Henning K, Ziller M, Niemczuck K, Roest H and Thiéry R, 2010. Development of harmonised schemes for the monitoring and reporting of Q-fever in animals in the European Union. EFSA Supporting Publications, 7, 2397–8325, 48E pp. Available online: https://efsa.onlinelibrary.wiley.com/doi/epdf/10. 2903/sp.efsa.2010.EN-48
- Sikkema RS, Schrama M, van den Berg T, Morren J, Munger E, Krol L, van der Beek JG, Blom R, Chestakova I, van der Linden A, Boter M, van Mastrigt T, Molenkamp R, Koenraadt CJ, van den Brand JM, Oude Munnink BB, Koopmans MP and van der Jeugd H, 2020. Detection of West Nile virus in a common whitethroat (Curruca communis) and Culex mosquitoes in the Netherlands, 2020. Euro Surveill, 25. https://doi.org/10.2807/1560-7917.Es.2020.25.40.2001704



- Siles-Lucas M, Casulli A, Conraths FJ and Müller N, 2017. Laboratory diagnosis of *Echinococcus* spp. in human patients and infected animals. Adv Parasitol, 96, 159–257. https://doi.org/10.1016/bs.apar.2016.09.003
- Széll Z, Marucci G, Pozio E and Sréter T, 2013. *Echinococcus multilocularis* and *Trichinella* spiralis in golden jackals (Canis aureus) of Hungary. Veterinary Parasitology, 197, 393–396. https://doi.org/10.1016/j.vetpar.2013.04.032
- Tamarozzi F, Akhan O, Cretu CM, Vutova K, Akinci D, Chipeva R, Ciftci T, Constantin CM, Fabiani M, Golemanov B, Janta D, Mihailescu P, Muhtarov M, Orsten S, Petrutescu M, Pezzotti P, Popa AC, Popa LG, Popa MI, Velev V, Siles-Lucas M, Brunetti E and Casulli A, 2018. Prevalence of abdominal cystic echinococcosis in rural Bulgaria, Romania, and Turkey: a cross-sectional, ultrasound-based, population study from the HERACLES project. Lancet Infect Dis, 18, 769–778. https://doi.org/10.1016/s1473-3099(18)30221-4
- Teunis P, Havelaar A, Vliegenthart J and Roessink G, 1997. Risk assessment of *Campylobacter* species in shellfish: identifying the unknown. Water Science and Technology, 35, 29–34. https://doi.org/10.1016/S0273-1223(97) 00230-8
- Teunis PF, Falkenhorst G, Ang CW, Strid MA, De Valk H, Sadkowska-Todys M, Zota L, Kuusi M, Rota MC, Simonsen JB, Mølbak K, Van Duynhoven YT and Van Pelt W, 2013. *Campylobacter* seroconversion rates in selected countries in the European Union. Epidemiol Infect, 141, 2051–2057. https://doi.org/10.1017/s0950268812002774
- Teunis PF, Sukhrie FH, Vennema H, Bogerman J, Beersma MF and Koopmans MP, 2015. Shedding of norovirus in symptomatic and asymptomatic infections. Epidemiol Infect, 143, 1710–1717. https://doi.org/10.1017/s095026881400274x
- Tozer SJ, Lambert SB, Strong CL, Field HE, Sloots TP and Nissen MD, 2014. Potential animal and environmental sources of Q fever infection for humans in Queensland. Zoonoses Public Health, 61, 105–112. https://doi.org/10.1111/zph.12051
- Van Walle I, Björkman JT, Cormican M, Dallman T, Mossong J, Moura A, Pietzka A, Ruppitsch W and Takkinen J, 2018. Retrospective validation of whole genome sequencing-enhanced surveillance of listeriosis in Europe, 2010 to 2015. Euro Surveill, 23. https://doi.org/10.2807/1560-7917.Es.2018.23.33.1700798
- Verhoeven V, Tsakitzidis G, Philips H and Van Royen P, 2020. Impact of the COVID-19 pandemic on the core functions of primary care: will the cure be worse than the disease? A qualitative interview study in Flemish GPs. BMJ Open, 10, e039674. https://doi.org/10.1136/bmjopen-2020-039674
- WHO (World Health Organization), 2007. WHO Guidelines on tularaemia. Available online: https://apps.who.int/iris/handle/10665/43793
- WHO (World Health Organization), 2020. COVID-19: Considerations for tuberculosis (TB) care. 12, Available online: https://www.who.int/docs/default-source/documents/tuberculosis/infonote-tb-covid-19.pdf
- WHO (World Health Organization), 2022. Third round of the global pulse survey on continuity of essential health services during the COVID-19 pandemic: November–December 2021: Interim Report, 7 February 2022. Available online: file:///C:/Users/ISS/Downloads/WHO-2019-nCoV-EHS-continuity-survey-2022.1-eng.pdf
- WOAH (World Organisation for Animal Health), 2018a. Manual of Diagnostic Tests and Vaccines for Terrestrial Animals. 8th Edition, 2018 Edition. Available online: https://www.woah.org/en/produit/manual-of-diagnostic-tests-and-vaccines-for-terrestrial-animals-2018/
- WOAH (World Organisation for Animal Health), 2018b. Verocytotoxigenic Escherichia coli. In. OIE Terrestrial Manual. Available online: https://www.woah.org/fileadmin/Home/eng/Health\_standards/tahm/3.09.10\_VERO\_E\_COLI.pdf
- WOAH (World Organisation for Animal Health), 2022. Report of the meeting of the scientific commission for animal diseases. Annex 10: report of the development of the case definition for infection with Coxiella burnetii (Q fever). Available online: https://www.woah.org/en/document/report-of-the-meeting-of-the-oie-scientific-commission-for-animal-diseases-scad/
- Young JJ, Haussig JM, Aberle SW, Pervanidou D, Riccardo F, Sekulić N, Bakonyi T and Gossner CM, 2021. Epidemiology of human West Nile virus infections in the European Union and European Union enlargement countries, 2010 to 2018. Euro Surveill, 26, 2001095. https://doi.org/10.2807/1560-7917.Es.2021.26.19.2001095
- Ziegler U, Santos PD, Groschup MH, Hattendorf C, Eiden M, Höper D, Eisermann P, Keller M, Michel F, Klopfleisch R, Müller K, Werner D, Kampen H, Beer M, Frank C, Lachmann R, Tews BA, Wylezich C, Rinder M, Lachmann L, Grünewald T, Szentiks CA, Sieg M, Schmidt-Chanasit J, Cadar D and Lühken R, 2020. West Nile virus epidemic in Germany triggered by epizootic emergence, 2019. Viruses, 12, 448. https://doi.org/10.3390/v12040448

#### **Abbreviations**

ADIS Animal Disease Information System

AE Alveolar echinococcosis

AHAW EFSA Panel on Animal Health and Welfare

BIOHAZ EFSA Panel on Biological Hazards

CA Competent authority
CE Cystic echinococcosis
CFT Complement fixation test
CFU Colony-forming unit

CHC Controlled housing conditions

18314732, 2022, 12, Downloaded from https://efs.aonlinelibrary.wiley.com/doi/10.2903j.d/sa.2022.72666 by Istituto Cooperinents dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18122023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/erens-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Cerawise Common.



dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms

CONTAM EFSA Panel on Contaminants in the Food Chain

COVID-19 Coronavirus disease 2019
DA Direct agglutination

DCF Data Collection Framework

DFS Disease-free-status
DHs Definitive hosts

EBLV European bat lyssavirus

ECDC European Centre for Disease Prevention and Control

EEA European Economic Area
EFTA European Free Trade Association

Eg Echinococcus granulosus

ELISA Enzyme-linked immunosorbent assay

Em Echinococcus multilocularis

ERCE European Register of Cystic Echinococcosis
EU-FORS European Union Foodborne Reporting System
EUOHZ European Union One Health Zoonoses Report

EURL European Union Reference Laboratory
EVD Emerging and vector-borne disease
FAO Food and Agriculture Organization

FBO Foodborne outbreak
FBOp Food business operator

FISH Fluorescent *in situ* hybridisation FNAO Food of non-animal origin

FSC Food safety criteria

FWD Foodborne and waterborne disease

g gram

GAPs Good Agricultural Practices
GMP Good Manufacturing Practices

GP Good Practices

HACCP Hazard analysis and critical control point

HUS Haemolytic-uraemic syndrome

i-ELISA Indirect enzyme-linked immunosorbent assay

IFA Immunofluorescence assay

IFAT Immunofluorescence antibody test

IHC Immunohistochemistry
IHs Intermediate hosts

ISO International Organization for Standardization

JEMRA Joint FAO/WHO Expert Meeting on Risk Assessment

LAT Latex agglutination test
LHT Low heat-treated
MS Member State

MSM Mechanically separated meat
MTBC Mycobacterium tuberculosis complex

N Number

NCP National control programmes
NMKL Nordic Committee on Food Analysis

NRCHC Not raised under controlled housing conditions

NT Not typable

OIE World Organisation for Animal Health

ORV Oral rabies vaccination
PCR Polymerase chain reaction
PHC Process hygiene criteria

RABV Rabies virus

RCHC Raised under controlled housing conditions

ROA Rapid Outbreak Assessments

RTE Ready-to-eat s.l. sensu lato



STEC Shiga toxin-producing *Escherichia coli* 

SoHO Substances of human origin

TB Tuberculosis

TBE Tick-borne encephalitis

TESSy The European Surveillance System

vs. Versus

WGS Whole-genome sequencing WHO World Health Organization

WNF West Nile fever WNV West Nile virus

WNND West Nile neuroinvasive disease
WOAH World Organisation for Animal Health

#### **Country codes**

Albania Austria Belgium Bosnia and Herzegovina Bulgaria Croatia Cyprus Czechia Denmark Estonia Finland	AL AT BE BA BG HR CY CZ DK EE FI
France	FR
Germany	DE
Greece	GR
Hungary	HU
Iceland	IS
Ireland	ΙE
Italy	IT
Latvia	LV
Liechtenstein	LI
Lithuania	LT
Luxembourg	LU
Mantanagra	MT ME
Montenegro Netherlands	NL
Norway	NO
North Macedonia	MK
Poland	PL
Portugal	PT
Romania	RO
Serbia	RS
Slovakia	SK
Slovenia	SI
Spain	ES
Switzerland	CH
Sweden	SE
United Kingdom	GB
United Kingdom (Northern Ireland)	ΧI



1831,4732, 2022, 12, Downloaded from https://ds.aonlinelibrary.wiley.com/doi/10.2903j.ds.a.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Coditiotics (https://onlinelibrary.wiley.com/terms

## Appendix A – Number of tested samples for *Listeria monocytogenes* for the main ready-to-eat (RTE) food categories, by reporting MS and non-MS countries, 2021

Country	RTE milk and milk products	RTE fish and fishery products	RTE meat and meat products		RTE food intended for infants and for medical purposes
Austria	905	176	742	688	1
Belgium	2,501	777	2,459	984	881
Bulgaria	4,518	421	932	4,309	1
Croatia	502	142	994	230	1
Cyprus	388	19	126	479	21
Czechia	2,196	156	6,128	728	2
Denmark	_	424	388	_	_
Estonia	84	227	174	183	_
France	1,427	912	2,367	1,937	18
Germany	8,764	3,991	11,111	9,493	316
Greece	512	102	166	737	6
Hungary	1,266	70	462	2,127	79
Ireland	931	198	1,329	1,678	200
Italy	11,324	967	300	1,629	101
Latvia	60	140	55	20	_
Lithuania	80	6	_	30	_
Luxembourg	53	2	304	621	10
Netherlands	3,520	1,218	633	1,264	107
Poland	8,251	14,972	54,532	15	_
Portugal	402	30	324	1,079	6
Romania	13,960	2,176	16,922	56,491	26
Slovakia	1,749	625	1,997	1,699	460
Slovenia	71	29	90	234	5
Spain	3,169	2,003	4,663	8,162	523
United Kingdom (Northern Ireland)	-	_	-	-	-
EU Total (27 + XI)	66,633	29,783	107,198	94,817	2,764
Albania	75	4	28	14	_
Iceland	_	30	_	_	_
Montenegro	1,381	3	261	_	_
North Macedonia	45	5	60	_	_
Serbia	779	39	424	121	_
Switzerland	3,410	_	_	_	_
Total non-EU countries	5,690	81	773	135	-
Total EU (27+ XI) + non-EU countries	73,323	29,864	107,971	94,952	2,764

<sup>-:</sup> no data available; RTE: ready-to-eat.

For each food category, the number of samples reported in the table was obtained without exclusion criteria.



# Appendix B – Occurrence of *Listeria monocytogenes* at all sampling stages combined in ready-to eat (RTE) food categories using a detection method, EU, 2019–2021

			202	1 <sup>(a)</sup>	20	20	201	9 <sup>(b)</sup>
RTE food category	Food subcategories	Sampling unit	N tested samples	Positive samples (%)	N tested samples	Positive samples (%)	N tested samples	Positive samples (%)
Fish and	Fish	Batch	5,089	5.4	124	1.6	22	0
fishery		Single	2,417	4.4	2,521	4.4	3,775	4.3
products	Fishery products	Batch	196	4.1	113	2.7	25	0
		Single	2,265	3.5	1,625	4.2	2,472	3.7
Milk	Pasteurised	Batch	156	0	153	0	144	0
		Single	1,215	0.08	297	0	813	0.12
	UHT	Batch	_	_	5	0	_	_
		Single	122	0	64	0	99	0
	Raw, intended	Batch	138	1.4	132	0	144	0.69
	for direct human consumption	Single	10	20	336	1.5	44	0
Hard cheeses	From cow milk	Batch	1,836	0.33	2,031	0.3	1,912	0.1
from		Single	1,123	0	805	0	1,273	0
pasteurised	From goat milk	Batch	3	0	8	0	107	0
milk		Single	37	0	42	0	105	0
	From sheep milk	Batch	12	0	20	0	4	0
		Single	29	0	45	6.7	30	0
Hard cheeses	From cow milk	Batch	450	0.67	542	1.1	541	0.55
from raw or		Single	259	0.39	516	0	727	1.1
low heat-	From goat milk	Batch	_		_		_	
treated milk	-	Single	32	0	34	0	29	3.4
	From sheep milk	Batch	_	-	5	0	_	-
		Single	259	4.6	240	5.4	170	3.5
Soft and semi-	From cow milk	Batch	40	0	71	0	172	0
soft cheeses		Single	2,673	0.86	844	1.1	1,465	0.55
from	From goat milk	Batch	18	0	32	0	17	0
pasteurised milk (including	-	Single	316	0.95	61	0	25	0
fresh cheese)	From sheep milk		53	0	23	0	19	0
,		Single	76	0	40	0	45	0
Soft and semi-	From cow milk	Batch	66	0	82	0	130	0
soft cheeses		Single	427	0.70	878	0.46	651	1.4
from raw or	From goat milk	Batch	_	_	_	_	_	_
low heat-		Single	26	0	15	0	42	0
treated milk (including fresh	From sheep milk		_	_	_	_	_	_
cheese)		Single	178	1.7	127	3.1	164	0
Meat products	From bovine	Batch	2	0	12	0	3	0
	animals	Single	2,215	3.9	844	7.5	1,245	4.3
	From broilers	Batch	2	100	11	0	_,	_
		Single	767	1.3	291	0.69	803	2
	From turkeys	Batch	3	0	5	0.05	3	0
	. rom carreys	Single	123	0	157	0.64	125	1.6
	From pigs	Batch	649	0.77	115	0.04	127	4.7

18314732, 2022, 12, Dawnloaded from https://ds.aonlinelibrary.wiley.com/dci/10.2903j.efsa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18122023]. See the Terms and Conditions (https://aninelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA a cardied are governed by the applicable Creative Commons of the conditions of the condi



18314732, 2022, 12, Downloaded from https://efs.aonlinelibrary.wiley.com/doi/10.2903j.efs.a.2022.7666 by Isituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/203]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.2903j.efs.a.2022.7666 by Isituto Zooprofilation Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/203]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.2903j.efs.a.2022.7666 by Isituto Zooprofilation Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/203]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.2903j.efs.a.2022.7666 by Isituto Zooprofilation Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/203]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.2903j.efs.a.2022.7666 by Isituto Zooprofilation Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/203]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.2903j.efs.a.2022.7666 by Isituto Zooprofilation Sperimenta (https://onlinelibrary.wiley.com/doi/10.2903j.efs.a.2022.7666 by Isituto Zooprofilation (https://onlinelibrary

			202	1 <sup>(a)</sup>	20	20	201	9 <sup>(b)</sup>
RTE food category	Food subcategories	Sampling unit	N tested samples	Positive samples (%)	N tested samples	Positive samples (%)	N tested samples	Positive samples (%)
		Single	24,102	2.7	6,470	3	13,908	4.2
Other RTE	Salads <sup>(c)</sup>	Batch	126	0.79	99	1	47	0
products		Single	844	0.95	2,664	2.3	2,993	3.6
	Bakery	Batch	4	0	3	0	22	0
	products <sup>(d)</sup>	Single	1,481	0.2	2,563	0.55	3,202	0.34
	Fruits,	Batch	24	4.2	17	0	66	0
	vegetables and juices <sup>(e)</sup>	Single	1,383	3	1,857	2.9	1,717	2.3
	Sauces &	Batch	4	0	4	0	_	-
	dressings <sup>(f)</sup>	Single	182	0	315	0.32	83	1.2
	Egg products	Batch	2	0	6	0	3	0
		Single	16	0	1	0	23	0
	Confectionery	Batch	_	_	2	0	3	0
	products & pastes <sup>(g)</sup>	Single	26	0	28	0	37	0
	Spices &	Batch	_	_	1	0	2	0
	herbs <sup>(h)</sup>	Single	115	0	113	0	139	1.4
	Other processed	Batch	96	2.1	123	0	252	0
	food products & prepared dishes	Single	21,001	0.34	10,365	0.39	10,585	1.1

- (a): Data from the United Kingdom (Northern Ireland) are taken into account for 2021. In accordance with the agreement on the withdrawal of the United Kingdom from the EU, and in particular with the Protocol on Ireland/Northern Ireland, EU requirements on data sampling are also applicable to Northern Ireland.
- (b): doi:Data from the United Kingdom are taken into account for 2019, because the United Kingdom was an EU MS, but it became a third country on 1 February 2020.
- (c): doi:Includes RTE salads containing mayonnaise.
- (d): doi:Includes bread, cakes, desserts and pastry.
- (e): doi:Includes fruits and vegetables pre-cut, products; fruit or vegetable juice, mixed juice.
- (f): doi:Includes sauce and dressings containing mayonnaise.
- (g): Includes confectionery products and pastes such as chocolate-based product and soft and hard candy.
- (h): Includes spices and herbs, either dried, fresh or frozen.



18314732, 2022, 12. Downloaded from https://drsa.onlinelibrary.wiey.com/doi/10.2903j.drsa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [1812.2023]. See the Terms and Conditions (https://onlinelibrary.wiey.com/erms

## Appendix C – Atlas of STEC serogroups in food, animals and humans in reporting MSs, EU, 2017–2021

			Food	<u> </u>			Aı	nima	ıls			Н	umai	ns	
Year	2021	2020	2019 <sup>(a)</sup>	2018 <sup>(a)</sup>	2017 <sup>(a)</sup>	2021	2020	2019 <sup>(a)</sup>	2018 <sup>(a)</sup>	2017 <sup>(a)</sup>	2021	2020	2019 <sup>(a)</sup>	2018 <sup>(a)</sup>	2017
N cases/sampling units tested	23,659	22,119	25,030	20,498	19,351	3,746		2,588	1,631	2,217	6,084	4,489	7,801	8,167	6,071
N STEC cases/isolates with an identified	239	146	214	161	115	137	92	112	70	115	2,500	1,900	3,718	4,264	3,344
serogroup Serogroup			∟~ ienc				requ							/ (%	 }
01		· cqc		, ( /			· cqc	Circ	( /	,	•	·cqc	iciic,	7 ( 70	
02															
03															
04															
O5															
06						.									
07															
08															
09															
010						-									
011															
012						-									
013															
014															
015															
016															
017															
O18															
O19															
O20															
021															
022															
023															
024															
O25															
O26															
O27															
O28															
O30															
O32															
034															
O36															
037															
O38															
O39															
O40															
041															



			Food	ı				Aı	nima	ls			Н	umai	ns	
Year	2021	2020	2019 <sup>(a)</sup>	2018 <sup>(a)</sup>	2017 <sup>(a)</sup>	100	1707	2020	2019 <sup>(a)</sup>	2018 <sup>(a)</sup>	2017 <sup>(a)</sup>	2021	2020	2019 <sup>(a)</sup>	2018 <sup>(a)</sup>	2017
N cases/sampling units tested	23,659	22,119	25,030	20,498	19,351		2,7	2,112	2,588	1,631	2,217	6,084	4,489	7,801	8,167	6,071
N STEC cases/isolates with an identified												2,500 6	1,900	3,718 7	4,264 8	3,344 6
serogroup	239	146	214	161	115	137		9/	112	2	115					
Serogroup	F	requ	ienc	y (%	<u>)</u>		Fr	requ	enc	<b>y</b> (%	<u>)</u>	F	requ	ency	/ (%	)
042																
O43																
044																
045							_									
O46																
O49																
O50																
O51																
O54																
O55																
O57																
O63																
O65																
O66																
O68																
O69																
O70																
071																
074																
075																
O76																
077																
078																
079																
O80																
O81																
O82																
O83																
084																
O85							T									
O86																
O87																
O88																
O89																
O90																
O91																
O93																
O96																
O98																

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903j.efsa.2022.7666 by Istituto Zooprofilativo Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [1812.2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rate of use; OA articles are governed by the applicable Creative Commons License



			Food				Aı	nima	ıls			Н	umai	ns	
Year	2021	2020	2019 <sup>(a)</sup>	2018(a)	2017 <sup>(a)</sup>	2021	2020	2019 <sup>(a)</sup>	2018 <sup>(a)</sup>	2017 <sup>(a)</sup>	2021	2020	2019 <sup>(a)</sup>	2018 <sup>(a)</sup>	2017
N cases/sampling units tested	23,659	22,119	25,030	20,498	19,351	3,746	2,112	2,588	1,631	2,217	6,084	4,489	7,801	8,167	6,071
N STEC cases/isolates with an identified serogroup	239	146	214	161	115	137	9/	112	70	115	2,500	1,900	3,718	4,264	3,344
Serogroup	F	requ	ency	<b>y</b> (%	)	F	requ	enc	y (%	)	F	requ	ency	/ (%	)
O100															
O101															
O102															
O103															
O104															
O105															
O106															
O107															
O108															
O109															
O110															
0111															
0112															
0113															
0114															
0115															
O116															
0117															
O118															
0119															
0121															
O123															
0124															
0125															
O126															
0127															
0128															
O130															
0132															
0133															
0134															
O136															
0137															
0138															
O139															
O140															
0142															
0143															
O145															

18314732, 2022, 12, Downloaded from https://efsa.onlinelibrary.wiley.com/doi/10.2903j.efsa.2022.7666 by Istituto Zooprofilativo Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [1812.2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rate of use; OA articles are governed by the applicable Creative Commons License



			Food	I			Aı	nima	ls			Н	umai	ns	
Year	2021	2020	2019 <sup>(a)</sup>	2018 <sup>(a)</sup>	2017 <sup>(a)</sup>	2021	2020	2019 <sup>(a)</sup>	2018 <sup>(a)</sup>	2017 <sup>(a)</sup>	2021	2020	2019 <sup>(a)</sup>	2018 <sup>(a)</sup>	2017
N		22,119					7				4	6			
cases/sampling units tested	23,659	2,1	25,030	20,498	19,351	3,746	2,112	2,588	1,631	2,217	6,084	4,489	7,801	8,167	6,071
N STEC	(7	(7	(4			(*)	(7	(7		()	Ψ.	4		- ω	
cases/isolates															
with an											0	0	∞	4	4
identified serogroup	239	146	214	161	115	137	9/	112	2	115	2,500	1,900	3,718	4,264	3,344
Serogroup		requ						ienc					္က iency	۱ (0/V	
O146	•	- Cqu	Circ	, ( ^	,	+ •	· cqu	Circ	, ( /		•	· cqc	iene,	/ ( //	,
0147															
0147															
0149															
0150															
0150															
0152															
0153															
0154															
0155															
0156															
0157															
0159															
O160															
0162															
O163															
0164															
O165															
O166															
0167															
O168															
O169															
O170															
0171															
0172															
0174															
O175															
O176															
O177															
O178															
O179															
O181															
O182															
O183															
O184															
O185															
O186															
O187															
O188															

Note: Presence and absence of STEC serogroups in food, animals and humans. Boxes are coloured according to the proportion of the STEC serogroup in each sample type category and year: red indicates values > 1%; orange, (0.1%–1%); yellow, (0.0001%–0.1%); white, absence of the serogroup.

1831/472, 2022, 12. Downloaded from https://dx.oonlinelibrary.wiley.com/dx/10.2903j.efxa.202.72666 by Isitiuto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18122023]. See the Terms and Conditions (https://aninebhary.wiley.com/rems-and-conditions) on Wiley Online Library for rules of use; OA attacks are governed by the applicable Creative Commons.

<sup>(0.0001%-0.1%);</sup> white, absence of the serogroup.

(a): Data from the United Kingdom are taken into account for 2017-2019, when the United Kingdom was still an EU MS. Since 1 February 2020, the United Kingdom has become a third country.



## Appendix D - STEC serogroups in food and animals, by reporting MS, EU, 2021

		1	l	I							1		1	l		I	l	1	1	l			
Reporting Country	Austria	Belgium	Bulgaria	Croatia	Cyprus	Czechia	Estonia	Finland	France	Germany	Hungary	Ireland	Italy	Latvia	Luxembourg	Netherlands	Poland	Portugal	Romania	Slovakia	Slovenia	Spain	Sweden
N sampling units tested	1,063	3,166	120	145	4	284	7	009	940	5,810	488	1,254	3,422	100	347	4,748	150	314	32	4	275	1,642	2,483
N STEC isolates with an identified serogroup	22	21	0	0		1	o Prop	48	8	8	0	0	22		0	160	0	6	0	0	2	36	
01																							
02																							
O5																							
O6																							
08																							
O9																							
O10																							
O11																							
O15																							
O21																							
O22																							
O26																							
O27																							
O28																							
O32																							
O36																							
O38																							
O46																							
O51																							
O54																							
O65																							
071																							
074																							
O75																							
O76																							
O78																							
O85																							
O88																							
O91																							
O98																							
O100																							
O103																							
O104																							



Reporting Country	j.	шn	aria	tia	SI	nia	nia	pu	e	lany	ary	pu		a	Luxembourg	Netherlands	Þ	ıgal	ania	ıkia	nia		en
	Austria	Belgium	Bulgaria	Croatia	Cyprus	Czechia	Estonia	Finland	France	Germany	Hungary	Ireland	Italy	Latvia	Luxe	Neth	Poland	Portugal	Romania	Slovakia	Slovenia	Spain	Sweden
N samples tested	1,063	3,166	120	145	4	284	7	009	940	5,810	488	1,254	3,422	100	347	4,748	150	314	32	4	275	1,642	2,483
N STEC isolates with an identified serogroup	22	21	0	0	0	1	0	48 (	3	8	0	0	22	0	0	160	0	6	0	0	2	36	4
O108																							
O109																							
0111																							
0112																							
0113																							
0116																							
0117																							
0121																							
O126																							
O128																							
O130																							
O136																							
0145																							
0146																							
0149																							
O150																							
0153																							
0154																							
O155																							
0156																							
0157																							
0163																							
O168																							
0171																							
0174																							
0176																							
0177																							
O178																							
O181																							
O182																							
O183																							
O185																							
O187																							
O188																							

Note: Presence and absence of STEC serogroups in food, animals and humans. Boxes are coloured according to the proportion of the STEC serogroup in each sample type category and year: red indicates values > 1%; orange, (0.1%-1%); yellow,  $(0.0001\%-\le 0.1\%)$ ; white, absence of the serogroup.

1831/472, 2022, 12. Downloaded from https://dx.oonlinelibrary.wiley.com/dx/10.2903j.efxa.202.72666 by Isitiuto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18122023]. See the Terms and Conditions (https://aninebhary.wiley.com/rems-and-conditions) on Wiley Online Library for rules of use; OA attacks are governed by the applicable Creative Commons.



18314732, 2022, 12. Downloaded from https://drs.onlinelibrary.wiley.com/doi/10.29031jd/sa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.29031jd/sa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.29031jd/sa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.29031jd/sa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.29031jd/sa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM). Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.29031jd/sa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM). Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.29031jd/sa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM). Wiley Online Library on [18/12/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/doi/10.29031jd/sa.2022.7666 by Istituto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM).

## Appendix E - Atlas of STEC serogroups in food and animals in reporting MSs, EU, 2021

						Food							Α	nima	ıls	
type/species	Bovine meat	Fruit and vegetable	Milk and dairy products	Mixed meat	Other food	Other meat	Other ruminants meat	Ovine and goat meat	Pig meat	Raw milk	Seeds	Cattle	Small ruminants	Other animals	Other ruminants	Pigs
N samples tested	7,399	4,027	2,999	467	2,419	1,992	231	664	1,407	1,095	959	3,316	48	210	121	51
N STEC cases/isolates with serogroup identified	135	7	9	LO.	ın	11	16	20	_	7	0	129	ıл	7	-	
Serogroup	рі	ropo	rtion	out	of to	tal s	erog	roup	ide	ntifie	ed		otal	rtion sero entif	grou	
01																
02																
O5																
06																
O8																
O9																
O10																
O11																
O15																
O21																
O22																
O26																
O27																
O28																
O32																
O36																
O38																
046																
051																
054																
O65																
071																
074																
075																
076																
078																
O85																
O88																
091																
O98																
O100																
0103																
0104																



						Food								Α.	nima		
						- 000								A	1111112	113	
type/species	Bovine meat	Fruit and vegetable	Milk and dairy products	Mixed meat	Other food	Other meat	Other ruminants meat	Ovine and goat meat	Pig meat	Raw milk	Seeds		Cattle	Small ruminants	Other animals	Other ruminants	Pigs
N samples tested	7,399	4,027	2,999	467	2,419	1,992	231	664	1,407	1,095	959		3,316	48	210	121	21
N STEC cases/isolates with serogroup identified	135	7	9	ю	ю	11	16	20	_	7	0		129	ю	7	-	
Serogroup	рі	ropo	rtion	out	of to	tal s	erog	roup	ide	ntifie	ed			otal		out grou ied	
O108																	
O109																	
O111																	
0112																	
0113																	
O116																	
0117																	
0121																	
O126																	
0128																	
0130																	
0136												-	•				
0145						•						-					
0146																	
O149 O150												-					
0153												-					
0154																	
0155																	
O156												-					
0157																	
0163												-					
O168																	
O171																	
O174																	
O176																	
0177																	
0178																	
0181																	
O182																	
O183																	
O185																	
O187																	
O188																	

Note: Presence and absence of STEC serogroups in food (left) and animals (middle). Boxes are coloured proportionally to the % of the STEC serogroup in the group and year, with red indicating values > 1%; orange, > 0.1% and  $\leq$  1%; yellow, > 0.0001% and  $\leq$  0.1%; white, absence of the serogroup.

1831/472, 2022, 12. Downloaded from https://dx.oonlinelibrary.wiley.com/dx/10.2903j.efxa.202.72666 by Isitiuto Zooprofilatico Sperimenta dell'Umbria e delle Marche (IZSUM), Wiley Online Library on [18122023]. See the Terms and Conditions (https://aninebhary.wiley.com/rems-and-conditions) on Wiley Online Library for rules of use; OA attacks are governed by the applicable Creative Commons.